PRINTED: 12/17/2002

1

ARIZONA DEPARTMENT OF MINES AND MINERAL RESOURCES AZMILS DATA

PRIMARY NAME: CHRISTMAS MINE

ALTERNATE NAMES:

INSPIRATION MS 4443 GILA COPPER SULPHIDE CO. SADDLE MT CO CLAIMS

GILA COUNTY MILS NUMBER: 20B

LOCATION: TOWNSHIP 4 S RANGE 16 E SECTION 30 QUARTER NE LATITUDE: N 33DEG 03MIN 30SEC LONGITUDE: W 110DEG 44MIN 40SEC TOPO MAP NAME: CHRISTMAS - 7.5 MIN

CURRENT STATUS: PAST PRODUCER

COMMODITY:

COPPER SULFIDE IRON MAGNETITE MOLYBDENUM GOLD SILVER ABRASIVE

BIBLIOGRAPHY:

ADMMR CHRISTMAS MINE FILE BLM MINING DISTRICT SHEETS 699 & 700 HARPER C M RECONN IRON RES IN AZ USBM IC 8236 1964 P 29-30 USBM WAR MINERALS REP #291 1944 HICKS C J MOLY OCC IN AZ 1979 P 16 ADMMR PUB PETERSON M P & R W SWANSON GEOL CHRIST MINE ROSS C P USGS BULL 771 1925 P 52

08/19/86

ARIZONA DEPARTMENT OF MINES AND MINERAL RESOURCES FILE DATA

PRIMARY NAME: CHRISTMAS CONCENTRATOR

ALTERNATE NAMES:

GILA COUNTY MILS NUMBER: 19

LOCATION: TOWNSHIP 4 S RANGE 16 E SECTION 20 QUARTER NW LATITUDE: N 33DEG 04MIN 20SEC LONGITUDE: W 110DEG 44MIN 20SEC TOPO MAP NAME: CHRISTMAS - 7.5 MIN

CURRENT STATUS: PRODUCER

COMMODITY:

COPPER SULFIDE SILVER GOLD

BIBLIOGRAPHY:

ADMMR CHRISTMAS MINE FILE EMJ JUNE 74 P 85 PART I, EMJ AUG 74 P 63 PT II (EXCELL EMPL 198 1974 EMJ MINE DIR 1974 (GOOD) EVALUATION IN 0040070052 MSHA YELLOW SHEETS USBM PROD TAB 1973

CHRISTMAS MINE

Mining World, June, 1957, p. 93 11 11 December, 1958, p. 58 11 11 May, 1960, p. 87 11 July, 1961, p. 42 11 11 April, 1962, p. 41 .. 11 July, 1962, p. 41 11 11 September, 1962, p. 28 11 11 February, 1963, p. 31 11 н April, 1963, p. 77

World Mining, July, 1964, p. 66 " " June, 1964, p. 163

Mining Congress Journal, September, 1965, p. 11 """ November, 1966, p. 5

Mining Engineering, December, 1966, p. 13 " April, 1973

Skilling Mining Review, April 14, 1973, p. 12 11 December 22, 1973, p. 19 11 11 11 Aug. 10, 1974, p. 19 (personnel) 11 11 11 Aug. 17, 1974, p. 23 (personnel) 11 Ħ 11 Aug. 24, 1974, p. 4 (safety record) p. 17 (personnel) n 11 11 February 28, 1975, p. 21

Metals Week, February 3, 1975, p. 2 (reduce cu prod.)

Mining Journal, February 7, 1975, p. 104 (prod. cutback)

GILA

CHRISTMAS MINE _ Gila County Location: Sec. 30, T4S, R16E

AEC 172-480, p. 162. In AEC files. No radioactivity.

RI 4293 RI 4559

"New Developments at the Christmas Mine," Ariz. Geological Society Digest, Oct., 1958.

Production Possibilities of the Marginal Copper mines in Ariz., 1941, p. 59-62

USGS Bull. 771 " " 1027-H

Economic Geology, November, 1966, p. 1244, 1245

ABM Bull. 125, p. 41-45 " " 129, p. 32-36, 40-42 " " 180, p. 105, 121, 137, 138, 145, 146, 235

MAPS - Upstairs in for flat file storage in the second drawer.

E/MJ, Vol. 128, No. 4, July 27, 1929, p. 148 11 April, 1959, p. 141 11 (complete story) January, 1963, p. 55-67 11 Vol. 165, No. 6, June, 1964, p. 200 11 Vol. 166, No. 10, October, 1965, p. 114 11 Vol. 167, December, 1966, p. 134 н March, 1967 н Aug., 1974, p. 63, 67, 68 (gen. info.) 11 November, 1974, p. 198 (leaching of near-surface, low-grade ores)

Metal	Mining	&	Processing,	February, 1964, p. 37
11	11		11	July, 1964, p. 45
11	11		11	April, 1965, p. 41

Mining Annual Review 1974, p. 303





18 46

ろうう

1

- (

Sec. 7

Sec. 8

1

		TAIL ING MS No I	TAILING MS No 2	
			5 6	┢
	MS 4728 TAILING MILLSITES	10 9	8 7	-
	Nos 3 Thru 41	13 14	15 16	
	Sec. 17	20 19	18 17	
	23 24 25	26 27	XMAS I XMAS 2.4	
	m 30 29 28	XMAS 3 XMAS 4. 4669 4669	XMAS 5 XMAS 6 4669 4669	
	52 33 34 35	XMAS 10, XMAS 9-4669	XMAS 8 5 XMAS 7	
Sec. 18	37 36 XMAS U XMAS U 4669 -	2 XMAS 13 XMAS 14 4669 4669	XMAS 15 XMAS 16	
	39 40 XMAS 23 XMAS 22 XMAS 2 4669 4669 4669	I XMAS 20 XMAS 19 4669 4669	XMAS 18 XMAS 17 4669 1	
	41 XMAS 24 XMAS 25 XMAS 26 XMAS 2 4669 4669 4669 4669	7 XMAS 28 XMAS 29 4669 4669	9 XMAS 30 XMAS 31 4669 4669	
XMAS 41, XMA 4669, 5, 4'4 68	S 40 XMAS 39 XMAS 38 XMAS 37 XMAS 3 9 4669 4669 4669	6 XMAS 35 XMAS 34 4669 4669 4669 4669 4669	4 XMAS 33 XMAS 32 4669 4669	
XMAS 42 XMAS 43 XM 4669 4 4669 4 4669 4 4669	S 44 XMAS 45 XMAS 46 XMAS 47 XMAS 4 9 4669 4669 4669	8 XMAS 49 XMAS 50 4669 5	2 XMAS 51 XMAS 52	
XMAS 63 XMAS 62 XMA 46634 - 4669 8 465	\$ 61 XMAS 60 XMAS 59 XMAS 58 XMAS 5 9 4669 4669 4669	7 XMAS 56 XMAS 50 4669 5	XMAS 54 XMAS 53 4669 4669	
XMAS 64 XMAS 63 XMA 4669 4659 50 465	\$ 66 XMAS 67 XMAS 68 XMAS 69 XMAS 7 9 4669 4669 4669		4 1 8 8 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1
XMAS 77 XMAS 76 XMAS	5 75 XMAS 74 XMAS 73 XMAS 72 XMAS 7	I B K	NO NO	Ļ
		INSPIRAT 1NSPIRAT 4443 1082 10	INSPIRAT 4443 4443 INSPIRAT	
]

MDS Sheet 699

ARIZONA DEPARTMENT OF MINES AND MINERAL RESOURCES

INFORMATION FROM MINE CARDS IN MUSEUM

GILA COUNTY <u>Christmas Mine</u> (file) MILS#20B 3.AKA' _R	MM 3160 Kinoite with Apophyllite MM 5264 Kinoite on Gangue MM-5316 Dioptase MM-9909 Kinoite with Apophyllite MM M 730 Dioptase		
Υ Υ Υ			

INSPIRATION CONSOLIDATED COPPER COMPANY

OFFICERS

H. Myles Jacob, President H. Carroll Weed, Vice President Edward F. Wendt, Secretary and Treasurer Robert F. Morison, Assistant Secretary and Assistant Treasurer

Operating Staff

At Inspiration, Arizona

H. Carroll Weed, Vice President and General Manager Abner H. Neal, General Superintendent Henry Allen, Jr., Assistant General Superintendent

James R. Watts, Mine Superintendent Bruce B. Whitney, Assistant Mine Superintendent Charles B. Kettering, Plant Superintendent H. Daniel Harper, General Foreman - Leaching Plant Albert J. Turk, General Concentrator Superintendent W. Harold Foard, Smelter Superintendent Howard M. Propper, Mechanical Engineer * Ethbert F. Reed, Chief Geologist Thomas E. Tizard, Chief Engineer - Power Plant William A. Mitchell, Chief Research Engineer Ralph V. Bamerio, Chief Industrial Engineer Robert F. Morison, Assistant Secretary and Assistant Treasurer * retired 5-27-66

At Christmas, Arizona

Richard R. Hyde, General Superintendent Thomas E. Bilson, Mine Superintendent Paul M. Musgrove, Concentrator Superintendent

March 1, 1966

⊹

NEW DEVELOPMENTS AT THE CHRISTMAS MINE

GILA COUNTY, ARIZONA

Ву

JOHN T. EASTLICK

Prepared for presentation at the Annual Meeting of the Arizona Section of the American Institute of Mining, Metallurgical and Petroleum Engineers

December 2, 1957

Not to be reproduced in any form without permission of the Inspiration Consolidated Copper Company.

NEW DEVELOPMENTS AT THE CHRISTMAS MINE, ARIZONA

By

JOHN T. EASTLICK

INTRODUCTION:

The Christmas Mine, located in the Banner Mining District, is in the southwestern corner of Gila County, Arizona, approximately 10 miles to the north of Winkelman along State Highway 77. The mine is situated on the eastern slope of the Dripping Springs Range about one mile to the west and 900 feet above the Gila River. Elevations range from 2100 feet at the Gila River to 4250 feet at the crest of the prominent ridges above the mine.

Discovered in the 1880's the mine has passed through the control and ownership of various mining companies. The isolated location of the property, and the fluctuation of metal prices caused intermittent operations until World War II. According to mine records, production through 1954 was 1,554,500 tons for a total of 54,969,573 pounds of copper.

The ore bodies at Christmas have drawn the interest of many mining companies and several exploration programs were initiated. Of these earlier programs, the exploratory work in the deeper horizons below the 800 level by the United States Bureau of Mines is most significant.

During the present three and one-half year program of development, preparation for a new shaft has been started; the No. 3 shaft has been deepened 534 feet with stations cut at the 1100, 1300, and 1400 levels; 16,800 feet of drifts and raises have been driven; and 73,060 feet of diamond drill holes have been completed.

GEOLOGIC SETTING:

Rocks in the Dripping Springs Range are represented by pre-Cambrian and Paleozoic sediments, intrusive diabase of Cambrian (?) age, extrusive andesite of Cretaceous age, intrusive quartz mica diorite of early Tertiary age, and intrusive basaltic and andesitic dikes of late Tertiary age.

Sediments of the Pre-Cambrian Apache Series outcrop to the northwest of Christmas and are the oldest rocks exposed in the area. Above the Mescal formation of the Apache group are Cambrian sediments composed of the coarse pebbly Troy quartzite with a thickness of approximately 400 feet, succeeded by about 175 feet of fine-grained, argillaceous, and limy quartzites. Uncomformably overlying the Cambrian quartzites are approximately 265 feet of Devonian Martin limestone comprised of shaly and dolomitic limestones in the basal member, massive limestone in the middle horizon, and shales and shaly limestone in the upper member. Above the Devonian Strata are 550 feet of massive Mississippian Escabrosa limestone, followed by approximately 1,000 feet of Naco limestone of Pennsylvanian and Permian (?) age consisting of interbedded shaly limestones, shales, and limestones with local cherty and quartzitic layers.

Intruded into the Mescal limestone and the Troy quartzite are irregular masses and sills of diabase. No exposures of diabase cutting rocks above the Troy have been noted in the area.

Resting on the Paleozoic sediments are a thick series of undifferentiated volcanics consisting of andesitic flows, tuffs, flow breccias, and conglomerates. In the vicinity of Christmas, the volcanics are fine to medium grained andesites consisting essentially of feldspar phenocrysts set in a matrix of fine, dark ferromagnesian minerals.

All the earlier rocks are cut by a generally east-west series of quartz mica diorite dikes. Numerous sills and irregular apophyses extend out into the surrounding rocks. These intrusives occur with several textural variations and are composed primarily of quartz, feldspar, and biotite. Generally, the smaller bodies contain varying amounts of hornblende and biotite. In the Christmas area, the most common quartz mica diorite occurs as a light gray, coarse-grained, even-textured rock becoming porphyritic near the outer margins with large phenocrysts of quartz and feldspar in a fine-grained, dark ground mass. Some narrow quartz mica diorite porphyry dikes cut

-2-

irregularly through the main intrusive masses, probably representing a late phase of the same magma.

In the Christmas mine, narrow post-mineral dikes of basalt and andesite cut through the Paleozoic sediments and the diorite intrusives. They commonly trend from slightly west to east of north with steep irregular dips. Two dikes along the north contact, varying from the general trend, strike approximately N60°E and dip steeply northwest. STRUCTURE:

The Dripping Springs Range is indicated by Ransome (2) to be a complexly faulted, anticlinal structure. The dominant structural features of the region are the generally east-west trending quartz mica diorite dikes, and the series of major northwest-trending faults. The orientation and distribution of the diorite dikes suggest that they were intruded along a system of steep, $N70^{\circ}W - N70^{\circ}E$ trending fissures approximately parallel to the axis of the regional deformation. The major faults are normal faults of the Basin and Range type with their hanging wall or down-thrown sides towards the valleys.

At Christmas one of these major structures, the Christmas fault, separates the area into two geologic settings. To the west, Naco limestone, capped by andesite, forms the prominent outcrops along the steep ridges above the mine, and on the east side andesite comprises the predominent rock eastward to and across the Gila River. The surface outcrops of the quartz mica diorite, intruding the limestones to the west and the volcanics to the east, form an irregular elliptical outline with the long axis trending about N70°E across the Christmas fault zone. Development work underground indicates the intrusive to consist essentially of two thick dikes, converging to the west towards No. 3 Shaft and to the east towards No. 4 Shaft, with numerous branching sills and interfingering smaller dikes. In the footwall of the Christmas fault, the greatest mass of quartz mica diorite is centered to the east of No. 3 Shaft between the 500 and 1100 levels where several thick sills and numerous irregular apophyses

-3-

extend into the adjoining limestones.

The sediments surrounding the intrusive contacts have generally south to southeasterly dips of $10^{\circ} - 20^{\circ}$. Much of the original structure is obscured by later postmineral faulting and by the intrusion of the quartz mica diorite, but it is apparent that some deformation preceded the intrusives. Along the south contact zone, the diorite cuts across the north limb of a small anticlinal fold, and the downward steepening of the sedimentary beds near the north contact suggests that the North dike intruded along the flank of a small flexure. Compressional stresses are also indicated by minor bedding-plane slips, small thrust faults, and local rolls along the bedding.

Pre-mineral fractures are evidenced by the numerous steep-dipping sulphide and quartz stringers in the diorite and in the surrounding rocks. These occur along definite conjugate pre-mineral fracture systems, one set consisting of essentially parallel fractures along the intrusive contacts, and the other set comprising of fractures approximately at right angles to the contacts.

Underground the Christmas fault is exposed on the 300, 400, and 800 levels on the north contact, and at the portal of the 400 level on the south contact. This fault, with a normal displacement of approximately 1200 feet down to the east, strikes generally $N20^{\circ} - 25^{\circ}W$ and dips $65^{\circ}-75^{\circ}$ northeast. Movement occurs along a 10 to 20 foot crushed zone with brecciated blocks and fragments of diorite, andesite, and limestone between several fault strands. Another paralleling structure, the Joker fault, has been discovered on the 800 level, 900 feet to the east of the Christmas fault. It strikes $N10^{\circ} - 25^{\circ}W$ and dips 75° northeast with a normal displacement down to the east of an unknown distance. Numerous smaller faults occur throughout the three major fault blocks. The majority, with normal movements of a few inches to 40 feet, have north to northwesterly strikes, trending obliquely to or parallel to the major structures. In the footwall area of the Christmas fault, two prominent east-west structures form the

-4-

borders of a characteristic graben. On the north side, the 1301 fault strikes generally $N70^{\circ}$ - 75°W and dips 55° - 60° southwest with movement down 60 feet to the south. The No. 3 Shaft fault along the south border strikes approximately N85°W and dips 60° - 70° northeast with a displacement of about 150 feet down to the north.

The age of faulting, particularly that of the major structures, has been the subject of much discussion. There can be no argument that at least part of the displacements along the Christmas and Joker faults are post-mineral, and evidence points strongly, both on a regional and a local scale, to the fact that probably the major part, if not all, the movement was later. Displacements along the 1301 and the No. 3 Shaft faults appear to be post-ore, but may reflect later post-mineral movement along pre-mineral fault or shear zones. The smaller northeasterly to northwesterly striking faults are definitely post-ore in age, cutting and displacing the sedimentary strata, the mineralized beds and stringers, the post-mineral basic dikes, and the diorite intrusive dikes and sills.

ORE DEPOSITS:

The known ore bodies of the Christmas mine are classified as pyro-metasomatic in type, occurring as replacements in metamorphosed limestones of the Naco, Escabrosa, and Martin formations. The relationship of the ore deposits to the intrusive is almost diagrammatic. The type and intensity of mineralization varies with distance from the intrusive contacts, with the degree of metamorphism, with the physical and chemical properties of the sedimentary rocks, and with the intensity of pre-mineral fracturing and shearing.

The sedimentary rocks near the intrusive contacts are highly altered and metamorphosed. In the Naco and Escabrosa limestones, garnet and marble are the principal contact-metamorphic products along with lesser amounts of epidote, wollastonite, idocrase, chlorite, and serpentine. In the Martin limestone, the lower beds are highly altered to serpentine, diopside, tremolite, and chlorite with garnet sparingly

-5-

present. Shale beds like the upper member of the Devonian are altered to fine-grained, banded hornfels and, in some of the thin-bedded limestones, silicification is almost complete. The numerous steep stringers and seams of sulphides which cut the ore deposits attest the fact that much of the metallization took place after development of the lime silicate minerals. Mineralizing solutions, undoubtedly traveling along the pre-mineral fracture zones near the limestone-diorite contacts, formed the extensive replacement ore bodies at intersections with favorable horizons in the metamorphosed limestones.

The principal metallic minerals are chalcopyrite, bornite, magnetite, pyrite, sphalerite, and pyrrhotite. Small amounts of galena and specular hematite are commonly present near the outer margins of the mineralized zones. Some minor molybdenite occurs sparsely in the mineralized beds, generally localized in the siliceous and silicified zones. Magnetite increases with depth, becoming a predominant constituent in ores in the Escabrosa and Martin limestones. Oxidation was almost complete above the 300 level, and extends locally to below the 800 level. Supergene ore minerals include chalcocite, native copper, copper oxides, and copper carbonates.

Naco Limestone:

Production up to 1953 was principally from the middle member of the Naco limestone. The deposits occur as gently dipping, tabular replacements in garnetized zones between thin shale and shaly limestone beds. Peterson and Swanson (1) identify nine distinct ore-bearing horizons along the north contact, and eleven beds on the south side. Viewed on a horizontal projection (Plate I), mineralized bodies in these favorable horizons, yielded ore along the entire lengths of the main intrusive contacts. The individual ore-bearing beds range from 5 to 12 feet in thickness and extend laterally 150 to 200 feet away from the contacts. Oxidation masks much of the character of the primary mineralization, but it is apparent that magnetite and pyrite were predominant against the intrusive contacts, yielding to chalcopyrite and bornite

-6-

in the intermediate zone, and grading to chalcopyrite, pyrite, and sphalerite with minor galena along the outer margins.

Ore bodies in the Naco limestones to the east of the Christmas fault are known only from diamond drill hole intersections and from a few exposures on the 800 level, but they show the same characteristics of the deposits to the west. A 1290 foot diamond drill hole, drilled from the 800 level near the center of the block between the Christmas and Joker faults, cut 955 feet of sediments which definitely can be identified as typical Naco beds.

Escabrosa Limestone:

Ore bodies in the Escabrosa limestone, in contrast to the tabular flat-lying deposits in the Naco limestone, occur as irregular, massive replacements near the quartz mica diorite contacts. Generally, the vertical dimensions are greater than the horizontal thicknesses, the ore bodies terminating abruptly into marbleized limestone away from the contacts. The known ore deposits, separated by a thick diorite sill, are in the upper and lower part of the formation.

Ore bodies developed in the Escabrosa upper horizon on and below the 800 level show the common characteristics of these deposits. Intrusive relationships are generally complex with ore deposition occurring in blocks of limestone completely or almost surrounded by diorite. Other deposits occur as replacements of limestones in embayments between diorite dikes projecting westward from the main intrusives, and as thick, shelllike masses against the outer intrusive borders. The ores are usually higher in grade than those of the Naco with chalcopyrite, bornite, pyrite, magnetite, and sphalerite occurring in a garnet gangue. Magnetite and pyrite commonly predominate near the intrusive contacts with sphalerite localized in the outer margins.

In the lower part of the Escabrosa, ore bodies are known only from diamond drill hole intersections, but they show the same characteristics of deposition peripheral to the intrusive contacts. Other important deposits occur between the main intrusive

-7-

masses.

Experience indicates that wherever the Escabrosa limestones are in contact with the main intrusives, sizeable ore bodies can be anticipated.

Martin Limestone:

The most extensive of the replacement ore bodies is found in the lower part of the Devonian limestones. Mineralization in this horizon, extending north and south from the main intrusive dikes, occurs as a flatly-dipping, massive tabular deposit. These lower limestones, developed to the north on the 1300 level (Plate 3) and to the south on the 1400 level (Plate 4) have proved to be consistently mineralized over an area 2700 feet in width across the intrusives, and 1400 feet in length along the intrusive contacts. Diamond drilling to the east and west of the developed area indicate extensions to 3000 feet along the south contact and to 2000 feet along the north contact. The lower 30 feet of the Martin limestone, consisting of thin-bedded dolomitic and shaly limestones, is the most favorable zone for replacement. However, adjacent to the intrusives, where the intensity of metamorphism and mineralization was greatest, the ore replaces up into the lower part of the massive limestones of the middle member.

Along the south contact, the deposit lies along the south limb of a small anticlinal fold which plunges gently to the west. Mineralization extends throughout a thickness of 65-80 feet for at least 1300 feet along the strike and for 600 to 850 feet down-dip, thinning abruptly to the south and to the west. Adjacent to the north dike, the thicker mineralization extends with a thickness of 55-75 feet for 1600 feet parallel to the intrusive contact, and for 200-400 feet up-dip along the bedding, becoming thinner to the northwest. Between the dikes the more favorable, thin-bedded, impure limestones are replaced, but mineralization above is spotty.

Magnetite forms the predominant metallic mineral throughout the deposit, comprising from 15 to 25 per cent of the total content. Steep-dipping seams of anhydrite and gypsum are common, and local occurrences of fluorite are noted. The sulphide minerals

-8-

commonly show both a vertical and lateral zonal arrangement. Laterally the mineralization grades from a pyrite -- chalcopyrite zone near the intrusive borders, to a chalcopyrite-bornite intermediate zone, and to a pyrrhotite-pyrite-sphalerite-chalcopyrite outer zone. Vertically in the thicker sections, pyrite, chalcopyrite, sphalerite, and sometimes galena generally border a chalcopyrite-bornite central zone.

CONCLUSION:

1. 2. 4

Inspiration Consolidated Copper Company has developed sufficient tonnage of ore to warrant a mine operation. Plans and preparations have been made for a new shaft in conjunction with a 2500 ton-per-day mill. Culmination of these plans is expected if and when the present copper situation improves.

ACKNOWLEDGEMENTS:

Acknowledgements are due the management of the Inspiration Consolidated Copper Company for their cooperation in making this report possible, and to the geological staff at Christmas for their aid in its preparation.

REFERENCES:

- 1. N. P. Peterson and R. W. Swanson, Geology of the Christmas Copper Mine, Gila County, Arizona: U. S. Geol. Survey Bull. 1027-H, 1956.
- 2. F. L. Ransome, <u>Copper Deposits of Ray and Miami, Arizona</u>: U. S. Geol. Survey Prof. Paper 115, 1919.









