



CONTACT INFORMATION

Mining Records Curator
Arizona Geological Survey
1520 West Adams St.
Phoenix, AZ 85007
602-771-1601
<http://www.azgs.az.gov>
inquiries@azgs.az.gov

The following file is part of the

Arizona Department of Mines and Mineral Resources Mining Collection

ACCESS STATEMENT

These digitized collections are accessible for purposes of education and research. We have indicated what we know about copyright and rights of privacy, publicity, or trademark. Due to the nature of archival collections, we are not always able to identify this information. We are eager to hear from any rights owners, so that we may obtain accurate information. Upon request, we will remove material from public view while we address a rights issue.

CONSTRAINTS STATEMENT

The Arizona Geological Survey does not claim to control all rights for all materials in its collection. These rights include, but are not limited to: copyright, privacy rights, and cultural protection rights. The User hereby assumes all responsibility for obtaining any rights to use the material in excess of "fair use."

The Survey makes no intellectual property claims to the products created by individual authors in the manuscript collections, except when the author deeded those rights to the Survey or when those authors were employed by the State of Arizona and created intellectual products as a function of their official duties. The Survey does maintain property rights to the physical and digital representations of the works.

QUALITY STATEMENT

The Arizona Geological Survey is not responsible for the accuracy of the records, information, or opinions that may be contained in the files. The Survey collects, catalogs, and archives data on mineral properties regardless of its views of the veracity or accuracy of those data.

PRINTED: 10-15-2012

ARIZONA DEPARTMENT OF MINES AND MINERAL RESOURCES AZMILS DATA

PRIMARY NAME: PALO VERDE MOUNTAINS

ALTERNATE NAMES:
POGUE PROSPECTS

PINAL COUNTY MILS NUMBER: 803

LOCATION: TOWNSHIP 5 S RANGE 2 E SECTION 15 QUARTER
LATITUDE: N 32DEG 59MIN 36SEC LONGITUDE: W 112DEG 08MIN 43SEC
TOPO MAP NAME: HALEY HILLS - 7.5 MIN

CURRENT STATUS: EXP PROSPECT

COMMODITY:
MICA
COPPER
GOLD

BIBLIOGRAPHY:
ADMMR PALO VERDE MOUNTAINS FILE
CLAIMS EXTEND INTO SEC. 16

POGUE PROSPECTS

PALO VERDE MOUNTAINS, PINAL COUNTY

Roy Pogue reported that he had made a bulldozer cut on the east side of his claims. According to him, the cut reached a maximum of 9 feet in depth and that at about 6 feet of depth, he uncovered limonite and some chalcocite. He said that at 6 feet he generally hit strong iron oxide staining. This checks well with the exposure secured in tunnel farther north, where "chalcocite" or "relief" limonite came in at 6 feet below the surface. Pogue said that he planned to make a similar cut on the west side of the claims. He had been contacted by Woodrow Simmons of Miami and a Phelps Dodge geologist for later visits.

Memo - LEWIS A. SMITH - 4-18-63 - Interview with Roy Pogue.

DEPARTMENT OF MINERAL RESOURCES

STATE OF ARIZONA
FIELD ENGINEERS REPORT

Mine Pogue Prospects
(1) Palo Verde Mountains, Pinal Co.
District (2) Sierra Estrella " " "

Date March 20, 1963

Engineer Lewis A. Smith

Subject: Mine visit with Roy C. Pogue.

Property: 5 unpatented lode claims.

Location: (1) Sections 15-16, T. 5 S., R. 2 E. (Palo Verde)
(2) " 6-7, T. 4 S., R. 2 E. (Sierra Estrella)

Owner: Roy C. Pogue, Box 24, Tolleson, Arizona (WE 6-3869).

Access: The Palo Verde mountains prospects were reached by driving on a graded road for 8 miles, parallel to the Southern Pacific R.R., west from Maricopa, thence due south for 6 miles on the Hidden Valley country road. The Sierra Estrella occurrence lies 1 mile north of the railroad and 9 miles west of Maricopa.

Minerals: Mica and copper, with minor gold.

Work: On the southern tip of the Palo Verde Mountains, and 100 to 125 feet above Hidden Valley, a series of trenches were dug. These extend over a length of 100 feet and are 1-5 feet deep and up to 4 feet wide. Over the ridge, 1/4 - 3/8 of a mile, two shallow shafts and a 15 foot adit drift were sunk. The deepest shaft is about 12 feet. The two shafts were on mica occurrences.

Geology: The Palo Verde mountains are essentially composed of fine grained light gray colored granite gneiss, which is overlain by schist "floaters" in numerous small masses, each of which has very limited horizontal area. The gneiss had intruded the schist in a series of ribs disrupting it by tilting, doming, and shattering. The gneiss at the contacts is pegmatitic, sometimes coarsely, but usually the pegmatites are relatively fine grained. The core of the gneiss mass is fine grained, closely foliated, and hard. The foliation lines are marked by fine grained quartz, biotite and orthoclase feldspar, giving the rock a black and light gray ribboned appearance. The coarser border pegmatitic zones contain muscovite-mica books up to 1-2 inches in diameter. These books were mined in places on the northeastern part of the claims. Much of the mica, however, is smaller in size, ranging from 1/8 to 1/2 inches and none appeared suitable for sheets. The pegmatites are roughly zoned, the central zone being composed of variously shaped dense white quartz masses, followed outward by feldspar and mica that are often inter-crystallized. Tongues of pegmatite and fine grained gneiss extend out from the mass of fine grained gneiss into the schist laminae for short distances, some of these tongues are up to 1 foot thick. The schists are of two major types - hornblende schist and quartz-mica schist. The predominant trend of the gneissic intrusives is nearly north, but locally they pinch and swell and vary in trend, and have been offset by the shears and faults. The schist now has no definite trend, its blocks being severely jumbled. The schist-gneiss contacts, in places, were mineralized by chrysocolla, garnet and epidote, particularly where E-W or NE-SW shear fractures cut them. The shears and transverse faults were also mineralized to some extent. In some cases, notably one on the east slope of the mountains, "chalcocite limonite" or "relief limonite"

Pogue Prospects (continued)

was found in a two foot gouge zone, along the 15 foot adit. This zone strikes NE-SW and dips 70 degrees N and reached 6 feet below the surface. Weathering has been intense in some places, whipping the copper oxides and limonite completely out for as much as 4-8 feet below the surface. Because of this, it was difficult to evaluate the area, no limonite being evident on or near the surface. The rocks, however, are mostly chemically neutral and would, except very locally, permit free downward and nearly vertical movement of leaching solutions. While the occurrences of copper as chrysocolla, are sporadic and local, the chance that the bulk of the copper may have been leached downward, is present. Superficial examination, such as this was, coupled with the deep outcrop weathering, makes it nearly impossible to conclude anything definite as to the worth of the prospect. Nevertheless the rocks' chemical neutrality, the presence of good limonite at some distance below the surface, coupled with an intimate structural pattern suggests that drilling, or geophysical tests, are in order. These copper showings are fairly sporadically distributed over an area 1/2 mile wide and 3/4 of a mile in length, but appear to be more evident in the zones affected by shears or faults. The maturity of the oxidized capping, where seen in the few workings, would indicate that sulphides, where present, would probably be well below the surface. The range topography in most of the area is precipitous and low angle holes could attain good depth rapidly. On the contrary, however, the rocks are intensely shattered locally, a fact that could cause much difficulty in coring. The drill sites could be had cheaply since the slopes up to the base of the ridge are relatively smooth. The ridge rises very abruptly in places. The topography is to a considerable extent regulated by the transverse faults and shears, most of the gullies having developed along these structures, serrating the range crest. Much of the north 5 miles at the Palo Verde mountain does not appear to be severely broken.

The Sierra Estrella mountains area is quite similar geologically to that of the Palo Verde mountains so will not be described separately. However, the topography here, is much less precipitous.