U.V.X. M-3 Zone
Vertical X-Sec. No. 2

Looking N45°W

1" = 20'

Don White - June, 1987

SW

NE

600

700

800

903

950

11,300N 6,900E

11,400N 7,000E

11,500N 7,100E

11,600N 7,200E

Tcgl

"Red bleaky schist"

Stope

$CuCO_3$ ore

Tertiary conglomerate

dt

"Schist"

Siliceous rock ("chert")

Primary silica sinter/exhalite &/or silica alt'n

si

Very ferruginous

Manway & ore pass

826, 827, 828 Stopes

828-2 Stope (1937)

Higher grade

Low grade ± Fe breccia

809 D.D.S. ~335' rear

806 D.D.S.

"Red bleaky schist" 800 Sill

824-N

10th Floor

Ore pass

X-Cut ~5' forward (Feb. 1936)

809-3 ~E.O.H. only ~5' rear in workings (365'-370') High grade (.37 oz Au/9.3 oz Ag) high Fe si br + gr-it

High grade Au+Ag in old low grade in drill (location problem?) + shear to grit zone (?)

Fe+

M-4 ~120'

Approx. penetration by D.D.H. 806-1, ~590', in low grade hem. si. br.

.5 oz Au/3.5 oz Ag = (1935)

903 Sill

903-N Drift

928-1 Stope (Mined Sept. 1935)

High grade Au+Ag in fault grit

Limonitic bxt. low Fe silica grit with high grade Au+Ag

Low grade Au+Ag ± Fe

Banded + massive chert (recrystallized) as found in drill holes using MAP casing of Morgan DDS location

M-9 ~125'

Limonitic silica grit with low grade Au+Ag

M-6 ~90'

M-3 ~82'

M-5 ~78'

Argillic-altered diorite

Chert-diorite contact as found in drill holes

Banded chert

Location Problem

Chert-limonitic contact with largest Au+Ag assays

May 1987

Morgan X-Cut

Morgan D.D.S. ~60' rear

Diorite within Verde Fault zone

Hanging Wall Verde Fault

Footwall Verde Fault

Cleopatra quartz porphyry member of Deception Rhyolite Fm.

qp

907 Drift