



CONTACT INFORMATION
Mining Records Curator
Arizona Geological Survey
416 W. Congress St., Suite 100
Tucson, Arizona 85701
602-771-1601
<http://www.azgs.az.gov>
inquiries@azgs.az.gov

The following file is part of the Grover Heinrichs Mining Collection

ACCESS STATEMENT

These digitized collections are accessible for purposes of education and research. We have indicated what we know about copyright and rights of privacy, publicity, or trademark. Due to the nature of archival collections, we are not always able to identify this information. We are eager to hear from any rights owners, so that we may obtain accurate information. Upon request, we will remove material from public view while we address a rights issue.

CONSTRAINTS STATEMENT

The Arizona Geological Survey does not claim to control all rights for all materials in its collection. These rights include, but are not limited to: copyright, privacy rights, and cultural protection rights. The User hereby assumes all responsibility for obtaining any rights to use the material in excess of "fair use."

The Survey makes no intellectual property claims to the products created by individual authors in the manuscript collections, except when the author deeded those rights to the Survey or when those authors were employed by the State of Arizona and created intellectual products as a function of their official duties. The Survey does maintain property rights to the physical and digital representations of the works.


QUALITY STATEMENT

The Arizona Geological Survey is not responsible for the accuracy of the records, information, or opinions that may be contained in the files. The Survey collects, catalogs, and archives data on mineral properties regardless of its views of the veracity or accuracy of those data.

STATE OF CALIFORNIA
EDMUND G. BROWN, *Governor*
THE RESOURCES AGENCY
HUGO FISHER, *Administrator*
DEPARTMENT OF CONSERVATION
DE WITT NELSON, *Director*

GEOLOGIC MAP OF CALIFORNIA
WEED SHEET

Scale 1:250,000
1964




DIVISION OF MINES AND GEOLOGY
IAN CAMPBELL, *State Geologist*
Ferry Building, San Francisco

Price \$1.50

INDEX TO CALIFORNIA GEOLOGIC MAP SHEETS


The date beneath each map sheet name indicates year of publication. If no date appears, the geologic map sheet had not been published at the time this cover was issued.



DIVISION OF MINES AND GEOLOGY
IAN CAMPBELL, CHIEF
GORDON B. OAKESHOTT, DEPUTY CHIEF


STATE OF CALIFORNIA
THE RESOURCES AGENCY
DEPARTMENT OF CONSERVATION

GEOLOGIC MAP OF CALIFORNIA
WEED SHEET



SEDIMENTARY AND METASEDIMENTARY ROCKS		IGNEOUS AND META-IGNEOUS ROCKS	
Qs	Dune sand	Qv	Recent volcanic: Qv - rhyolite; Qv - andesite; Qv - basalt; Qv - pyroclastic rocks
Al	Alluvium		
Sc	Stream channel deposits		
Fd	Fan deposits		
Bd	Basin deposits		
Sl	Salt deposits		
Ql	Quaternary lake deposits		
Qm	Quaternary marine and marine terrace deposits		
Qn	Quaternary nonmarine terrace deposits		
Qp	Quaternary nonmarine		
Pt	Plio-Pliocene nonmarine		
U	Undivided Pliocene nonmarine		
Uu	Upper Pliocene nonmarine		
Um	Upper Pliocene marine		
Ml	Middle and/or lower Pliocene nonmarine		
Mm	Middle and/or lower Pliocene marine		
Uo	Undivided Miocene nonmarine		
Uo	Upper Miocene nonmarine		
Uo	Upper Miocene marine		
Mo	Middle Miocene nonmarine		
Mo	Middle Miocene marine		
Lo	Lower Miocene marine		
Oo	Oligocene nonmarine		
Oo	Oligocene marine		
EO	Eocene nonmarine		
EO	Eocene marine		
PO	Paleocene nonmarine		
PO	Paleocene marine		
CO	Cenozoic nonmarine		
CO	Cenozoic marine		
Tn	Tertiary nonmarine		
Tn	Tertiary lake deposits		
Tn	Tertiary marine		
UO	Undivided Oligocene marine		
UO	Upper Oligocene marine		
LO	Lower Oligocene marine		
JO	Upper Jurassic marine		
JO	Middle and/or Lower Jurassic marine		
JO	Triassic marine		
PC	Pre-Cretaceous metamorphic rocks (in - limestone or dolomite)		
PC	Pre-Cretaceous metamorphic rocks		
PC	Pre-Cretaceous granitic and metamorphic rocks		
PC	Pre-Cretaceous marine (in - limestone or dolomite)		
PC	Paleozoic metamorphic rocks		
PC	Permian marine		
PC	Permian metamorphic rocks		
PC	Undivided Carboniferous marine		
PC	Carboniferous metamorphic rocks		
PC	Pennsylvanian marine		
PC	Mississippian marine		
PC	Devonian marine		
PC	Devonian metamorphic rocks		
PC	Silurian marine		
PC	Devonian and pre-Devonian metamorphic rocks		
PC	Pre-Silurian metamorphic rocks		
PC	Pre-Silurian metamorphic rocks		
PC	Ordovician marine		
PC	Ordovician metamorphic rocks		
PC	Cambrian marine		
PC	Cambrian - Pre-cambrian metamorphic rocks		
PC	Undivided Precambrian metamorphic rocks		
PC	Later Precambrian sedimentary and metamorphic rocks		
PC	Earlier Precambrian metamorphic rocks		

TOPOGRAPHIC BASE MAP
Prepared by the Army Map Service (AGS), Corps of Engineers, U.S. Army, Washington, D.C. Contours from 1957 from United States Quadrangle, 1:48,000; 1:50,000; 1:62,500; U.S. Geological Survey and AMS, 1953-54. Topographic maps, photogrammetric, reliability good. Control by USGS, USGS and USFS. Map field checked, 1958.
Line not shown by U.S. Geological Survey.
Submarine contours derived from Shepard and Emery Special Paper No. 31, Dept. Soc. America.
Major corrections and additions to contour by California Division of Mines and Geology, 1963.



CONTOUR INTERVAL 200 FEET
WITH SUPPLEMENTARY CONTOURS AT 100 FOOT INTERVALS
GEOLOGIC MAP OF CALIFORNIA
OLAF P. JENKINS EDITION
WEED SHEET
COMPILATION BY RUDOLPH G. STRAND 1963



INDEX TO GEOLOGIC MAPPING
COMPLETE INDEX ON EXPLANATORY DATA SHEET
1. ...
2. ...
3. ...
4. ...
5. ...
6. ...
7. ...
8. ...
9. ...
10. ...
11. ...
12. ...
13. ...
14. ...
15. ...
16. ...
17. ...
18. ...
19. ...
20. ...
21. ...
22. ...
23. ...
24. ...
25. ...
26. ...
27. ...
28. ...
29. ...
30. ...

HEAVY BORDER ON BOXES INDICATES UNITS THAT APPEAR ON THIS SHEET

STRATIGRAPHIC NOMENCLATURE— WEED SHEET

AGE	STATE MAP SYMBOL	STATE MAP UNIT <small>State Map Units listed here are not necessarily in stratigraphic sequence; the sequence used has been standardized for all sheets of the Geologic Map of California</small>	STRATIGRAPHIC UNITS AND CHARACTERISTIC LITHOLOGIES <small>(The formally named formations grouped within an individual State Map Unit are listed in stratigraphic sequence from youngest to oldest.)</small>
QUATERNARY	Qs	RECENT DUNE SAND	Dune sand and older eolian sand along the coast.
	Qal	RECENT ALLUVIUM	Recent stream channel, flood plain and alluvial fan deposits. Fluvio-glacial deposits (may be Pleistocene in part), and mud flow deposits, notably along Mud Creek on southeast flank of Mt. Shasta.
	Qrv	RECENT VOLCANIC ROCKS: UNDIFFERENTIATED	"Shastina" pyroxene andesite—dark gray or black andesitic basalt rich in bright green granules of pyroxene (these lavas are "identical" with the early flows of Shasta and hence it is not certain that they originated from the Shastina vent, Williams, 1934, p. 245).
	Qrv ^a	ANDESITIC	Hypersthene-rich vesicular andesite and basaltic andesite of Goozenest volcano. Dark vesicular, glass-rich pyroxene andesite of Deer Mountain. Pale gray porphyritic pyroxene andesite lava, the latest flow from Shasta. Pale gray andesite lavas (containing long slender prisms of brownish hornblende) of Black Butte dome. Hornblende andesite talus breccia in Diller Canyon on west side of Mt. Shasta.
	Qrv ^b	BASALTIC	Pluto's Cave Basalt—black, vesicular olivine-rich augite basalt. Olivine-augite basalt at Copco Lake. Blocky, scoriaceous, augite-olivine basalt along Alder Creek. Black, vesicular, olivine-augite basalt of Little Deer Mtn. volcano. Dark, vesicular, olivine basalt of The Whaleback. Gray and black, glassy, pyroxene-rich andesitic basalt of Shastina.
	Qrv ^p	PYROCLASTIC	Dark tuff, lapilli, and bombs of vesicular glass, on Mt. Shasta. Red, brown and black basaltic cinders in the Macdoel 30' quadrangle.
	Ql	QUATERNARY LAKE DEPOSITS	Recent diatomite at Copco Lake. Semiconsolidated clay, volcanic ash, diatomite, and sand in Butte Valley. Glacial lake sediments near the town of Mt. Shasta.
	Qg	QUATERNARY GLACIAL DEPOSITS	Late (Morris Meadow), middle (Rush Creek), and early (Alpine Lake) Wisconsin (age) moraines in the Trinity Alps area. Undifferentiated glacial deposits on Mt. Shasta and elsewhere. Glacial deposits shown on this map are generalized; in some places the areas are exaggerated in size and may include less than 10% glacial debris.
	Qt	QUATERNARY NONMARINE TERRACE DEPOSITS	Unconsolidated sand, clay, and gravel terrace deposits.
	Qm	PLEISTOCENE MARINE DEPOSITS AND MARINE TERRACE DEPOSITS	Battery Formation—buff and blue sand units interbedded with white, bluish-gray, brown, and blue clay units.
PLEISTOCENE	Qc	PLEISTOCENE NONMARINE SEDIMENTARY DEPOSITS	"Older" alluvium, fan and terrace deposits. "Older" surficial agate-bearing deposits northwest of Copco Dam. "Older" dune sand in the Trinidad quadrangle.
	Qpv	PLEISTOCENE VOLCANIC ROCKS: UNDIFFERENTIATED	Andesitic basalt and dense gray pyroxene andesite (early flows forming the base of Mt. Shasta).
	Qpv ^r	RHYOLITIC	Pale gray rhyolitic dacite characterized by large phenocrysts of plagioclase. White pumiceous lava (probably a glassy dacite) forming the domes south of Gray Butte on Mt. Shasta.
	Qpv ^a	ANDESITIC	Porphyritic pale-gray and brown, pyroxene andesites having distinctive platy or slabby habit and abundant basic inclusions, present on Mt. Shasta. Fine-grained andesite or andesitic basalt of the Bear Butte and Spring Hill lava cones. Pale gray andesite and dacite of Gray Butte plug dome. Colorless glass-rich andesite of the twin domes on the north side of Mt. Shasta.
	Qpv ^b	BASALTIC	Butte Valley Basalt—smooth crusted, black to gray, vesicular olivine basalt (in part Recent). Pale gray to black, dense, massive to vesicular basalts that emanated from the Everett Hill volcano and flowed down the Sacramento River Valley.
	Qpv ^p	PYROCLASTIC	Well consolidated lapilli tuff and tuff-breccia in the Macdoel 30' quadrangle, (Pleistocene to Recent). Volcanic mud flow deposits and tuff-breccias exposed in walls of Mud Creek Canyon (perhaps oldest exposed rocks of Mt. Shasta). Black tuff, lapilli, and volcanic bombs on Mt. Shasta.
	QP	PLIOCENE-PLEISTOCENE NONMARINE SEDIMENTARY DEPOSITS	Probable Plio-Pleistocene deposits in the Orick quadrangle. Pleistocene or Late Tertiary terrace deposits that have been slightly deformed and contain deeply weathered boulders (Hoopa quadrangle).
	☼	QUATERNARY AND/OR PLIOCENE CINDER CONES	Pliocene (?) to Recent cinder cones.
	Pu	UPPER PLIOCENE MARINE SEDIMENTARY ROCKS	Small areas of highly fossiliferous marine, near-shore sands and gravels deposited on very irregular surface along the present day coast.
	PLIOCENE	Pv ^a	ANDESITIC
Pv ^b		BASALTIC	Olivine basalt and basaltic andesite in the Macdoel 30' quadrangle (possibly Plio-Pleistocene to late Pleistocene).
Pv ^p		PYROCLASTIC	Low mounds of red cinders, remnants of cinder cones in Macdoel quadrangle.
Tm		TERTIARY MARINE SEDIMENTARY ROCKS	St. George Formation—dull gray-blue, massive fossiliferous siltstone and shale containing irregular lenses of sand (Pliocene).
TERTIARY	Mu	UPPER MIOCENE MARINE SEDIMENTARY ROCKS	Wimer Formation—friable yellow shale and siltstone that weathers brown.
	Tc	TERTIARY NONMARINE SEDIMENTARY ROCKS	Late Tertiary auriferous gravel deposits in the Gasquet and Klamath quadrangles.
	Ti	TERTIARY INTRUSIVE (HYPABYSSAL) ROCKS: UNDIFFERENTIATED	Intrusive rocks exposed in Hornbrook and Dunsmuir areas.
	Ti ^r	RHYOLITIC	Massive gray and olive-green porphyritic pyroxene dacite at Cedar Lake. Albite rhyolite porphyry dikes in the Gasquet quadrangle.
	Ti ^a	ANDESITIC	Andesitic volcanic necks.
	Ti ^b	BASALTIC	Basaltic volcanic neck in vicinity of Copco Dam.
	Tv	TERTIARY VOLCANIC ROCKS: UNDIFFERENTIATED	Wason Formation, Roxy Formation, Colestine Formation ¹ —hypersthene augite andesite, basalt, some dacite flows, volcanic conglomerate and sandstone (considered to be upper Eocene to upper Miocene by H. Williams).
	Tv ^r	RHYOLITIC	Basaltic flows in the Gazelle area and in the southeast corner of the map area (age uncertain; may prove to be Quaternary).
	Tv ^b	BASALTIC	White to pale cream very fine-grained rhyolite plug domes in the Macdoel 30' quadrangle.
	Tv ^p	PYROCLASTIC	White to cream, and pale blue-green rhyolite tuff, glassy pumiceous dacite tuff, and andesitic tuff-breccia.
CRETACEOUS	Ku	UPPER CRETACEOUS MARINE SEDIMENTARY ROCKS	Hornbrook Formation—greenish-gray massive arkosic sandstone, conglomerate, and bluish-gray shale, buff and white massive sandstone (nonmarine in part), cross-bedded sandstone, and coal. Coarse- to medium-grained sandstone, angular boulder conglomerate, and light-gray shale in the Sunlight Peak area.
	Kjf	FRANCISCAN FORMATION	Franciscan Formation—graywacke, interbedded shale, minor conglomerate, thin-bedded chert, some undifferentiated basaltic or siltitic rocks that have been altered to greenstone, and small masses of glaucophane schist. KJf = Dothan Formation—dark gray hard metagraywacke, metasandstone, thin-bedded metabasalt, and intercalated green metavolcanic rocks (probably Late Jurassic, but believed by F. Wells and G. Walker (USGS Map GQ21, 1915) to be older than the Galice Formation shown as Ju on this map; the Franciscan Formation and the lithologically similar Dothan Formation have been extended from their type areas from the south and from the north to the Oregon-California line where the rocks are known by either name).
	KJfv	FRANCISCAN VOLCANIC AND METAVOLCANIC ROCKS	Metamorphosed basic igneous rocks within the Franciscan Formation.

STRATIGRAPHIC NOMENCLATURE— Continued

AGE	STATE MAP SYMBOL	STATE MAP UNIT <small>State Map Units listed here are not necessarily in stratigraphic sequence; the sequence used has been standardized for all sheets of the Geologic Map of California</small>	STRATIGRAPHIC UNITS AND CHARACTERISTIC LITHOLOGIES <small>(The formally named formations grouped within an individual State Map Unit are listed in stratigraphic sequence from youngest to oldest.)</small>
MESOZOIC	gr	MESOZOIC GRANITIC ROCKS UNDIFFERENTIATED	Quartz diorite, diorite, gabbro, granite and granodiorite that have not yet been differentiated in the mapping. Locally includes some ultramafic rocks.
	gr ^a	GRANITE AND ADAMELLITE (QUARTZ MONZONITE)	Adamellite at Yellow Butte. Granite and adamellite in the Seiad Valley quadrangle.
	gr ^g	GRANODIORITE	Hornblende-biotite granodiorite, and leucocratic granodiorite in the Etna and Coffee Creek quadrangles. Granodiorite in the Seiad Valley quadrangle.
	gr ^t	TONALITE (QUARTZ DIORITE) AND DIORITE	Trondhjemite in Caribou Mtn. area (Coffee Creek quadrangle), tonalite, hornblende diorite, biotite diorite, and pyroxene diorite in the Etna and Coffee Creek quadrangles. Tonalite in the Seiad Valley quadrangle.
	bi	MESOZOIC BASIC INTRUSIVE ROCKS	Hornblende gabbro and related dark diorite, augite-hornblende gabbro, hornblende gabbro and hornfels complex in the Coffee Creek quadrangle. Hornblende gabbro, and undifferentiated gabbro and related dark diorite in the Etna quadrangle. Gabbro, undifferentiated gabbro and diorite, and minor granodiorite elsewhere.
	ub	MESOZOIC ULTRABASIC INTRUSIVE ROCKS	Predominantly serpentinized peridotite. Includes some unaltered peridotite, dunite, and pyroxenite, also landslide deposits of ultramafic rocks in the Coffee Creek quadrangle. Most of the ultramafic rock in the eastern part of the map area may be part of a once-continuous sheet that intruded a low-angle fault along which Paleozoic strata to the east were thrust westward over the Abrams and Salmon Formations (Irwin and Lipman USGS Prof. Paper 410-C Article 67, 1962).
	Ju	UPPER JURASSIC MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS	Galice Formation—dark gray to black slate, phyllite, and interbedded thin to massive, light gray, tuffaceous sandstones; includes some interbedded andesitic metavolcanic rocks.
	Jml	MIDDLE AND/OR LOWER JURASSIC MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS	Potem Formation—argillite, tuffaceous sandstone and shale, and limestone; Arvison Formation—pyroclastic rocks deposited in a marine environment; volcanic breccia and agglomerate containing interbedded tuff, tuffaceous sandstone and conglomerate; some limestone.
	Jrv	JURASSIC AND/OR TRIASSIC METAVOLCANIC ROCKS	Metavolcanic rocks in the northwestern part of the Weed sheet (may be in part correlative with the Rogue Formation in Oregon and are in part interbedded within the Galice Formation). Bagley Andesite—flow rocks, volcanic breccia and pyroclastic deposits (Middle Jurassic). Pic Formation, (volcanic part)—Keratophyre, light to dark colored flow rocks, mafic flows and some pyroclastic rocks (Triassic).
	R	TRIASSIC MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS	Modin Formation—shale, sandstone, tuff, agglomerate, andesite flow rocks, limestone, and conglomerate; Brock Shale(?)—thin dark shale, in part calcareous; Hosselkus Limestone—gray, lenticular, fossiliferous limestone; Pic Formation—late, siltstone and sandstone and some undifferentiated volcanic rocks.
JURASSIC	m	PRE-CRETACEOUS METAMORPHIC ROCKS, UNDIFFERENTIATED	Chiefly undivided phyllite, chert, and metavolcanic rocks of the Western Paleozoic and Triassic Belt of Irwin (1960), (in part correlative with the Applegate Group of southwestern Oregon); includes rocks of the so-called Southwestern Devonian and Southwestern Carboniferous Belts of Diller ² the Blue Chert and part of the Lower Slate series of Hershey ³ and the Grayback Formation of Maxson (1933). In Scott Bar and Conroy Mtn. quadrangles, includes amphibolite, quartzite, and marble. Stuart Fork Formation—phyllite, quartzite, graphitic quartz-mica phyllite, greenstone and related tuff, minor marble (formerly included within the Abrams Mica Schist, Davis and Lipman, G.S.A. Bull., Dec. 1962. This formation is present in the Coffee Creek quadrangle and is probably correlative with similar rocks elsewhere in the central metamorphic belt.)
	ls	ls = LIMESTONE AND/OR DOLOMITE	White, coarsely crystalline limestone (marble) in the Seiad quadrangle and the Marble Mountains.
	ms	PRE-CRETACEOUS METASEDIMENTARY ROCKS	Quartzites, metachert, and mica schist on Yellow Butte (Macdoel 30' quadrangle). Phyllite, blue-gray thin-bedded chert, some volcanic rocks, and minor limestone have been considered as correlative with the Triassic Applegate Group. Rocks considered to be more highly metamorphosed equivalents of the Galice Formation, includes Weitchpec Schist of Oscar Hershey ³ (1906).
	mv	PRE-CRETACEOUS METAVOLCANIC ROCKS	Greenstone and greenstone schist having metasandstone interbeds of chert, argillite and limestone (these rocks may be correlative with the Applegate Group). Metabasite intrusive rock south of Gunsight Peak northward to Paradise Crags in the Hornbrook quadrangle. The "mv" unit north of Seiad Valley depicts coarse-grained, foliate and/or lineate black amphibolite of metamorphic grade similar to that of the adjacent "ps" unit. However, F. Wells (personal communication to W. Dickinson) believes that the black amphibolite resembles metamorphic phases of undoubted Applegate Group, whereas the hornblende schist ("ps" unit) is unlike anything known to be Applegate Group or its equivalents.
	Rv	PERMIAN METAVOLCANIC ROCKS	Dekkas and Noonan Formations undifferentiated (Bollibokka Group)—indurated tuff-breccia, conglomerate, and green and maroon mudstones.
	R	PERMIAN MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS	McCloud Limestone—light gray, coarsely crystalline limestone, some dark gray massive or thin-bedded limestone.
	C	UNDIVIDED CARBONIFEROUS MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS	Baird Formation—Metamorphosed maroon and green mudstone, multicolored conglomerate, sandstone, tuff, and dark gray limestone (probably Mississippian).
	CM	MISSISSIPPIAN MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS	Bragdon Formation—Metamorphosed shale, siltstone, mudstone, sandstone and coarse conglomerate.
	D	DEVONIAN MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS	Kennett Formation—Metamorphosed black shale and mudstone, and dark gray dense coral reef metamorphosed limestone.
	PALEOZOIC	Dv?	DEVONIAN AND PRE-DEVONIAN ? METAVOLCANIC ROCKS
S		SILURIAN MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS	Gazelle Formation—Essentially unmetamorphosed volcanic graywacke, dark gray to black siltstone and mudstone, siliceous and feldspathic grit, chert conglomerate, limestone and limestone conglomerate.
O		ORDOVICIAN MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS	Duzel Formation ⁴ —Pale gray-green schistose graywacke, phyllite, limestone, chert, and greenstone.
ps		PRE-SILURIAN ? METAMORPHIC ROCKS, UNDIFFERENTIATED	Lineated hornblende schist and medium to dark green plagioclase-chlorite schist in the Seiad Valley quadrangle. Chlorite-quartz-muscovite schist in the Conroy Mtn. and Seiad Valley quadrangles. Hornblende schist in the Preston Peak quadrangle. Hornblende and chlorite schists in the Scott Bar quadrangle. These units have been considered as equivalents of the Salmon Hornblende Schist and Abrams Mica Schist, however they may be nothing more than more highly metamorphosed equivalents of Triassic or Paleozoic rocks to the south. Dickinson (1962) written communication) believes that the amphibolites shown as "mv" north of Seiad Valley are of approximately equivalent metamorphic grade to the adjacent "ps" schists.
pSs		PRE-SILURIAN ? METASEDIMENTARY ROCKS	Quartz-mica schists that have been considered as Abrams Mica Schist. Grouse Ridge Formation (Davis and Lipman, G.S.A. Bull., Dec. 1962)—micaceous and feldspathic quartz schist, almandine-hornblende rocks, hornblende schist, hornblende gneisses, and calc-schists. Rocks shown as pSs? (gneissic amphibolite) are considered by Lipman (1962) as part of the Salmon Hornblende Schist.
pSv		PRE-SILURIAN ? METAVOLCANIC ROCKS	Salmon Hornblende Schist—lineated hornblende schist probably formed by the metamorphism of basaltic rocks. Included in pSv is a "transitional" Stuart Fork-Salmon unit of Lipman and Davis; the mapped northward extension of this unit is however, a greenstone that has a similar metamorphic history to that of the Stuart Fork Formation. Davis and Lipman (G.S.A. Bull., Dec. 1962) have postulated that the Stuart Fork-Salmon contact is a major low angle thrust fault.
NOTES			
¹ These units are equivalent to a part of the Western Cascade Volcanic Series of H. Williams. ² Diller, J. S., 1903. Klamath Mountain section, California. Am. Jour. Sci. 4th ser., vol. 15, pp. 342-362. ³ Hershey, O. H., 1906. Metamorphic formations of northwestern California. Am. Geologist, vol. 27, pp. 225-245. ⁴ Hershey, O. H., 1906. Some western Klamath stratigraphy. Am. Jour. Sci. 4th ser., vol. 21, pp. 58-66. ⁵ The Duzel Formation was first described by F. Wells, G. Walker, and C. Merriam (G.S.A. Bull., May 1899) as Upper Ordovician (?) based upon a fossil fauna at Horsehoe Gulch. This age was determined by Helen Duncan (of the U.S.G.S.) who recognizes the less likely possibility that this fauna may be Early Silurian (written communication from Miss Duncan, 1963). C. W. Merriam (written communication, 1963) believes that a different faunal assemblage which he collected at a later date at Horsehoe Gulch is Silurian; however, he recognizes the possibility that both Ordovician and Silurian strata may be present.			