CONTACT INFORMATION
Mining Records Curator
Arizona Geological Survey
416 W. Congress St., Suite 100
Tucson, Arizona 85701
602-771-1601
http://www.azgs.az.gov
inquiries@azgs.az.gov

The following file is part of the Grover Heinrichs Mining Collection

ACCESS STATEMENT

These digitized collections are accessible for purposes of education and research. We have indicated what we know about copyright and rights of privacy, publicity, or trademark. Due to the nature of archival collections, we are not always able to identify this information. We are eager to hear from any rights owners, so that we may obtain accurate information. Upon request, we will remove material from public view while we address a rights issue.

CONSTRAINTS STATEMENT

The Arizona Geological Survey does not claim to control all rights for all materials in its collection. These rights include, but are not limited to: copyright, privacy rights, and cultural protection rights. The User hereby assumes all responsibility for obtaining any rights to use the material in excess of "fair use."

The Survey makes no intellectual property claims to the products created by individual authors in the manuscript collections, except when the author deeded those rights to the Survey or when those authors were employed by the State of Arizona and created intellectual products as a function of their official duties. The Survey does maintain property rights to the physical and digital representations of the works.

QUALITY STATEMENT

The Arizona Geological Survey is not responsible for the accuracy of the records, information, or opinions that may be contained in the files. The Survey collects, catalogs, and archives data on mineral properties regardless of its views of the veracity or accuracy of those data.

GEOLOGY OF THE CASTLE MT.MINING DISTRICT, MONT.

W.H.Weed & L.V.Pirsson.USGS Bull.139, 1896.

For location see Tectonic Map of W.U.S.

Rock Formations:See accompanying plan.

where slaty
Algonkian: Mainly shale,rather well indurated,a.t slaty cleavage. A cliff-maker. Has limestone beds,and sandy horizons.

Cambrian: Lower,Flathead shales, 75-100' qtzite at base,above this.400' sha shale,50' ls.,260' upper shales. Above Flathead,Gallatin ls.pure.fossilif., about 1200'.

Silurian: Jefferson formation.massive ls. 550',dark gray,overlain by 165' dark fetid ls.

Devonian: Three Forks shales,here fissile ls,thin-bedded. 145'.

Carboniferous:

Quadrant Formation: 600' sandstones,shales,with pure and impure ls.
Madison
Massive ls. 400'
Thin-bedded ls 800'. Mt-forming ls.

Juratrias: ~~xxxxx xxx~~ Ss at vase,overlain by dense white ls.,then by ss;capped by red earthy shales.90' total.

Cretaceous: Kootanie? Thin-bedded ss,red earthy shale,coal seam near top, beneath Dakota qtzite.300'.

Dakota? Heavy ss bed or qtzite,at base,overlain by shaly ss and earthy shal Qtzite a cliff-maker.separates Carboniferous,ore-bearer,from shaly K beds in which prospecting NG. Basal ss s.t. cgl.

Colorado: Benton,black carbonaceous shales. Niobrara,sandy shales & impure ss.Total 2000'.

Montana:Pierre shales,2000'.Fox Hills ss. Sandy shales,ss at top.

Laramie: 500' of massive light-colored ss.with intercalations of gray shal and coal seams.

Livingston (now classed as Lance,latest K,and Ft.Union,Paleocene).Great thickness of dark ss,grits,cgls.with IB shales.

Granite: Main mass forms core of Castle Mt. Porphyritic texture. Cuts beds except in W part.(?) Near town of Castle Cumberland mine workings show contact dips 55° outward. Intruded sheets in seds.traced into main mass. CM.

Diorite: Probably pre-granite.CM.

Rhyolite flows,tuffs,breccias indicate extrusive phase of granite,

Structure,Geologic History.- E part of Rockies in Montana consists of 2 main areas of uplift, southern one the Beartooth-Showy,broad anticline with Archean core.Second uplift to NW of 1st,broad anticlinal folds of the Belt ranges,with Algonkian cores. Castle Mt.on border of 2nd uplift,forming connection between the two Belt Mt.anticlines.

Belt area on W slopes of Castle Mt. part of E end of Big Belt anticline. Anticline that forms SW part of mt. is one of the small sharp folds that mark the dying away of the greater fold. N part of MT.part ofbroad anticlinal area of the little Belt Range.

Maps shows 2 prominant anticlines ,one on E,one on W sides of mt. Axes NW,pitch SE. Biggest plunging anticline on E.Smaller one,interrupted by granite plug,on W.Overturned to E.

History: Close of Laramie area raised above sea.Mt.-building progressed during Livingston period,cordillera fully developed. Thismorogeny continued through Eocene,giving the great folds of the mt.regions,upon whose flanks the lately deposited Livingstom grits etc. were upturend and folded. At this time intrusive bosses of Crazy Mts.formed,radial dikes etc.This period was in general one of prolonged erosion.At close of Eocene,diorite boss formed. Granite boss or stock formed,cutting the folds with so little disturbance that no great amount of folding can be ascribed to the intrusion itself,though its uprise may have been precded by by an uplifting and fracturing of the beds above it.p.28. Granite shown as laccolitic massintruded in the folded Algonkian shales. Uplifting of the beds rather than replacement Violent explosive action later took place,much of sedimentary cap blown off,

rhy tuff and breccia deposited. Thicknesses to 4008 on immediate mt.slopes. Exposures of rhy now are mere remnants,

Ore Deposits.-Silver-lead, Main mine,Cumberladm produced

1890 500,000 lbs.argentiferous lead,20,000 ozAg.
1891 5,000,000 lbs.lead.
1892 300,000 lbs. lead Closed 1892,drop in Ag proce.

At 1 oz.Ag to 25 lbs.Pb, silver production,1891,200,000 oz., 1892, 12,000,total,232,000 oz.Ag. 5,800,000 lbs.Pb. At acerage price Pb, 4 c.,and Ag,95 c, total gross production about $250,000 probably half a million.

Ore arg.galena and oxid,products,ccpy,py. Gange siliceous jasper.s.t. grades to qtz.

Deposits in altered ls in CM zones about granite and diorite bosses.

Cumberland Mine:1893 panic,drop in ailver caused shut-down. For approximate location see plan. A manto,following ls.bed dipping 60°;axial length about 1200',cross*sections,80-100'(strike width) x 50-60'(perp.to dip).At bottom ore passes into narrow zone of pyritic vein-matter between a porphy.dike and the ls.On 500' level crosscut runs W to granite mass. Granite contact dips 65o under ls.,roughly paralleling bedding. Crosscut cut two decomposed porph dikes.Ls much more altered at depth,due to granite mass underlying and to dikes or intrusive sheets,more abundant at depth.

Blackhawk Mine:(Approx.location on map).130' shaft,E and W drifts at 100'.Water stopped work.OB said to be large,& to lie between a porphyry wall and altered ls.Dip 50o. Drifts seem to match in trend the E-W strike of the Cambrian,just W of Blackhawk. Ore was black,manganiferous,5-15 oz. Ag. Might turn to lead at slightly greater depth.

Legal Tender:In Blackhawk area. Shaft on smamm OB not carefully explored. Here strike is N-S;crosscuts run E-W.Ore moxed galena and cerusite,

Judge:See plan.In metamorphosed Cambrian.Two-comp.shaft 240'40 to 50' of crosscutting.Was being worked by 2 men,1894,sort and sack ore.OB said to have "outcropped at the surface".15' wide,but pay ore "forms a chute" 8" to 4' wide,20' long.May be manto.

Alice:Blackhawk area. Prospect shaft,85',exposes manto 3 x 6' in X-sect.

Iron Chief:Same area. Gossan.Ore carries 50% Pb,25 oz.Ag.Two-comp.shaft 250'.

Wherever the white Paleo limestones are found in the contact zone about the granite mass there are prospect pits.Esp.along N flanks

Grasshopper Mine:See plan.Ore manganif.with some PbS(cf.Blackhawk mine). In pocket in slightly altered black Silurian ls.

Copperopolis: Only random prospecting.In Belt shales.Small fissure veins s.t. on shear planes These veins do not pass unpward into Cambrian,hence pre*Middle Cambrian(?);perhaps only roots of OBs. But ores are rich.Ccpy,py.

Notes by E.W. Paper very poor.No mines located on plan. Sections referred to in text largely do not correspond with those givenPlate IV.

Granite stock probably extends UB to NE,indicated by abundant qp sheets there,by diorite stock,probably initial phase of intrusion of a composite stock,and by obvious modification of earlier NW open folds plunging SE. (Cf. Bingham, Marysville).

Stock arches strata as seen in NEc sections given,cuts across strata as would be seen in NW section. Probably came up between and /or along two NE master fracturesProbably another structural factor,possibly the chief factor in localization of the granite stock is the curving reverse fault that appears blotted out to the south by the ctock. This fault appear on the Tectonic Map,W.U.S. and may be a marginal thrust from the main Big Belt uplift to the SW. Here as at Bingham are two directions of deformation necessary to localize intrusions and mineralization.

Only the stock contacts could serve as channels for ore solutions ascending thru the soft Belt shales.

District looks like a "sheeper".

I-B. Limestone replacement deposits.

Castle Mt.,Montana. W.H.Weed, L.V.Pirsson:Geol.of Castle Mt.Dist.,Mont. USGS Bull.139,1896.

NE flank of Big Belt anticlinal uplift.CR Beltian, Paleo,Meso. Intrusives:granite,diorite;many dikes,sills.

Structure: Two NW anticlines plunging to SE. The composite stock elongated NE has gearlty widened the plunging anticline on the SW, strongly choked syncline bwteen the two anticlines. Localized by (1) NE fractures now largely blotted out;(2) NW reverse fault which stock has blotted out;probably marginal thsrust off Big Belt anticline.

As at Bingham,two structural features needed to localize intrusion. Stock arched Beltian shale seen in NE section;cuts across,seen in NW section.

Silver-lead replacements in altered massive Cambrian and Carbonif. ls around stock. Cumberland mine near Castle, along SE border of stock,a manto in ls dipping 60° SE parallel to stock contact here. Near porph sill or dike other deposits localized along sills or dikes also.

Only stock contacts could have furnished channels for ore solutions asceinding through them many thousand of ft.of Beltian shale.

Compare with Bingham.

(*) Long boring paper!

Castle Mtn Mining Dist., Mont. USGS Bull 139 Weed + Pirsson 1896

(1) @ Eastern Rockies;



in Belt uplift

(PE cores) + (intr. rx acidic)
(in anticlines)

(2) Pb-Ag product. ~ 1890

↓ Beartooth uplift (80 mi center to center)

(3) granite mass (8 mi max)
in folded seds



WY40
x yellowstone

(4) Note Tyin paleoz. sect



(5) Diorite pre granite

(6) QP dikes, + rhyo, tuff from granite

---

apparently a laccolithic (or ? Turnip) shaped mass of gr., acc. to W+P. close to surface, during period of Eocene uplift, folding, erosion (intr. attend) very little folding ascribable to gr. intrus. itself
→ { Same old question! stoping? assimilation? }
but - is on Lg. (see sec) rev. fit
ended in explosive volcanic activity

Weed + Pirsson say it is laccolithic, uplifted beds id, as shown in ~~strong~~ SW-NE section (✎)

BUT - gr. CUTS beds in SE-NW section (✎) poss. two NE fits?

or? - composite growth, as is partly developed on NE side of gr, w. QP dikes, Dior. stock

The curving rev. fit may also localized it

Discussion

NW - SE sec shows: ↖ ↘



MAIN PT.

(*) granite is actually at: acute intersection of a synclinal axis and reverse flt of great displacement

(*) some uplift of beds on NW flank attributable to granite but basically it, and SE, are cut off by intrusion

(*) poor evidence for NE flts, acc. to mapping at least → during intrusion

(*) uplift on E normal for sole or f.wall of thrust

(*) uplift on W is problem: ↘ due to this →?



certainly, acc. to mapping, is poor for this



(1) No (concentric or radial folding)

(2) No satellitic QP etc

(3) No apparent disruption or bending of the beds on W.

any "Tensional" area at Time of Intrusion was on THE EAST

STATE OF MONTANA
Tim Babcock, *Governor*

BUREAU OF MINES AND GEOLOGY
E. G. Koch, *Director*

Bulletin 64
January, 1968

GEOLOGY AND ORE DEPOSITS of the CASTLE MOUNTAIN MINING DISTRICT MEAGHER COUNTY, MONTANA

By Allen S. Winters

STATE OF MONTANA
BUREAU OF MINES AND GEOLOGY
E. G. Koch, Director

BULLETIN 64

GEOLOGY AND ORE DEPOSITS OF THE CASTLE MOUNTAIN MINING DISTRICT, MEAGHER COUNTY, MONTANA

by

Allen S. Winters

MONTANA COLLEGE OF MINERAL SCIENCE AND TECHNOLOGY
Butte, Montana
January 1968

For sale by

Montana Bureau of Mines and Geology

Room 203-B, Main Hall

Montana College of Mineral Science and Technology

Butte, Montana 59701

Price 35 cents.

CONTENTS

	Page
Abstract	1
Introduction	2
Purpose and scope	2
Field work	2
Previous work	2
Acknowledgments	3
Location and accessibility	3
Surface features	3
Topography	3
Drainage	4
Climate	4
Vegetation	4
Glaciation	6
General geology	6
Sedimentary rocks	6
Precambrian	8
Cambrian	8
Devonian	9
Mississippian	9
Mississippian-Pennsylvanian	10
Jurassic	10
Cretaceous	10
Quaternary	12
Igneous rocks	12
Intrusive rocks	12
Extrusive rocks	13
Structure	14
Regional folding and faulting	14
Folds	14
Fractures and faults	15
Precambrian-Cambrian problem	16
Mining history and production	16
Ore deposits	22
Classification	22
Contact metasomatic	22
Metasomatic replacement	23
Fissure veins	23
Replacement veins	23
Paragenesis	23
Oreshoots	25
Depth of mineralization	26
Structural controls	26
Stratigraphic controls	26

Contents--contd. Page

Alteration 27
Contact metamorphism 27
Hydrothermal alteration 29
Geochemical studies of the district 29
Grid survey 29
Analysis of heavy-metal anomalies 30
Analysis of zinc anomalies. 31
Traverse surveys. 32
Traverse 1, Powderly 32
Traverse 2, Judge 35
Traverse 3, Belle of the Castle. 38
Traverse 4, Yellowstone 38
Miscellaneous traverses. 38
Descriptions of mines and prospects 40
Cumberland 40
Yellowstone 43
Great Eastern and Great Western. 44
Jumbo 45
Judge 46
Blackhawk and Alice. 47
Legal Tender. 49
Iron Chief 49
Powderly. 50
California 50
Homestake. 52
Merrimac 53
Hidden Treasure 54
Silver Star. 54
Solid Silver 54
Broadway. 55
Belle of the Castle 56
Copper Bowl and Copper Kettle. 56
Milwaukee 58
Ruby adit 58
Princess 59
Golden Eagle. 60
Antelope. 60
Etta 60
Felix Crexent 60
American 60
Grasshopper 61
Suggestions to prospectors 61
Summary and conclusions 61
Appendix--Geochemical analyses, Yellowstone mine 63
References cited 64

ILLUSTRATIONS

Page

Plate

1. Geologic map of Castle Mountain mining district . . . . (in pocket)
2. Plan and sections of Cumberland mine. . . . . (in pocket)
3. Plan of Yellowstone mine . . . . . (in pocket)
4. Claim map of Castle Mountain mining district . . . . . (in pocket)

Figure

1. Index map showing location of Castle Mountain district . . . 5
2. Geochemical anomaly map, heavy metals 33
3. Geochemical anomaly map, zinc 34
4. Traverse 1, Powderly 36
5. Traverse 2, Judge 37
6. Traverse 3, Belle of the Castle 39
7. Traverse 3, Belle of the Castle 40
8. Map of surface workings, Blackhawk area 47
9. Map of Powderly mine 51
10. Map of Hamden adit 53
11. Map of Corliss adit 55
12. Surface map of Copper Bowl-Copper Kettle 57
13. Map of Copper Bowl adit 59
14. Map of Ruby adit 59

TABLES

Table

1. Stratigraphic column of the Castle Mountain district 7

GEOLOGY AND ORE DEPOSITS OF THE CASTLE MOUNTAIN MINING DISTRICT, MEAGHER COUNTY, MONTANA

by

Allen S. Winters*

ABSTRACT

The Castle Mountain mining district is a mountainous area on the southeastern flank of the Castle stock, which rises to an altitude of 8,606 feet above sea level. Sedimentary rocks ranging from Precambrian to Cretaceous have been uplifted by intrusion of two separate stocks, which produced both longitudinal and radial fractures. Metamorphism is not strong, but igneous and sedimentary rocks both have been hydrothermally altered by low and medium temperature processes.

Metalliferous deposits contain lead, zinc, and silver and minor copper, manganese, and gold. The sulphide minerals are commonly associated with jasper and magnetite, and are found mainly in altered Paleozoic limestone.

Geological maps and geochemical dispersion patterns of the district and a few of the properties are presented, along with production data and historical information.

---

*Present address, The Anaconda Company, Butte, Montana.

INTRODUCTION

The Castle Mountain mining district was discovered in the 1880's and gained considerable prominence as a producer of lead, silver, and zinc. Since the turn of the century, however, the district has had an erratic history of production and activity.

In recent years, in expectation of better metal prices, lessees and owners have begun to reopen and develop a few of the larger mines. The small production, however, has all come from the Cumberland and Yellowstone mines.

PURPOSE AND SCOPE

The aim of this report is to present the results of two separate studies. The first included geological mapping and a study of the district's mineral deposits, ores, and economic potential. Intended to determine the feasibility of geochemical prospecting, the second study included a series of selected traverses and a grid soil-sampling program that covered the district.

FIELD WORK

Field work was done during the summers of 1963 and 1964. Virtually all the known mineral deposits were investigated, and accessible workings were mapped. Air photos were used in mapping and to find the various mines in the district. The laboratory investigations were made during the winter and spring months of 1963 and 1964.

PREVIOUS WORK

The oldest published report on the Castle Mountain district is that of MacKnight (1892), who briefly described some of the mines in the district. Published by the U.S. Geological Survey in 1896, the first geological study of the district and surrounding area was made by W. H. Weed and L. V. Pirsson in 1894. The report includes studies of the lithology, petrology, and general geology of the area. Stone (1909) mapped the Eagle Formation near Warm Springs Creek; Stone and Calvert (1910) made stratigraphic studies of the Livingston Formation; Gardner and others (1946) measured and described the Cretaceous, Jurassic, and Carboniferous formations; and Hanson (1952) studied the Cambrian stratigraphy of the mountains. Tanner (1949) made the first detailed geological map of the Castle Mountains, but his main interest was the correlation of the upper Cretaceous strata. Describing the iron and nonferrous mineral deposits respectively, Goodspeed (1945) and Roby (1950) made the only economic studies of the district.

ACKNOWLEDGMENTS

It is a pleasure to express gratitude for the cooperation and assistance given by supervisors, operators in the district, and colleagues; to Mr. U. M. Sahinen, associate director of the Montana Bureau of Mines and Geology, for sponsorship and consultation; to Dr. F. N. Earll for advice, guidance, constructive criticism, and consultation during the course of the field work; to Mr. C. R. Oliphant of White Sulphur Springs, Mr. Carl Voss of Helena, Mr. Jack Oliver of Harlowton, and Messrs. George Volseth, Kay Burg, Paul Grande, and Claude Wessel of Martinsdale, who were most generous with their time and who allowed free access to their properties and private files; and to Messrs. D. C. Lawson, F. P. Jones, and R. B. Holmes for invaluable assistance rendered in assaying, rock analysis, and map preparation.

LOCATION AND ACCESSIBILITY

The Castle Mountain district, Meagher County, Montana (Fig. 1), as the term is used in this report, lies principally within T. 8 N., R. 8 E., plus the western and southern sections of T. 8 N., R. 9 E., and T. 9 N., R. 8 E. (Pl. 1). The central part of the district is approximately 10 miles northwest of Lennep, at lat 46° 28' N., and long 110° 41' W.

The district is accessible by four gravel roads; two from Lennep, one from Checkerboard, and one from White Sulphur Springs, the county seat of Meagher County. The only maintained road, however, is from Castle to Lennep, which is connected to White Sulphur Springs by 35 miles of improved gravel and paved roads.

The main line of the Chicago, Milwaukee, St. Paul, and Pacific Railroad passes through Lennep and Ringling, where a branch line from White Sulphur Springs connects.

SURFACE FEATURES

TOPOGRAPHY

The Castle Mountain mining district is a mountainous area on the eastern slope of the Castle Mountains. In the map area, the mountains have a maximum relief of approximately 2,600 feet, from Castle at about 6,000 feet to the summit of Elk Peak at 8,606 feet. Most of the mines are in upland gulches and on rounded ridges at altitudes between 6,300 and 7,500 feet.

The topography of the map area is closely related to the paths of glaciers and to drainage patterns subsequently produced by active

down-cutting streams, which occupy narrow canyons. Steep slopes gradually grade into high-level parks, some of which are as much as 1 mile wide and 5 miles long.

The topography is also related to the folding of the sedimentary rocks. Anticlinal and synclinal ridges are clearly developed near Lennep, but in the map area the weaker sedimentary rocks and folds have provided channels only for the tributaries; the main creeks, at least in part, have been controlled by glacial paths and fracturing.

The igneous rocks are resistant to erosion, and their contacts are distinguished by a sharp steepening in slope. Of the sedimentary rocks, the Kootenai, Mission Canyon, and Belt form conspicuous ridges, whereas the Colorado, Big Snowy, and Park occupy topographic lows.

DRAINAGE

Numerous streams flow through the district and should provide ample water for all present and future mining and milling needs. The water originates in many mountain springs, and although some of the tributaries are dry in late summer, the main streams all have a permanent flow.

CLIMATE

The fairly moist climate of the Castle Mountains is noticeably different from the generally semiarid climate of the rest of Meagher County. Weather data have not been systematically collected in the map area, but on the basis of the statistics from Findon and Kings Hill the annual precipitation is estimated between 17 and 28 inches, including an annual snowfall of about 180 inches. The mean annual temperature is about 37 degrees, maxima in the 90's coming during the late summer months and minima in the -40's in January or February (Roby, 1950, p. 6).

In general, the climatic conditions are by no means prohibitive but are not very favorable for year around small operations.

VEGETATION

The slopes in the vicinity of the mining district are timbered with a dense growth of lodgepole pine and some fir and spruce. Native range grasses cover the open ridges and parks and is accompanied by sagebrush at the lower elevations.

Present and future mine operators should have little trouble in finding an ample supply of lodgepole, which if properly treated, makes acceptable mine timber.



Figure 1.--Index map showing location of Castle Mountain district.

GLACIATION

Numerous glacial erratics cover the lower slopes adjacent to Alabaugh Creek. This debris is the most conspicuous evidence of glaciation in the district.

Valley glaciers originated on the flanks of Elk Peak where the ice accumulated in small recesses and carved out small cirques and glacial troughs. Following pre-existing stream-cut valleys, the glaciers removed a few of the surface irregularities but only slightly modified the original landscape.

Weed and Pirsson (1896, p. 144) stated that glacial evidence is most prominent from 7,000 to 5,400 feet, where both lateral and end moraines can be recognized. The lateral moraines are by far the more abundant. These deposits should more properly be termed glacial debris, as they are not distinguishable landforms. The debris consists chiefly of granitic rocks but includes some Precambrian and Paleozoic blocks. End moraines can be observed on the valleys of Boulder, Hamilton, and Fourmile Creeks.

GENERAL GEOLOGY

SEDIMENTARY ROCKS

Sedimentary rocks cover a large part of the map area and include a wide variety representing a vast interval of time--Precambrian to Cretaceous (Table 1).

The combined Jefferson and Three Forks, the Kibbey and Otter, and the Rierdon, Swift, and Morrison Formations were mapped as three units because of their small thicknesses or poor exposures. Belt shale and Lower Cretaceous shale were mapped as the Piegan and Colorado Groups respectively.

No major angular unconformities were observed within the district. The only angular unconformity reported in the area is that between the Cambrian and Precambrian strata. Tanner (1949, p. 11) reported that this angle ranges from 0 to 12 degrees. Major unconformities separate the Jurassic from the Pennsylvanian, and the Devonian from the Cambrian.

Stratigraphic sections were not measured within the district, as numerous studies have already been made in and near the map area. Thicknesses of the formations in this report are only rough estimates, hence may differ from those reported by previous workers (Weed and Pirsson, 1896, p. 30-55; Gardner and others, 1946, p. 30-36; Hanson, 1952, p. 35-36; and Tanner, 1949, p. 11-47).

Table 1. --Stratigraphic column, Castle Mountain mining district, Meagher County, Montana.

Age		Stratigraphic unit		Thickness (feet)	Lithology
Quaternary	Recent	Alluvium			Silt, sand, and gravel
	Pleistocene	Glacial debris			Unsorted sand, gravel, and boulders
Cretaceous	Upper	Montana Group	Hell Creek	1,000±	Gray tuff and sandstone
			Lennep	300	Dark sandstone and shale
			Bearpaw	900±	Black shale and siltstone
			Judith River	1,200	Brown tuff and sandstone
			Eagle	450	White sandstone and gray siltstone
	Lower	Colorado Group		2,300	Interbedded shale and sandstone
		Kootenai		380	Sandstone and red siltstone
Jurassic		Morrison		350	Brown siltstone
		Ellis Group	Swift	42	Brown sandstone
			Rierdon	15	Gray limestone
Pennsylvanian(?)-Mississippian		Amsden		400	Pinkish-gray limestone
		Big Snowy	Otter	150	Green shale
			Kibbey	120	Red siltstone
		Mission Canyon		400	Massive gray limestone
		Lodgepole		800	Blue-gray limestone
Devonian		Three Forks		145	Thin-bedded limestone
		Jefferson		520	Massive buff to brown dolomitic limestone
Cambrian		Red Lion (Dry Creek Member)		60	Tan siltstone and shale
		Pilgrim		280	Blue-gray limestone and limestone conglomerate
		Park		360	Black shale
		Meagher		160	Alternating limestone and siltstone
		Wolsey		400	Green micaceous shale
		Flathead		125	Pink sandstone and quartzite
Precambrian	Belt Series	Piegan Group			Red argillite and gray limestone

Precambrian

Belt Series. --Named and described by Peale (1893, p. 12), the Belt rocks are generally thought to be Algonkian in age. Tanner (1949, p. 13) correlated the 2,500 feet of Belt argillite within the map area with the Spokane Formation, but these rocks may represent the Spokane, Empire, Helena, and March Formations recognized in the nearby Belt Mountains (Ross, 1963, p. 3). No attempt to differentiate these sediments was made; they are referred to the Piegan Group.

Mostly chocolate color to deep purple, the Belt argillite near Blackhawk is intensely altered and forms sharp ridges. Several prospects have investigated iron and copper in the calcareous part, which possibly represents the Helena Formation.

Cambrian

Flathead Formation. --The Flathead Formation is composed of fine- to coarse-grained quartzitic sandstone. In most of the map area the formation is completely metamorphosed to metaquartzite, individual grains being indistinguishable with the naked eye; scattered hematite grains give the rock a speckled appearance. Farther from the intrusive bodies the formation can be divided into two units. Composed of large pink quartz grains, the basal 50 feet of the formation is cross bedded. The upper 75 feet is composed of fine clear to white quartz grains. Both members are cemented by silica, and the formation forms a conspicuous ridge near Blackhawk.

Wolsey Formation. --The Wolsey Formation can be divided into three units--a lower 50 feet of dark-brown fissile shale and siltstone, a middle 250 feet of tan to brown thin-bedded limestone and siltstone, and an upper 100 feet of tan to olive-green micaceous shale and calcareous siltstone. The formation is nonresistant and occupies a depression between the Flathead and Meagher Formation.

Meagher Formation. --In the map area, the contact between the Meagher Formation and the Wolsey is poorly defined and gradational. The Meagher is composed of 40 feet of alternating limestone and siltstone beds overlain by 120 feet of dirty limestone. The lower thin-bedded limestone and siltstone beds are mostly tan to buff but contain a few olive-green shale beds similar to those in the Wolsey.

Park Shale. --Approximately 360 feet of gray to black Park Shale crops out in the map area. The shale is metamorphosed to hornfels in the vicinity of the Yellowstone mine, where it resembles Belt argillite. Farther from the intrusive body the formation is nonresistant and its outcrop is marked topographically by a slight depression.

Pilgrim Limestone. --The Pilgrim Limestone is approximately 280 feet thick in the map area. The lower 75 feet of the formation is easily distinguished by the pebble conglomerate beds. The upper limestone beds are buff and display intraformational conglomerate limestone lenses. Close to intrusive bodies the limestone is recrystallized, and cherty seams and lenses weather in relief. Some ore was deposited in the metamorphosed limestone.

Red Lion Formation. --The Red Lion Formation is represented in the map area by the Dry Creek Member, which is composed of 60 feet of tan to green shale interbedded with red siltstone layers. Near the Judge mine, the shale is metamorphosed to chocolate-color hornfels that forms a ridge 2 to 4 feet high.

Devonian

Jefferson Limestone. --The Jefferson Limestone is composed of approximately 520 feet of massive dolomitic limestone generally characterized by a fetid odor. The limestone grades from buff to brown away from intrusive contacts. Where the limestone is recrystallized, some ore deposits may occur.

Three Forks Shale. --The Three Forks Formation is not a mappable unit in the Castle Mountains. Weed and Pirsson (1896, p. 38) assigned 145 feet of thin-bedded limestones to this formation, but in the map area the formation could not be distinguished, as intrusive rocks have obscured its normal character.

Mississippian

Lodgepole Limestone. --Approximately 800 feet of blue-gray to dark-gray thin-bedded limestone constitutes the Lodgepole Limestone in the map area. These beds are very fossiliferous and together with the Mission Canyon Formation, they form large outcrops and support many ridges. Near the intrusive bodies, the formation is completely recrystallized, and ores mined from deposits in the marble have contributed much of the district's production. Except where the rocks are metamorphosed, the upper contact with the Mission Canyon Formation is distinct.

Mission Canyon Formation. --Massive buff to gray limestone of the Mission Canyon Formation overlies the Lodgepole Limestone. It is 400 feet thick, generally light in color, and is characterized by numerous concretions of brown chert. The formation is very cavernous, and two small natural bridges can be seen on Warm Springs Creek. Where sharply folded, the limestone is brecciated and iron stained. Mines in the Mission Canyon Formation have contributed at least half of the district's production.

Kibbey Formation. --The Kibbey Formation consists of 120 feet of thin-bedded limestone, sandstone, and siltstone. Generally marked by a belt of red soil, the lower contact with the Mission Canyon is very distinct. Less definite is the upper contact, where the red siltstone and limestone layers grade into the buff limestone and pale-green shale of the Otter Formation.

Otter Formation. --The Otter Formation is composed of 150 feet of alternating layers of shale and thin-bedded limestone. Near the lower contact the shale is green, but it gradually darkens upward to gray and black toward the upper contact with the Amsden Limestone. The limestone beds are generally buff or mottled pink.

Mississippian-Pennsylvanian

Amsden Limestone. --Approximately 400 feet of pink to blue-gray Amsden Limestone crops out in the district. This limestone is thin-bedded and mottled pink near the lower contact, but toward the upper contact it is slightly dolomitic and darker. An accurate age has not been established for the formation in the map area, but probably both Mississippian and Pennsylvanian rocks are included.

Jurassic

Rierdon(?) Formation. --The presence of the Rierdon Formation within the map area is questionable, but 15 feet of dark-gray fossiliferous limestone was assigned to the formation.

Swift Formation. --The Swift Formation consists of 42 feet of calcareous sandstone, which can be divided into two units, a lower 3 feet of brown conglomerate and an upper 39 feet of reddish sandstone. The basal conglomerate contains both calcareous and siliceous pebbles and has a greenish cast, owing to its contained glauconite.

Morrison Formation. --The Morrison Formation is approximately 350 feet thick in the district and consists of reddish-brown siltstone in the lower half and gray siltstone in the upper half and contains many thin beds of sandstone. The formation is nonresistant. Tanner (1949, p. 45) pointed out that seams of shale and coal can be found near the contact with the Cretaceous sandstone. This relationship was not observed within the district, but was observed on Checkerboard Creek, where a coal bed has been worked.

Cretaceous

The Cretaceous sediments are unmineralized and were not studied except for the purpose of establishing contacts. Contacts are the same as those thosen by Tanner (1949), who studied the beds in detail.

Kootenai Formation. --The Kootenai Formation is composed of approximately 380 feet of sandstone and siltstone. Within the map area the formation can be divided into two separate lithologic units. The lower unit seems to rest unconformably on the Morrison Formation and is composed of nearly 100 feet of cross-bedded gray to reddish sandstone, conglomeritic near the base and locally metamorphosed to a metaquartzite that forms a sharp ridge. The upper unit is composed of approximately 280 feet of red to purple siltstone and silty sandstone.

Colorado Shale. --Consisting of about 2,300 feet of black shale and alternating beds of sandstone and siltstone, the Colorado Shale has not been studied in enough detail to permit subdivision of the group into separate formations. From Tanner's (1949, p. 47) description, three major lithologic units persist, a lower unit of black fissile shale interbedded with lenses of sandstone, a middle unit of drab to green shale, and an upper unit of gray to tan fissile shale.

Eagle Sandstone. --The Eagle Sandstone can be divided into two separate units. The lower 300 feet of the formation consists of white cross-bedded coarse-grained sandstone interbedded with some siltstone beds. Numerous coal seams in the siltstone beds have been prospected without success on Warm Springs Creek. The upper 150 feet of the formation is composed of gray siltstone and calcareous sandstone, which forms a small ridge near Warm Springs Creek.

Judith River Formation. --Sediments mapped as the Judith River Formation are actually the Judith River and Claggett Formations combined, but the two cannot be separated in the map area (Stone and Calvert, 1910, p. 745). Approximately 1,200 feet thick, the formation is composed of brown to gray tuff and fine-grained sandstone.

Bearpaw Shale. --Approximately 900 feet of black fissile shale and siltstone make up the Bearpaw Formation in the map area. Tanner (1949, p. 70) divided the formation into two units, the lower composed of black fissile shale and the upper composed of dark-gray claystone and siltstone.

Lennep Formation. --The Lennep Formation is approximately 300 feet thick in the district and is composed of dark tuffaceous sandstone and dark shale (Stone and Calvert, 1910, p. 746). The formation generally forms conspicuous orange ridges and contains several gabbroic sills.

Hell Creek Formation. --The Hell Creek Formation is reported to be 900 to 1,000 feet thick near the map area (Tanner, 1949, p. 17). Only a small patch of Hell Creek beds crop out in the area. These beds consist of medium to dark-gray tuff and fine-grained sandstone.

Quaternary Deposits

Unconsolidated deposits composed of silt, sand, gravel, and large boulders are common in the district. The ridges are commonly covered by glacial debris, and stream channels have been filled with alluvial deposits.

IGNEOUS ROCKS

The igneous rocks of the district are generally acidic and consist mainly of varieties of the granite group. Basic dikes are present in the district and generally represent a late stage intrusion of an early differentiate.

Several periods of igneous activity can be recognized in the district. The oldest intrusive body, the Blackhawk stock, is composed of diorite. A few basic dikes and possibly some andesitic flows outside the district are associated with this early intrusion. After the initial igneous activity had ceased, the main mass of granite was forced into the sedimentary rocks west of the diorite stock. The granite of the Castle stock is exposed over an area of approximately 33 square miles, and is roughly ten times the size of the Blackhawk stock. Associated with the Castle stock are large sheeted zones and radial dikes. Violent eruptions followed the intrusions, and masses of rhyolite and rhyolite breccia were ejected.

Intrusive Rocks

Granite. --The Castle stock is composed of fine- to medium-grained granite and granite porphyry. The porphyritic nature of the rock is best exhibited near Elk Peak, where numerous feldspar phenocrysts exceed $\frac{1}{2}$ inch in length. The groundmass, however, is generally fine-grained and composed of subhedral grains of feldspar and dark quartz.

Orthoclase, oligoclase, quartz, biotite, and hornblende are the chief rock-forming minerals, named in decreasing order of abundance. On fresh exposures, the rock is light colored and locally has a pinkish tint, owing to the orthoclase phenocrysts. The feldspars are somewhat kaolinized, and all biotite is slightly altered to chlorite. Accessory minerals include magnetite, zircon, and a small amount of apatite. Weed and Pirsson (1896, p. 100) reported that tourmaline and fluorite are very common in the stock to the north of the map area.

That the stock should be named the Castle Mountain is understandable, as the granite weathers into high crags or castles.

Diorite. --The chief rock of the Blackhawk stock is medium-grained diorite but toward the southern and eastern boundaries of the stock the rock is granodiorite, which weathers much the same as the granite of the Castle stock.

In the central portion of the stock the rock is composed of oligoclase, the mineral present in greatest quantity, and orthoclase, biotite, and some augite. A few rounded grains of quartz can be found. Toward the contact with sedimentary rocks, the diorite is more acidic, and many grains of quartz can be observed.

The eastern section of the stock near Blackhawk resembles pinkish-gray granite. Tanner (1949, Pl. 1) mapped the diorite and granodiorite facies as separate stocks, but both masses seem to be parts of a single stock. Tanner also showed Cambrian limestones separating the stocks, but what he observed may have been a small roof pendant.

Dacite Porphyry. --The dacite porphyry intrusive bodies were formerly classified as quartz porphyry by Tanner (1949) and Weed and Pirrson (1896). These rocks are found as intruded sheets and dikes around the edge of the Castle stock.

The rocks are felsitic and mostly gray. Near the Castle stock the rock is speckled by dark subrounded quartz phenocrysts but farther from the contact the quartz phenocrysts are less abundant.

The rock is composed of oligoclase, orthoclase, quartz, biotite, and hornblende. Feldspars are partly kaolinized, and the biotite is altered to chlorite. Near the Yellowstone mine the rock assayed 66 percent SiO_2 compared to 69 percent SiO_2 near the granite stock. Weed and Pirsson (1896, p. 99) reported local sericitization and alteration of the biotite and hornblende to magnetite.

Two separate stages of dacite intrusion were observed. In the Hensley Creek area, strongly mineralized dacite porphyry dikes cut older dacite sheets.

Syenite Porphyry. --Four sills form conspicuous ridges south of the townsite of Castle where they have intruded Cretaceous sedimentary rocks. The igneous rocks consist of a dense gray groundmass of feldspar containing large phenocrysts of hornblende and pink orthoclase. Biotite, partly altered to chlorite, is also present. Tanner (1949, p. 57) pointed out that the orthoclase feldspar phenocrysts are aligned parallel to the strike of the sills.

Extrusive Rocks

Rhyolite. --Rhyolite, exposed along Fourmile and Bonanza Creeks, is typically gray to yellow and contains a few phenocrysts of quartz and white feldspar. The composition ranges from rhyolite to possibly rhyodacite, depending on the silica content, which seems to differ considerably in different parts of the same flow.

STRUCTURE

Regional Folding and Faulting

The Smith River intermontane valley is bordered on the east by the Little Belt Mountains and on the west by the Big Belt Mountains. Within this valley and northwest of the Crazy Mountains, rise the Castle Mountains as a distinct geographic unit related to neither of the Belt ranges.

Flanking the Castle Mountains on the east and west are two large anticlinal folds, which plunge to the southeast. On the west lies the Warm Springs anticline, which is cut in the center by the Castle stock. This anticline is overturned to the east on its southern end near Warm Springs Creek. Weed and Pirsson (1896, p. 23) believed that this anticline is related to the southeastern end of the Big Belt anticline. On the eastern flank of the stock is a lateral offshoot of the Little Belt anticline.

The Castle Mountains also lie near the eastern border of the Rocky Mountain disturbed belt. Thrust and normal faults trending northwest are common north and east of the map area.

Folds

Many small folds are found within the map area. The largest of these is the Corral Creek anticline. This anticline, whose axial trace trends S. 24° E., plunges about 16° to the southeast. Flows of rhyolite cover the central part in the map area but the Mission Canyon Limestone is exposed south of the flows. Farther north the Blackhawk stock interrupts the anticline but the effects of folding can be observed in Belt argillite north of this stock.

Along the same trend as the Corral Creek anticline lies a smaller fold just east of the townsite of Blackhawk. The small anticline plunges about 32° to the southeast. Belt rocks are exposed in the core of the anticline, and a longitudinal fault cuts its western flank.

Along the south border of the diorite are several smaller folds, which seem to have been produced by the intrusion. The axial traces of these folds generally trend parallel to the Corral Creek anticline. Broadly, however, the area between Corral Creek and Warm Springs Creek is a north-trending syncline. The synclinal fold is excellently exhibited in the Cretaceous beds 2 miles east of Castle where the fold trends S. 18° W., and plunges about 42° to the south.

Approximately 1 mile east of Castle Lake, Alabaugh Creek cuts through the center of a small anticlinal fold whose axial trace trends S. 20° E. This anticline, which is cut off at the north by the Castle stock, has Big Snowy

beds exposed in its core near the granite contact. The anticline plunges about 31° to the southeast. Adjoining the anticline on the west is a tightly folded syncline. Plunging about 22° to the southeast, the syncline trends N. 54° W. Altered shale of the Colorado Group is exposed in the core of the syncline near Castle Lake.

Fractures and Faults

Faults and fractures in the district can be placed in four separate groups according to their age and strike. The oldest faults in the area are longitudinal faults associated with the Blackhawk stock. The largest fault of this group is a northwest-trending vertical fault north of the townsite of Blackhawk. It has a length of approximately 2 miles. Locally, the fault may be a high-angle reverse fault, as the Pilgrim Limestone is slightly overturned on the western limb of the Blackhawk anticline, owing to drag folding. This fault also has some rotational movement, the northern part having greater vertical displacement. Where the fault cuts across the East Fork of Checkerboard Creek, Meagher Limestone is in contact with Belt rocks. This relationship represents a total displacement of not less than 300 feet. The pivot point of the fault is on the North Fork of Bonanza Creek east of the townsite of Blackhawk.

Radial faults and fractures also were produced while the sedimentary rocks were being domed by intrusion of the Blackhawk stock. The largest of these radial fractures are north of the townsite of Blackhawk, and they have been filled with material similar to the granodiorite of the stock. Three larger dikes can be traced for several miles to the east, but little or no displacement of the sedimentary rocks was noticed.

The second group of faults and fractures is associated with the Castle stock. Radiating from the center of the stock, the structures are generally vertical. They have been filled with dacite porphyry and are excellently exhibited in the Paleozoic limestone near Blackhawk and in the Belt argillite between Hensley and Robinson Creeks.

Longitudinal faults generally parallel the eastern edge of the Castle stock. Hamilton, Hensley, and Robinson Creeks have actively exploited these planes of weakness, which generally trend north to northwest. There is little strike-slip movement on the faults but the Hamilton Creek fault has a vertical displacement of at least 150 feet.

The fourth group of faults and fractures includes those that have been mineralized. In general these faults offset the older dikes and slips but have little strike-slip displacement. In general, the veins are believed to be younger than the dike-filled fractures because the dikes are believed to have provided the channels for the mineralizing solutions. The fault vein on which

the Blackhawk and Alice mines are located is observed to offset the northeast-trending dikes approximately 5 to 10 feet to the right. In the Corliss mine, however, the dike-filled fault offsets the vein 3 to 4 feet to the left.

Precambrian-Cambrian Problem

An interesting structural problem is provided by the absence of the Flathead and at least part of the Wolsey in the central portion of the district. Near Boulder Creek, the Meagher Formation is separated from Beltian rocks by a sheet of dacite porphyry, which is generally less than 100 feet thick.

Several explanations for this peculiar situation are possible, but because of the poor exposures no definite solution to the problem is offered. Nondeposition seems very likely because the basal Cambrian formations are known to have extreme variations in thickness. Because of the complicated series of faults required, faulting seems less likely to account for the absence of the sediments. Compressive squeezing could possibly account for the absence of part of the Wolsey Formation but is unlikely to explain the disappearance of the Flathead Formation. Assimilation of the sediments is also possible, but the dacite sheet would be expected to become more acidic with the addition of silica from the Flathead. Additional field work would be needed before an adequate explanation is apparent.

MINING HISTORY AND PRODUCTION

The Castle Mountains remained relatively unknown until they were first noticed by Captain James Clift, who passed by the mountains on his exploratory march across Montana in 1869. The igneous core and other rock formations of the mountains, however, were not recognized until Captain W. H. Ludlow, accompanied by geologists G. B. Grinnel and E. S. Dana, made their exploratory trip along the Musselshell in 1873 (Weed and Pirsson, 1896, p. 17).

A few prospectors may have passed through the mountains on their way to the copper veins at Copperopolis in the 1870's, but these men failed to recognize the favorable geologic conditions in the mountains. The first prospector known to have really examined the area was C. Barnes, the U.S. Postmaster of White Sulphur Springs, who first began to prospect the hills and gulches of the Castle Mountains in 1881. Realizing the mineral potential of the district, he continued to prospect and in 1884 located the Blue Bull claim on Robinson Creek and the Princess on Alabaugh Creek. News of these two locations aroused a few other prospectors, but it was not until the Hensley family discovered the rich Cumberland and Yellowstone oreshoots that a sincere interest in the area began.

Reports of the Hensley's strike spread rapidly, and virtually overnight three booming mining camps sprang to life. Castle Town, the largest of these camps, was located on Castle Creek, a mile below the Cumberland mine. At the peak of its career Castle Town boasted a population exceeding 1,500, and numerous business establishments were set up to accommodate the tremendous influx of people (personal communication, C. R. Oliphant).

Numerous partnerships and several companies were immediately formed to work the claims as soon as they were discovered. The ore was first shipped by ox and mule team 75 miles to Livingston, Montana. From Livingston, the ore was then shipped to eastern smelters for processing. It was soon realized that if the mines were to continue to develop, the ores would have to be smelted locally.

In 1887 construction of the first smelter began on Alabaugh Creek. The smelter was built to treat the ore from the Yellowstone mine. The following year a newer and more modern smelter was constructed at the Cumberland mine site to treat the large tonnage that was being extracted from this property. This smelter processed ore from 1889 to 1893, and at the peak of its production could treat 35 tons of oxidized lead ore per day. As the mines developed, however, the ore changed from oxide to sulphide, and the smelters could not efficiently treat it, owing to increasing zinc content with depth.

G. C. Swallow (Swallow and Trevarthen, 1890, p. 17), inspector of mines for Montana, stated that in 1889 the Connellsville coke used in the furnaces cost the smelters $45 per ton delivered at the plants, that refining costs ran $16 per ton, and that the expense of shipping to Aurora, Illinois, was $22 per ton. He then estimated that if a railroad were built into the area the mines could save $15 to $20 a ton. The smelters soon closed, and the residents of Castle then began trying to promote a railroad into "Leadville" as they fondly called the district.

Immediate dreams of a railroad, however, were shattered when Congress demonetized silver in 1893. This action quickly put an end to the district's brief but vigorous life, and the mines closed in swift succession.

R. A. Harlow finally promoted enough money to construct a railroad from Loweth to Leadboro, half a mile below Castle Town. The line was known as the Jawbone because of the amount of persuasion and discussion it took to raise the funds. After the railroad was built, the district experienced a small revival from 1896 to 1898. Most of the ore shipped by the Jawbone, however, came from the stockpiled ore left on the dumps, and the largest producer, the Cumberland, never reopened, owing to the low prices of silver and lead (Roby, 1950, p. 17).

After the dumps from the Cumberland and Yellowstone mines had been shipped to East Helena, the district again lapsed into a state of inactivity and Castle Town soon became a ghost town. Sporadic mining continued on a small scale throughout the years, but not until 1957 were regular shipments again produced from the district. Since 1957 the Cumberland and the Yellowstone have both been active and have shipped considerable ore.

The following chronological account of mines and production of ore in the Castle Mountain district is compiled from annual volumes of the Report of the Director of the Mint (1888-1903), Minerals Resources of the United States (1905-1931), Minerals Yearbook (1932-1963), and from smelter returns recorded in East Helena. Only the largest producers are mentioned, and the ensuing notes are restricted to mines described in this report.

1888--Considerable development took place in the district, and the Cumberland, Yellowstone, and Great Eastern mines shipped most of the district's ore to eastern smelters. Meagher County produced $100,000 in silver, most of which came from the Castle Mountains.

1889--The two smelters in the district processed the Cumberland and Yellowstone ores and produced $36,355 in silver, $16,550 in gold, and 2,120,000 pounds of lead bullion. Other producers shipped several cars of lead ore, which yielded 76,459 ounces of silver.

1890--The Cumberland smelter produced 500,000 pounds of lead bullion and more than 20,000 ounces of silver from the Cumberland mine.

1891--Becoming Montana's largest lead producer, the Cumberland mine produced more than 5,000,000 pounds of lead bullion from 13,000 tons of ore. This mine produced almost half of the district's total production.

1892--The Cumberland mine produced only 300,000 pounds of lead. The district's production of 3,279,811 pounds came mostly from the Yellowstone, Jumbo, Great Eastern, and Judge mines.

1893--Except for the Cumberland mine, the district's mines were worked only by lessees, who produced several tons of lead and manganese ore from near-surface workings.

1894--Jay Anderson worked the Judge mine and produced most of the 98,094 pounds of lead recorded from Meagher County.

1895--Lessees produced most of the 383,385 pounds of lead from shallow shafts and dumps.

1896--The Jawbone Railroad was constructed to Leadboro, and considerable ore was removed from the mine dumps. The county production is given as 501,620 pounds of lead.

1897--R. A. Harlow shipped most of the district's ore from the Cumberland and Yellowstone dumps. Of the ore, 8,486 tons from the Cumberland dump yielded 5,952,776 pounds of lead and 133,542 ounces of silver. The Great Eastern also reported a production of 35 to 40 tons per day and the Powderly mined a high-grade oreshoot on Robinson Creek.

1898--The Judge mine, together with the Jumbo, accounted for most of the Meagher County production of 6,000,000 pounds of lead and 99,207 ounces of silver.

1899--The Jumbo, Judge, and Powderly mines produced nearly all of Meagher County's 7,500,000 pounds of lead and 405,630 ounces of silver.

1902--Lessees produced a small tonnage from the Judge and Jumbo mines. The Copper Bowl and Copper Kettle claims also produced 2,000 tons of iron-copper ore.

1903, 1908, 1911, 1912--A small production is reported from the district, and the Powderly is credited with being the most active mine.

1914--The district produced 12,308 pounds of zinc from oxidized ore near the Robinson area.

1916--A small tonnage of argentiferous copper ore was produced from the Milwaukee and Copper Kettle claims.

1917--The Milwaukee and Ruby adit claims produced argentiferous copper ore, and the Yellowstone shipped a few cars of lead-silver ore.

1918--Copper ore was produced from the Ruby adit. Some iron ore was also reported from the Hensley Creek area.

1919--Five mines produced 41 tons of lead ore valued at $1,706. The Ruby adit also shipped one car of copper ore.

1920--A few small lots of copper ore were reported to have been shipped from the Hensley Creek area.

1922--The Copper Bowl claim reported some development ore.

1924--James Perry shipped 892 tons of lead slag from the Cumberland dump, which assayed 0.47 ounce silver per ton, 5.9 percent lead, and 2.6 percent zinc.

1925--Lead slag was shipped from the Cumberland dump, and the Castle Mountain Mines, Inc., shipped development ore from the Homestake group of claims. This company completed considerable drifting and raising and planned to be on a milling basis in 1926.

1926--The Yellowstone, Jumbo, Homestake, and Cumberland mines were all active. The mines and dumps produced 2,412 tons of slag and 443 tons of ore, which yielded 2,070 ounces of silver, 5,532 pounds of copper, and 413,001 pounds of lead. Most of this production came from the Yellowstone property.

1927--Fourteen hundred tons of slag from the Yellowstone smelter was shipped to East Helena.

1930--No production was recorded, but the Great Western group reported 1,500 feet of tunnel work.

1933-1937--Lessees reported development work on several properties.

1939--The Belle of the Castle claim shipped a small lot of copper ore, and the Great Eastern produced 1,532 pounds of lead and 31 ounces of silver.

1943--From a small shaft on the Cumberland claim, John Oliphant produced 178 tons of ore, which yielded 94,900 pounds of lead and 1,845 ounces of silver.

1944--Lessees shipped 108 tons of ore from the Cumberland waste dump, which yielded 44,300 pounds of lead and 180 ounces of silver.

1946--The Yellowstone mine produced 48 tons of lead-silver ore, which yielded 1 ounce of gold, 193 ounces of silver, 15,000 pounds of lead, and 1,500 pounds of zinc.

1947--The district produced 2,500 pounds of lead and 53 ounces of silver.

1949--The district produced 17 tons of ore.

1950--The Silverton Mines, Inc., operated the Cumberland mine. The Yellowstone mine also operated, and together they produced 51 tons of ore, which yielded 8,000 pounds of lead, 2,000 pounds of zinc, and 294 ounces of silver.

1951--The Cumberland and Yellowstone mines produced 188 tons of ore, which yielded 108,000 pounds of lead, 6,000 pounds of zinc, and 1,885 ounces of silver.

1952--Producing 321 tons of ore, the Cumberland and Yellowstone mines were both active.

1953--A small output of lead ore was reported from the Cumberland and Great Eastern mines.

1954--Glenn Franklin produced lead ore from the Cumberland mine.

1955--From the Copper Kettle claim the D. & V. Mining Company produced 36 tons of copper ore, which assayed 2.5 percent copper and 0.5 ounce silver. The Cumberland mine also reported 89 tons of development ore.

1956--A small lot of lead ore was reported from the Cumberland mine.

1957--The H. O. Mining Company opened 80 feet of shaft, drove 100 feet of new development drift, and sank 50 feet of winze at the Cumberland mine. From these workings, the company produced 59,689 pounds of lead.

1958--The Cumberland mine was operated by the H. O. Mining Company, who shipped 1,103 tons of ore, which yielded 497,785 pounds of lead.

1959--Operating the Cumberland mine, the H. O. Mining Company and the Cumberland Mines Company produced 552 tons of ore.

1960--HOCO, Inc., operated the Blackhawk and Cumberland mines, and the Hamilton Mines Inc. operated the Yellowstone mine. The district produced 350,000 pounds of lead, nearly 4,000 ounces of silver, and 44,000 pounds of zinc. Of this amount, the Cumberland produced 340,342 pounds of lead.

1961--Operating the Cumberland mine, HOCO, Inc., shipped 559 tons of ore, which yielded 288,483 pounds of lead. The Yellowstone mine produced the rest of the district's production.

1962--HOCO, Inc., mined 653 tons of ore from the Cumberland mine; it yielded 4 ounces of gold, 4,182 ounces of silver, 266,000 pounds of lead, and 16,000 pounds of zinc.

1963--Producing until August, HOCO, Inc., operated the Cumberland mine. The Yellowstone mine, operated by George Voldseth, uncovered

a new oreshoot south of the shaft. The Copper Bowl and the Powderly were also active.

1964--The Yellowstone and Powderly mines produced more than 200 tons of lead ore and accounted for the district's production. At the Cumberland mine, considerable development was done, but as of September, no ore had been shipped.

Records of the mineral production from the various districts in Meagher County are either incomplete or not available for the years prior to 1932. Roby (1950, p. 16) compiled all the recorded production figures from 1883 to 1947 and stated that the county had produced not less than 4,250,000 ounces of silver, 29,439,740 pounds of lead, and 34,207 pounds of zinc. Virtually all of the county's lead and most of its silver production is known to have come from the Castle Mountain district, as it is the only major producer of these metals in the county. The district has also produced a considerable part of the county's zinc, copper, and manganese.

Since 1948, the district is credited with 1,444,700 pounds of lead, 22,752 ounces of silver, and 129,700 pounds of zinc. Considering that 90 percent of the lead, 50 percent of the silver, and 50 percent of the zinc recorded in the county prior to 1948 was produced from the Castle Mountain mines, the district is credited with not less than 27,940,466 pounds of lead, 2,147,752 ounces of silver, and 146,803 pounds of zinc.

ORE DEPOSITS

CLASSIFICATION

Irregular replacement bodies associated with bedding-plane fractures, jointing, and favorable beds occur adjacent to or at some distance from igneous intrusive bodies. Fissure veins also occur in the district. All of the mineable deposits were formed through hydrothermal processes.

Contact-Metasomatism

Ore deposits of the contact-metasomatic type are found along the contacts between Paleozoic limestone and intrusive dacite porphyry, and are by far the most abundant type of deposit in the district. The best example of a contact-metasomatic deposit is in the Yellowstone mine, where several oreshoots occur adjacent to a dacite porphyry sill in warped Pilgrim (Cambrian) limestone.

Metasomatic Replacement

Deposits of the metasomatic replacement type are very irregular and discontinuous, and most are confined by bedding planes. Considerable development work is required per ton of ore. The largest producer in the district, the Cumberland mine, is an excellent example of metasomatic replacement.

Fissure Veins

Production veins of the fissure type have been of minor importance in the district because of their lower grade. These veins occur in the Robinson Creek area, the Powderly veins being the most important.

Replacement Veins

Replacement deposits occur along single fissures or bedding-plane cracks in favorable limestone. Replacement of the favorable bed may persist for hundreds of feet. Referred to as jasper leads, the veins range from 2 to 10 feet in width, and jasper is the main mineral.

PARAGENESIS

The sequence of mineral deposition has been interpreted from open space fillings, mineral intergrowths, textures, and mineral replacements. Specimens were collected and studied from the various mines in the district. As the district covers a considerable area, the paragenetic relationships may differ in different mines, but it was found that the various sequences of mineral deposition are similar except for a few minor variations.

The earliest mineralization involved the introduction of iron-rich solutions, which replaced favorable limestone units in the Precambrian and lower Paleozoic sedimentary rocks. In the vicinity of Hensley and Robinson Creeks, large masses of magnetite crop out. The magnetite represents a high-temperature stage of mineralization and is found only close to intrusive bodies. Samples collected from the Belle of the Castle and Iron Chief mines show three distinct types of iron deposition. The earliest is nearly pure magnetite, which has a distinct cuneiform fracturing pattern. The second stage is pyrite, and the third again magnetite. The pyrite is cut by the second stage magnetite, and much of this magnetite is characterized by exsolution laths of hematite. Hematite veins are also believed to be related to this late magnetite deposition and probably represent the low-temperature phase of the iron mineralization. The relationship between the late-stage sulphide and magnetite-hematite deposition is not clearly understood, but may be related to the various periods of igneous activity.

Jasperoidization followed the early iron deposition. Large areas of favorable limestone were altered to jasperoid and jasper-filled open

fractures and cavities. Jasper cemented the fractured magnetite and possibly obtained its iron coloration from the magnetite. This siliceous stage continued intermittently throughout the sulphide deposition.

Following the large initial silica deposition came a normal sequence of sulphides. Pyritohedral pyrite formed in the open spaces, and massive pyrite replaced large volumes of the limestones adjacent to solution channels. Preservation of bedding was observable on a large scale but not in individual specimens. After deposition of the pyrite the ores were fractured.

Chalcopyrite was probably the next major mineral to be deposited. Bornite and an unknown mineral believed to be a variety of the enargite group were also deposited during this period, but their relationships could not be determined. Chalcopyrite is the main mineral and chalcopyrite and sphalerite began to be deposited before second-stage pyrite deposition.

Deposition of pyrite followed the copper mineralization; pyrite has partly replaced all the pre-existing minerals. Relict grains of these minerals occur as small blebs and irregular masses in the pyrite, producing a poikilitic texture.

Introduction of a minor amount of crystalline quartz interrupted the sulphide deposition. Some fracturing followed this brief silica stage.

Sphalerite was probably the next mineral to form. Crystal aggregates fromed in open fractures, and massive sphaleritie filled large voids and partly replaced the pre-existing minerals. The sphalerite contains much iron, and that in the Blackhawk area may be termed marmatite. Pyrrhotite(?) as small blebs within the sphalerite was also noticed, and it may have separated out of the solution because of an excess of iron. Chalcopyrite is also common as blebs and stringers within the sphalerite. These linear blebs show an emulsion texture suggesting exsolution. As chalcopyrite and sphalerite exsolve at about 400° C, the deposits probably formed at about this temperature (Buerger, 1934, p. 528).

Galena followed sphalerite in order of deposition, and it fills the open fractures of the early minerals. Galena has massively replaced all the pre-existing minerals and is the most abundant sulphide mineral.

No silver minerals were observed in the polished sections, but secondary native silver and silver chlorides have been reported from the Merrimac and American claims. The primary minerals that may have provided the silver are the silver sulfosalts or the tetrahedrite group.

Iron and manganese carbonates were also found at a few of the mines. They were deposited later than the sulfides, but their interrelationship is unknown, as only one specimen of primary manganese carbonate was found.

Rhodochrosite was observed at the Yellowstone mine, where it formed crusts that coated fractures in galena. The manganese deposition is believed to represent a late low-temperature stage of mineralization in a peripheral zone surrounding the sulphides. Siderite was found at several different mines and was deposited as a late gangue mineral.

Late chalcedonic quartz occurs as films, and fills small fractures in the jasper and sulphides. It is not certain, however, whether this mineral is related to the hypogene mineralization.

Enrichment is minor in the district, and it has been reported that there is little difference in the silver content of the oxidized and the primary ore. Weed and Pirsson (1896, p. 154) did state, however, that the gold content is greater in the oxidized portion of the ore in the Cumberland mine. At most of the mines some copper enrichment is indicated by the presence of chalcocite and covellite, which have replaced chalcopyrite and galena.

Oxidation has proceeded incompletely to a depth of nearly 500 feet at the Cumberland mine. Oxidation depths at the other mines differ considerably, but at most of them oxidation extends well below the 100-foot level. Zinc and copper are almost completely removed or oxidized to smithsonite, malachite, azurite, chrysocolla, cuprite, tenorite, and scarce native copper. Pyrite is converted to goethite and limonite, and the manganese minerals are completely oxidized to psilomelane and pyrolusite. Galena is oxidized to cerussite and anglesite as far down as the 460-foot level of the Cumberland mine. Cerussite has been the chief mineral mined from the district and accounts for most of the ore value.

ORESHOOTS

High-grade oreshoots occur in the mineralized zones as individual pipes and pods. The pipes are fairly continuous and conformable to the bedding. The largest pipe mined in the district lay along the hanging wall of the Cumberland mine and was mined from the surface to a point below the 500-foot level.

Where the dip changes, the pipes may be considerably enlarged. The changes in dip of the favorable limestone create swells and pinches in the pipes. Irregular pods of various sizes lay between the hanging and footwall of the Cumberland deposit. Many of these pods were only a mine set in size. Most were found near axes of change in dip of the limestone. Other pods project into the walls with no apparent control.

No suitable explanation for the oreshoots within the veins or brecciated zones can be given, owing to the inaccessibility of the workings. The position of solution cavities in limestone and the period of time required for solidification of the silica may have played an important role in localizing the oreshoots.

DEPTH OF MINERALIZATION

The type of mineralization in the Castle Mountain district suggests that the ores were deposited under mesothermal conditions. The vertical range for these deposits has not been established, but the largest mine in the district, the Cumberland, reached a zone characterized by a large ratio of pyrite to valuable sulphides at approximately 700 feet. This change in mineral content is believed to represent the bottom of the oreshoot. Other mine structures in the district are not as large as the Cumberland and are not believed to contain consistent mineral deposits for depths much greater than 700 feet (Weed and Pirsson, 1896, p. 155).

STRUCTURAL CONTROLS

It is well-known that belts of igneous intrusions are belts of ore deposits (Bateman, 1958, p. 304). Although much the greater portion of the igneous intrusive rock is nonmineralized, the ore deposits are intimately associated with such rock. The most important igneous rock type related to mineralization in the Castle Mountain district is the dacite porphyry, which has sheeted and cut the limestone beds. The porphyry, in general, merely provided deep-seated channels for the mineralizing fluids.

Local folding of the limestone seems to be the second most important factor localizing mineralization. Tension cracks, formed by the folding of the strata, provided channels along the crests and troughs of the individual folds. Where these folds are associated with intrusive dacite, ore may be found. The crescent-shaped Cumberland and Yellowstone oreshoots illustrate sulphide replacement along these flexures.

Brecciation and fracturing have also played an important part in localizing mineralization at the Blackhawk, Iron Chief, and other properties. Where such zones occur near intrusive dacite, ore may be found.

Filling of solution cavities may be more important than realized. Cavities produced by the modification of pre-existing openings, such as joints and fissures, have provided adequate space for ore deposition. At least in part, this feature is believed to have controlled the emplacement of the Corliss vein.

Open fractures have also provided space for ore deposition, the most important veins of this type being in the Robinson Creek area.

STRATIGRAPHIC CONTROLS

No one formation has proved to be more favorable than any other. Three formations seem to stand out, however, for the size and value of ore deposits contained. They are the Mission Canyon, the Jefferson, and the Pilgrim.

Several features common to the favorable units within these formations can be recognized. Where sulphides have replaced these units, the limestone tends to be slightly dolomitic. Bleaching and decomposition of the favorable units by the intrusive rock is also a noteworthy feature. The rocks are believed to have been brittle enough to crack readily under slight stress. This characteristic produced the permeability necessary for replacement.

In most mines, tough beds adjacent to the brittle units may or may not have provided impervious base rocks in the troughs of folds. The hard impervious footwall of the Cumberland and possibly the Yellowstone illustrate this feature.

The acidity of the sedimentary rocks seems to have influenced the type of deposition. Copper is the major mineral in the impure Precambrian limestone, which is much more siliceous and altered than the nearby Paleozoic limestone units that favored replacement.

ALTERATION

Both contact and hydrothermal alteration were noted in the map area. Contact metamorphism is widespread near the intrusive bodies, but hydrothermal alteration is restricted to the mineralized zones.

CONTACT METAMORPHISM

Metamorphism of the sedimentary rocks adjacent to intrusive rocks is widespread and easily recognized. The most striking effects occur between Blackhawk and Robinson where, adjacent to the diorite, the limestone has been bleached and recrystallized into marble for considerable distances. Near Blackhawk the marble is coarsely granular, and broad cleavage plates of calcite can be found.

Near the contact the marble contains agglomerations of contact minerals. Green garnet in grains and nests is commonly associated with diopside and phlogopite. Weed and Pirsson (1896, p. 93) stated that masses of brown vesuvianite are common in the nests of garnet.

The adjacent shale, especially that in the Belt Series and the Dry Creek, is baked into tough hornstone, which forms topographic ridges above the Paleozoic limestone. Argillization and possibly some silicification seem to be the dominant processes affecting these strata.

The diorite, in general, shows few effects near the chill zone. The only noticeable differences were finer groundmass and fewer phenocrysts. Weed and Pirsson (1896, p. 93) stated that chloride has replaced biotite



CASTLE MOUNTAIN DISTRICT, MONTANA

0 1 2 3 4 5 Mi.

LEGEND

Sediments and Extrusives

QUATERNARY	Alluvium:	QAI	
TERTIARY	Basalt:	L b LL	
	Rhyolite:	v r v	
	" Breccia, Tuff:	^rb^	
	Paleocene Fort Union } Livingston Formation:	L	Sandstone, grits, cgls., shale
MESOZOIC	Upper Cret. Lance }		
	" " " Dakota-Colorado-Montana:	K	Shales, Sandstones.
	Jurassic:		Shale, ss., ls.
PALEOZOIC	Carboniferous:		Sandstone, Shale Limestone
	Siluro-Devonian:		Limestone
	Cambrian:		Shale Quartzite
PROTEROZOIC	Algonkian Belt	Ab	Mainly Shale; ls., ss.
	Intrusives		
TERTIARY	Eocene?: Acid & basic dikes & sheets:		
	Granite:		Quartz Porphyry: gp
	Diorite:	d	





Scale: 1" = 2 Mi.



Base from U. S. Forest Service Maps 46-110-310 and 46-110-240

Geology by A. S. Winters -1964-





GEOLOGIC MAP OF THE CASTLE MOUNTAIN MINING DISTRICT MEAGHER COUNTY, MONTANA

EXPLANATION

SEDIMENTARY ROCKS

CENOZOIC — Quaternary
- Qd — GLACIAL DEBRIS

MESOZOIC — Cretaceous
- Kh — HELL CREEK FORMATION
- Kl — LENNEP SANDSTONE
- Kb — BEARPAW SHALES
- Kj — JUDITH RIVER FORMATION
- Ke — EAGLE SANDSTONE
- Kc — COLORADO GROUP
- Kk — KOOTENAI FORMATION

MESOZOIC — Jurassic
- J — MORRISON, SWIFT, RIERDON FORMATIONS

PALEOZOIC — Mississippian
- Ma — AMSDEN LIMESTONE
- Mb — BIG SNOWY GROUP
- Cmmc — MISSION CANYON FORMATION
- Cml — LODGEPOLE LIMESTONE

PALEOZOIC — Devonian
- D — THREE FORKS, JEFFERSON FORMATIONS

PALEOZOIC — Cambrian
- Cdc — DRY CREEK MEMBER
- Cp — PILGRIM LIMESTONE
- Cpk — PARK FORMATION
- Cm — MEAGHER LIMESTONE
- Cw — WOLSEY SHALE
- Cf — FLATHEAD SANDSTONE
- Cu — CAMBRIAN UNDIVIDED

PRECAMBRIAN — BELT SERIES
- pCp — PIEGAN GROUP

IGNEOUS ROCKS

CENOZOIC — Tertiary
- sy — SYENITE
- rh — RHYOLITE
- dp — DACITE PORPHYRY
- gr — GRANITE
- di — DIORITE

20° TRUE NORTH / MAGNETIC NORTH

SYMBOLS
- MINE
- PROSPECT
- ADIT
- CONTACT
- FAULT
- ANTICLINE
- SYNCLINE
- STRIKE AND DIP (67)

0 1/2 1 MILE SCALE

IMPORTANT MINES AND PROSPECTS

(1)	Cumberland	(12)	Iron Chief
(2)	Jumbo	(13)	California
(3)	Broadway (Lead)	(14)	Homestake
(4)	Yellowstone	(15)	Powderly
(5)	Great Eastern	(16)	Cleopatra
(6)	Great Western	(17)	Legal Tender
(7)	Copper Bowl	(18)	Judge
(8)	Golden Eagle	(19)	Blackhawk
(9)	Belle of The Castle	(20)	Alice
(10)	Milwaukee	(21)	Princess
(11)	Hidden Treasure	(22)	American



SECTION A-A'



COMPOSITE PLAN OF KNOWN LEVELS

0 10' 20'





GEOLOGY OF ADIT LEVEL



PLAN AND SECTIONS OF THE CUMBERLAND MINE

SEC. 14, T. 8 N., R. 8 E., MEAGHER COUNTY, MONTANA

500' Level adapted from Arthur Lakes 1950

dump
boiler
shaft
ore
blue lms
86
dump
rse
ore?
shop
shaft #2 caved 290'
ore
shaft #1 caved 250'
direction of major jointing
4" py
4" Fe stained clay
2" Fe oxides
0-2"
small seams of Fe stained clay
large sol. channel-some Zn ore from cave
4" Fe oxides
4' shear zone
10" Fe stained clay - contact
many small stringers of py in grey xtal lms
Fe stained sugary lms
direction of major jointing
contact
2" Fe oxides & clay
grey - green argillite & silicified lms. & dolomite
2" clay
2" py
zone of many py stringers & replacement
py & Fe oxides along jointing
0-2" py
1" gouge
1-3" gouge
grey-tan mottled lms
approx locacation of soil samples
approx contact on surface
surface cut- sugary lms near contact
gossan
2.8% Pb
Yellowstone Claim Line
N

YELLOWSTONE MINE

SEC. 11, T.8 N., R.8 E., MEAGHER COUNTY, MONTANA

0 50'

T. 9 N.

T. 8 N.

33 34 35 36

4 3 2 1

9 10 11 12

16 15 14 13

21 22 23 117 24

28 27 26 25

Monitor 3225
3317
O'Connor 3224
St Paul 3226
Legal Tender
Legal Tender 2392
Judge 2439A
Felix
Crescent 3223
Annie Maude 2424
Marie Marten 2455
Alice
Mother 2454
Caledonia 3491
Carolus 2456
Bondholder 2423
2439B
Blackhawk 2425
Alpena 3512
A.S.C. 3229
Granite Mountain 3230
Nevada 3146
Maggie 3149
Dwarf 3147
Hensle 3148
Bonanza Cr.

Golden Eagle 3492
Copper Bowl
Horseshoe Frac.
Copper Kettle
Hillbilly
Milwaukee 3051
Belle of the Castle 3066A
3736
Little Nellis 3733
Saturday 3734
Evening Bell 5677
Copper Bell 5695
Silver Bell
Antelope
Gem
First National
Hidden Treasure 2935
Iron Chief 3050
Iron Chief
Conscript
Washington
Relude (Lomar 3069A)
Kelley Fraction
Monaxx 2422
Armeda
Volunteer
Iron Reserve
Boomerang
Yellowstone 3065
Solid Silver 3219
Roland Fraction
Hendricks 3539
California 3538
Tom Pain 3098
Reserve
Little Great Eastern 3217
Bamboo Chief
Compromise
Blue Bird 3215
Homden 3216
Homden 2216
Comis 3218
Mills Bill 3222
Baltimore 6158
Homestake 3220
Hillside 3221
Phoenix
Silver Star 3131
Powder
Missouri
Crown Point
Cachie
Helena
Great Western 3068
Mary Anderson 3361
Elkhorn 3616
Contact
Hillside No 2
Graphic
Big Sarah
Fox Hound
Montana
Severance 3925
Montana Central
St Louis
Revenue
Grey Hound
Merrimac #1 3494
Grand Central 3310
Lee Lewis 3324
Ontario 3266
Mint 3267
A C M E
Bozeman
Epic
CASTLE MTN.
George 4700
Etta
Broadway 4699
Lead 2866
Jumbo Fraction
Jumbo 3265
Jefferson Davis 3268
Bone of Contention 3270
Louisville
Contact
Cumberland 2865
Stonewall Jackson 2867A
Potomac 3337
Assippee 3355
Germania
Monument 2864
Cumberland No 2
Granite Castle #2 3017A
Cornelius Shirley Placer 3336
Clar... Lode 3374
Lode 3373
Lewis and Clark
Castle Townsite
Leland Stone Placer 3249
3065A
3065B
RRG Pat.
King Add.
3173B
3065B
Black Rock 3442
3347 Castle Clay Placer
Castle Lake
Boulder Cr.
Castle Cr.
Hietman Fraction 3791
Spencer
Princess 3331
Sherman 3213
Cyclone 3792
HES 393
624915 D/C
EX Sur. 1257
1114817 D/C
1114817 D/C
1112636 D/C
R.8E.
R.9E.
Hensley Cr.
Alabaugh Cr.
To Lennep
Robinson Cr.
Forest
National
North Star 3132
Cleopatra 3362
Pike Load 2663
L.P Load 2662
Yule Dam 2664
Corral Cr.

N

1200 600 0 600 1200
Scale in Feet

CLAIM MAP OF THE CASTLE MOUNTAIN MINING DISTRICT, MEAGHER COUNTY, MONTANA