CONTACT INFORMATION
Mining Records Curator
Arizona Geological Survey
416 W. Congress St., Suite 100
Tucson, Arizona 85701
602-771-1601
http://www.azgs.az.gov
inquiries@azgs.az.gov

The following file is part of the Grover Heinrichs Mining Collection

ACCESS STATEMENT

These digitized collections are accessible for purposes of education and research. We have indicated what we know about copyright and rights of privacy, publicity, or trademark. Due to the nature of archival collections, we are not always able to identify this information. We are eager to hear from any rights owners, so that we may obtain accurate information. Upon request, we will remove material from public view while we address a rights issue.

CONSTRAINTS STATEMENT

The Arizona Geological Survey does not claim to control all rights for all materials in its collection. These rights include, but are not limited to: copyright, privacy rights, and cultural protection rights. The User hereby assumes all responsibility for obtaining any rights to use the material in excess of "fair use."

The Survey makes no intellectual property claims to the products created by individual authors in the manuscript collections, except when the author deeded those rights to the Survey or when those authors were employed by the State of Arizona and created intellectual products as a function of their official duties. The Survey does maintain property rights to the physical and digital representations of the works.

QUALITY STATEMENT

The Arizona Geological Survey is not responsible for the accuracy of the records, information, or opinions that may be contained in the files. The Survey collects, catalogs, and archives data on mineral properties regardless of its views of the veracity or accuracy of those data.

DEPARTMENT OF THE INTERIOR
UNITED STATES GEOLOGICAL SURVEY

CENOZOIC GEOLOGY AND SECTIONS OF THE IONE QUADRANGLE, NYE COUNTY, NEVADA

By
Charles J. Vitaliano

MINERAL INVESTIGATIONS
FIELD STUDIES MAP MF-255

PUBLISHED BY THE U. S. GEOLOGICAL SURVEY
WASHINGTON, D.C.
1963

MAP MF-255

DEPARTMENT OF THE INTERIOR
UNITED STATES GEOLOGICAL SURVEY

MINERAL INVESTIGATIONS
FIELD STUDIES MAP MF-255



Base map by Topographic Division
U.S. Geological Survey, 1948

Cenozoic geology by Vitaliano,
1945, 1946, and 1954





CENOZOIC GEOLOGY AND SECTIONS OF THE IONE QUADRANGLE, NYE COUNTY, NEVADA

By

Charles J. Vitaliano

SCALE 1:62 500

1 ½ 0 1 2 3 MILES

Contour interval 40 feet
Datum is mean sea level

1963

MINERAL INVESTIGATIONS
FIELD STUDIES MAP MF-255

INTERIOR—GEOLOGICAL SURVEY, WASHINGTON, D. C.—62360

For sale by U. S. Geological Survey, price 50 cents

EXPLANATION

Qal — Alluvium
Sand and gravel; restricted to valley floors; covers large areas in Ione and Indian Valleys

Qw — Hill wash and terrace gravels
Fine to coarse gravel, mainly older than alluvium. Forms prominent terraces in Reese River Valley, and subdued terraces in Ione Valley

QUATERNARY

QTv — Volcanic rocks
Andesite and trachyandesite lavas and volcanic plugs. Lavas are olivine bearing, and associated with white-, gray-, and lavender-colored pyroclastic rocks of intermediate composition. Limited to the Shoshone Mountains

TERTIARY OR QUATERNARY

EROSIONAL UNCONFORMITY (?)

QTi — Intrusive rocks
Intrusive dikes and plugs: andesites, rhyolites, and trachyandesites ranging in age from Tertiary to Quaternary

TERTIARY AND QUATERNARY

Pliocene (?) — Upper volcanic sequence

Tvuu — Upper division
Thin-bedded, dark-red to black, vesicular, aphanitic to meagerly porphyritic andesite and trachyandesite. Andesites are labradorite, hypersthene, and augite-hypersthene varieties. Trachyandesites contain olivine. Thickness approximately 300 feet

EROSIONAL UNCONFORMITY (?)

Tvt — Toyabe Quartz Latite
Light-colored pyroclastic rocks, mainly ignimbrites, breccias, and associated lavas and intrusive (?) rocks. In the Shoshone Mountains, south of Ione Canyon, dacite, dacite-breccia, and rhyodacitic ignimbrites in lower and middle members, overlain by rhyolitic vitrophyre and flows in upper member; north of Ione Canyon, rhyolite ignimbrites and sillar-type deposits with occasional rhyolite flows. In the Paradise Range, rhyolitic ignimbrites. Thickness approximately 2,000 feet

EROSIONAL UNCONFORMITY (?)

Tvo — Oddie Rhyolite
Flows, tuffs, and coarse-grained volcanic breccias, white to reddish-white to tan. Rhyolite breccia at base, overlain by rhyolite flows, lithic tuffs, and associated aphanitic facies, followed by younger rhyolite which is distinctly intrusive in many localities. Thickness approximately 1,000 feet

EROSIONAL UNCONFORMITY

Upper Miocene and lower Pliocene — Middle volcanic sequence

Tve — Esmeralda Formation
Upper division, fine-grained white to buff sandstone containing some poorly preserved fossils; very poorly represented in quadrangle. Thickness approximately 4,000 feet. Lower division coarsely fragmental, white to red breccia with associated lithic and ashy tuffs, moderately well cemented and cut by calcite veinlets; lithic fragments are composed of all earlier rocks in the area

EROSIONAL UNCONFORMITY

Lower volcanic sequence

Tvlu — Upper division
Dark-red to black, distinctly porphyritic lavas with large hornblende, biotite, pyroxene, and plagioclase phenocrysts in various combinations in aphanitic groundmass; interbedded pyroclastics. Keratophyre (albitized trachyte), trachyte, rhyolite, quartz latite, dacite, andesite, and trachyandesite. Interbedded pyroclastics make up a small percentage of this formation. Intense propylitic alteration is widespread in these and older rocks. Related intrusive rocks cut this and older rocks, but not the youngest andesites in this formation. Thickness approximately 1,000 feet

EROSIONAL UNCONFORMITY (?)

Tvll — Lower division
Generally white to purple lavas with associated pyroclastics. Rhyolite, rhyolite tuff, trachyte, trachyte tuff, and quartz latite; sheared and contorted, with development of low-grade metamorphic textures and minerals. Presence of many quartz veinlets suggests silicification. Interbedded sediments and reworked tuffs occur just beyond the southern boundary of quadrangle. Thickness approximately 1,500 feet

TERTIARY

Unconformity

pTu — Pre-Tertiary rocks, undifferentiated
(See Silberling, N. J., 1959, U.S. Geol. Survey Prof. Paper 322, 67 p.)

Contact
Dashed where inferred

U / D — Fault
Dashed where inferred, dotted where covered
U, upthrown side; D, downthrown side