



CONTACT INFORMATION
Mining Records Curator
Arizona Geological Survey
3550 N. Central Ave, 2nd floor
Phoenix, AZ, 85012
602-771-1601
<http://www.azgs.az.gov>
inquiries@azgs.az.gov

The following file is part of the Cambior Exploration USA Inc. records

ACCESS STATEMENT

These digitized collections are accessible for purposes of education and research. We have indicated what we know about copyright and rights of privacy, publicity, or trademark. Due to the nature of archival collections, we are not always able to identify this information. We are eager to hear from any rights owners, so that we may obtain accurate information. Upon request, we will remove material from public view while we address a rights issue.

CONSTRAINTS STATEMENT

The Arizona Geological Survey does not claim to control all rights for all materials in its collection. These rights include, but are not limited to: copyright, privacy rights, and cultural protection rights. The User hereby assumes all responsibility for obtaining any rights to use the material in excess of "fair use."

The Survey makes no intellectual property claims to the products created by individual authors in the manuscript collections, except when the author deeded those rights to the Survey or when those authors were employed by the State of Arizona and created intellectual products as a function of their official duties. The Survey does maintain property rights to the physical and digital representations of the works.

QUALITY STATEMENT

The Arizona Geological Survey is not responsible for the accuracy of the records, information, or opinions that may be contained in the files. The Survey collects, catalogs, and archives data on mineral properties regardless of its views of the veracity or accuracy of those data.

05/12/87

ARIZONA DEPARTMENT OF MINES AND MINERAL RESOURCES FILE DATA

① Dripping Spgs. Dist.

PRIMARY NAME: RATTLER MINE , *See Also Rents Group and Dripping Springs*

ALTERNATE NAMES:

TROY COPPER CO. PROPERTY

(Columbia) Mines

Also Check Buckeye Mine

PINAL COUNTY MILS NUMBER: 134

LOCATION: TOWNSHIP 3 S RANGE 14 E SECTION 26 QUARTER C
LATITUDE: N 33DEG 08MIN 31SEC LONGITUDE: W 110DEG 53MIN 08SEC
TOPO MAP NAME: SONORA - 7.5 MIN


CURRENT STATUS: PAST PRODUCER

COMMODITY:

COPPER

BIBLIOGRAPHY:

ADMMR RATTLER MINE FILE
ADMMR TROY MINE FILE
RANSOME, F.L., RAY FOLIO 1923, P. 22
ADMMR U FILE PINAL CU21 (USBM NO 463.2/15083)
USGS MAP GQ 1021; 1971



Sonora 7.5'



ROAD CLASSIFICATION

Heavy-duty ——— Light-duty
Medium-duty - - - Unimproved dirt

○ State Route

SONORA, ARIZ.

NW/4 RAY 15' QUADRANGLE
N3307.5—W11052.5/7.5

1964

AMS 3850 III NW—SERIES V898

(HAYDEN)
3850 III SE

2.
PL

* GENERAL REFERENCES

REFERENCE 1	P1 < USGS GQ 1021, 1971
REFERENCE 2	P2 < AGENT-USDM FILE DATA
REFERENCE 3	P3 < ADNR TRAY AND RATTLE MINE FILES
REFERENCE 4	P4 < RANSOME, F-L. USGS: GEOLOGIC ATLAS OF THE U.S., RAY FOLIO, 1922 MAP AND P 22.

DIOL PRODUCTION INCLUDED UNDER TEST IN AGENT-USER FILE DATA

uncls 134

U.S. CRIB-SITE FORM

RECORD IDENTIFICATION

RECORD NUMBER B10 <_____> RECORD TYPE B20 <X.1.M> DEPOSIT NUMBER B40 <_____>
REPORT DATE G1 <6.8.70.3> INFORMATION SOURCE B30 <1.2> FILE LINK IDENT. B50 <USGM-004-0211003>
YR. MO.
REPORTER(SUPERVISOR) G2 <GEST. DON E _____>
(last, first, middle initial) (last, first, middle initial)
REPORTER AFFILIATION G4 <ABENT _____> SITE NAME A10 <RATTLE MINE>
SYNONYMS A11 <TROY, TROY - MANHATTAN, TROY-ARIZONA>

LOCATION

WINING DISTRICT/AREA	A50	< DRIPPING SPRINGS DISTRICT			
COUNTY	A60	< PINAL	STATE	A80	< A.Z
PHYSIOGRAPHIC PROV	A63	< I.P.M.			COUNTRY
DRAINAGE AREA	A62	< 1,50,50.1 00.W. LOWER COLORADO			
QUADRANGLE NAME	A90	< SONORA			LAND STATUS
SECOND QUAD NAME	A92	< (1, 1, 9, 6, 4)			A64
ELEVATION	A107	< (3, 7, 50, W.F.T.)			QUADRANGLE SCALE
					A100
					SECOND QUAD SCALE
					A91

JTM
NORTHING A120<3.6.6.6.8.4.0.>
EASTING A130<5.1.0.7.0.0.>
ZONE NUMBER A110<+1.2>

*ACCURACY

ACCURATE ACC (circle)

ESTIMATED EST CENTER OF GRAVITY OF 4 ADITS

GEODETIC
 *LATITUDE A70 < _ _ _ _ N >
 *LONGITUDE A80 < _ _ _ _ W >

CADAstral

TOWNSHIP(S) A77<0.03.S.:N.:E.:S.:W.:> RANGE(S) A78<0.14.E.:N.:S.:W.:>
SECTION(S) A79<26.:>
SECTION FRACTION(S) A76<>
MERIDIAN(S) A81<GILA AND SALT RIVER>

POSITION FROM NEAREST PROMINENT LOCALITY A82 < 1 1/2 MILBS NE OF TIGER MOUNTAIN >
LOCATION COMMENTS A83 < 1/4 MILE W OF PINAL-GILA COUNTY LINE

ESSENTIAL INFORMATION
ESSENTIAL SOMETIMES OR HIGHLY RECOMMENDED

GENERAL COMMENTS GEN _____

INVENTORY OF THE RATTLER MINE EQUIPMENT
AS OF FEBRUARY 1, 1945

3	Redwood tanks	\$150.00
2	Steel tanks	100.00
1	2" deep well pump	150.00
1	1-1/2" Pressure pump	100.00
1	2" Pressure pump	200.00
1	1-1/2 H.P. Fairbanks Morse Engine	100.00
1	1-1/2 H.P. White Engine #340540	100.00
1	8 H.P. Stover Engine #94997	200.00
1	30 H.P. Foss Hoist & Cable	500.00
1	Grizzly complete	50.00
1	Water tank 450 Gals.	75.00
1	D.C. elec. Generator & Board #221551 Type B.C.	150.00
1	Light plant & batteries, Engine #17536	200.00
1	Model A. Ford Truck 1-1/2 ton	200.00
1	International Dump Truck 1-1/2	150.00
1	V8 Ford pick-up '36	350.00
1	Blacksmith forge & tools	150.00
	Mining tools, Misch.	250.00
	Hand tools, Misch.	250.00
3	Mine cars 18" gauge	100.00
	1-1/2 Ton mine rails	60.00
1	Tent house	150.00
1	Tool house	100.00
1	3-room house	350.00
1	Air receiver	100.00
3	Ore buckets	100.00
1	Shave wheel	35.00
	Leaching plant already installed	
5	Wooden tanks for precipitation	150.00
1	Furnace Burner	25.00
1	Circulating pump & engine #B-51147	50.00
	Foundation for tanks and labor	250.00
	Labor 2 men installing leaching plant	650.00

\$5,545.00

374-5389

~~Simple best way 1" = 200'~~

~~Don't hole for map for notes to Peter Berger~~

call McKenna

- Call Bob Golden

Call Gila Indians River Indians

call Mike Siakoski

~~JS, R 7B Sec. 35, 36~~

Air photos for Two Area, Diggins, Lewis Dist - Paul & Bela Counties, AZ

do sectional report

58

T 35 R 0E

Sec. 31

A2-78 DC

2-3-68

2-2-82

9" x 9" contact print + \$12.00 color

left of center

adjoints 2-3-68 on the west)

balance

per order
vol

\$ 1

36 x 36 = \$115.00

40 x 40 = \$118.00

all

enlargement include

color balance - not need
the \$12.00

Need to check
ownership - Globe, 1:100,000

DEP_NAME: TROY RANCH PROSPECT
STATE_CODE: AZ
COUNTY: PINAL
COMMODITY: CU MO
MINE_DIST: DRIPPING SPRINGS DISTRICT
TOWNSHIP: 003S
RANGE: 014E
SECTION: 23
SECT_FRACT: S/
POSITION: 1/2 MI N OF RATTLER MINE
LOC_COMM:
MINE_TYPE: S
PROD_SIZE: N
EXPL_COMM: STAKED BY C. MOORE OF GLOBE IN LATE 1960'S. TESTED
BY SHALLOW
DEP_TYPES: PORPHYRY COPPER
DEP_SHAPE:
OBY_STRIKE:
OBY_DIP:
DEP_DESCR:
WKGS_COMM:
AGE_HOST: CRET
HOST_ROCK: GRANODIORITE RHYODACITE PORPHYRY DIKES
AGE_IGN: CRET
IGN_ROCK: GRANODIORITE RHYODACITE PORPHYRY DIKES
AGE_MINER: CRET ABOU
MINERALOGY: QUARTZ SERICITE PYRITE K-SPAR BIOTITE
ORE_MATS: CHALCOPYRITE MOLYBDENITE PYRITE BORNITE CU
OXIDES
ORE_CONTR: E-W VEINS DIKES
SIG_ALTER: QUARTZ-SERICITE-PYRITE VEINS CUTTING
K-SPAR-BIOTITE-CHALCOPYR
FORM_AGE:
FORM_NAME:
GEOL_COMM:
GEOL_ENV:
GEOL_NOTES:
GEN_COMMS: THIS REPORT WAS TAKEN FROM RECORD M030475 OF JAN
WILT IN MOLY
REFERENCES: CORNWALL ET AL 1971 USGS MAP GQ-1021|KEITH S.B.
UNPUB DA
ALT_NAME: MARY ALICE CLAIMS 8-21 NEARBY CLAIMS 2 3 4 7
QUADRANGLE: SONORA
SCALE: 24000
ALTITUDE:
YR_FST_PRD:
YR_LST_PRD:
LAST_OPER:
CUM_PROD:
CUM_P_COMM:
COMMENTS:
LATITUDE: 33-08-53N
LONGITUDE: 110-53-03W
UTM_N: 3667500
UTM_E: 510040
UTM_ZONE: +12
COUNTRY: US
INFO_SRCE: 1
REPORTER: PETERSON JOCELYN A
REP_AFFIL: USGS
REP_DATE: 83 04
UPDATE:
REC_TYPE: X1M
REC_NO: M030475

DEP_NAME: BLUE COPPER MINE
STATE_CODE: AZ
COUNTY: PINAL
COMMODITY: CU AGAU
MINE_DIST: DURHAM-SUIZO DISTRICT
TOWNSHIP: 008S
RANGE: 012E
SECTION: 17
SECT_FRACT: SW
POSITION: 1 MILE SE OF NORTH HILL; NORTHERN MOST OF THE
DURHAM HILLS
LOC_COMM: UTM IS WHERE 2 ADITS ARE SIDE BY SIDE IN SW 1/4
SECTION 17
MINE_TYPE: B
PROD_SIZE: S
EXPL_COMM: OPERATORS INCLUDED DRAKE ENTERPRIZES GUZMAN AND
SCHWARTZ '61
DEP_TYPES: VEIN/SHEAR ZONE
DEP_SHAPE: IRREGULAR
OBY_STRIKE: NW
OBY_DIP: 10-
DEP_DESCR: SOUTH ORE BODY 1200 FT LONG WIDTH IN RANGE OF 75 TO
150 FT-AL
WKGS_COMM: 2 SHAFTS OLD; TUNNEL DRIVEN WESTWARD INTO HILL.
SHALLOW AN
AGE_HOST: PREC
HOST_ROCK: SCHIST GRANITE
AGE_IGN: CRET-PALEO
IGN_ROCK: QUARTZ DIORITE APLITE DIKES
AGE_MINER: CRET-PALEO
MINERALOGY: ALONG FRACTURES
ORE_MATS: CHALCOCITE CHRYSOCOLLA MALACHITE AZURITE
CHALCOPYRITE
ORE_CONTR: SCHIST- GRANITE CONTACT FRACTURES WITH VEINLETS
SIG_ALTER: KAOLINIZATION AND SILICIFICATION
FORM_AGE: PREC
FORM_NAME: PINAL SCHIST
GEOL_COMM: ORE HAS HIGHALUMNA CONTENT
GEOL_ENV:
GEOL_NOTES:
GEN_COMMS:
REFERENCES: ADMR BLUE COPPER FILE|C.F. BARTER 1962. GEOLOGY
OF THE OML
ALT_NAME: DURHAM HILLS MINE BLUEHILL COPPER MINE BIG BULL
MINE BUSY B
QUADRANGLE: TORTOLITA MOUNTAINS (1959)
SCALE: 62500
ALTITUDE: 2720 FT
YR_FST_PRD: 1948
YR_LST_PRD: 1962
LAST_OPER: MCFARLAND AND MULLINGER 1966
CUM_PROD:
CUM_P_COMM:
COMMENTS:
LATITUDE: 32-43-51N
LONGITUDE: 111-07-56W
UTM_N: 3621250
UTM_E: 487600
UTM_ZONE: +12
COUNTRY: US
INFO_SRCE: 2
REPORTER: GEST DON E.
REP_AFFIL: ABGMT
REP_DATE: 82 03
UPDATE:
REC_TYPE: X1M
REC_NO: M899898

DEP_NAME: BLUESTAR MINE

CUM_PROD:
CUM_P_COMM:
COMMENTS:
LATITUDE: 32-55-19N
LONGITUDE: 116-50-48W
UTM_N: 3642453
UTM_E: 514336
UTM_ZONE: +11
COUNTRY: US
INFO_SRCE: 1
REPORTER: GEST DON E.
REP_AFFIL: ABGMT
REP_DATE: 82 04
UPDATE:
REC_TYPE: X1M
REC_NO: M241198

UTM_E: 507170
UTM_ZONE: +12
COUNTRY: US
INFO_SRCE: 1
REPORTER: GEST DON E.
REP_AFFIL: ABGMT
REP_DATE: 82 03
UPDATE:
REC_TYPE: X1M
REC_NO: M899899

DEP_NAME: ALICE MINE
STATE_CODE: AZ
COUNTY: PINAL
COMMODITY: CU AU AG
MINE_DIST: DRIPPING SPRINGS DISTRICT
TOWNSHIP: 003S
RANGE: 014E
SECTION: 33
SECT_FRACT: NO
POSITION: 1 MILE W OF TIGER MTN IN HACKBERRY GULCH
LOC_COMM: LOCATED AND NAMED ON QUADRANGLE. POSSIBLY THE PRATT
TUNNEL IS
MINE_TYPE: U
PROD_SIZE: S
EXPL_COMM: MINE DID NOT PAY EXPENSES CLOSED QUICKLY SOMETIME
BEFORE 192
DEP_TYPES: VEIN/SHEAR ZONE
DEP_SHAPE:
OBY_STRIKE: NE
OBY_DIP:
DEP_DESCR:
WKGS_COMM: INCLINED SHAFT 45 DEG. THREE LEVELS (1923) 400 FT
IN 1916. C
AGE_HOST: PREC
HOST_ROCK: QUARTZITE LIMESTONE
AGE_IGN: CRET-PALEO
IGN_ROCK: RHYODACITIC PORPHYRY DIKE
AGE_MINER: CRET-PALEO
MINERALOGY:
ORE_MATS: CHALCOPYRITE NATIVE GOLD NATIVE SILVER
ORE_CONTRL: FISSURES AND VEINS NEAR RHYODACITE DIKE
SIG_ALTER:
FORM_AGE: PREC|PREC
FORM_NAME: MESCAL LIMESTONE|TROY QUARTZITE
GEOL_COMM:
GEOL_ENV:
GEOL_NOTES:
GEN_COMMS:
REFERENCES: RANSOME F.L. 1923. USGS GEOLOGIC ATLAS OF THE
U.S. RAY FOL
ALT_NAME: TROY PROPERTY PRATT TUNNEL
QUADRANGLE: SONORA (1964)
SCALE: 24000
ALTITUDE: 3080 FT
YR_FST_PRD: 1900'S
YR_LST_PRD: 1910'S
LAST_OPER:
CUM_PROD:
CUM_P_COMM:
COMMENTS:
LATITUDE: 33-07-56N
LONGITUDE: 110-55-23W
UTM_N: 3665750

DEP_NAME: BUCKEYE MINE GROUP
STATE_CODE: AZ
COUNTY: PINAL
COMMODITY: CU AG AU
MINE_DIST: DRIPPING SPRINGS DISTRICT
TOWNSHIP: 003S
RANGE: 014E
SECTION: 27
SECT_FRACT: SW
POSITION: ONE HALF MILE NW OF TIGER MOUNTAIN
LOC_COMM: UTM IS LOCATION OF BUCKEYE SHAFT OTHER SHAFTS IN
GROUP ARE NOR
MINE_TYPE: U
PROD_SIZE: S
EXPL_COMM: OPERATORS INCLUDED TROY ARIZONA COPPER TROY
MANHATTAN COMPAN
DEP_TYPES: VEIN/SHEAR ZONE
DEP_SHAPE:
OBY_STRIKE: WNW
OBY_DIP:
DEP_DESCR:
WKGS_COMM: 3 LEVELS AND 150 FT SHAFT IN 1923 ONE LEVEL 1000
FT LONG. SHA
AGE_HOST: PREC
HOST_ROCK: DIABASE LIMESTONE QUARTZITE
AGE_IGN: CRET-PALEO
IGN_ROCK: GRANODIORITE RHYODACITE DIKES
AGE_MINER: CRET-PALEO
MINERALOGY: QUARTZ SERICITE
ORE_MATS: CHALCOPYRITE CHALCOCITE NATIVE SILVER NATIVE GOLD
ORE_CONTRL:
SIG_ALTER:
FORM_AGE: PREC|PREC
FORM_NAME: MESCAL LIMESTONE|TROY QUARTZITE
GEOL_COMM: MINERALIZATION ASSOCIATED WITH GRANODIORITE
INTRUSION OPENED
GEOL_ENV:
GEOL_NOTES:
GEN_COMMS:
REFERENCES: USGS GQ 1021 1971|ABGMT-USBM FILEDATA|ADMR TROY
AND RATTLER
ALT_NAME: CLIMAX TROY PROPERTY TROY-ARIZONA
QUADRANGLE: SONORA (1964)
SCALE: 24000
ALTITUDE: 3600 FT
YR_FST_PRD: 1903
YR_LST_PRD: 1951
LAST_OPER:
CUM_PROD:
CUM_P_COMM:
COMMENTS:
LATITUDE: 33-08-14N
LONGITUDE: 110-54-40W
UTM_N: 3666320
UTM_E: 508290
UTM_ZONE: +12
COUNTRY: US
INFO_SRCE: 1
REPORTER: GEST DON E.

REP_AFFIL: ABGMT
REP_DATE: 82 03
UPDATE:
REC_TYPE: X1M
REC_NO: M899870

DEP_NAME: NINETY-ONE MINE
STATE_CODE: AZ
COUNTY: PINAL
COMMODITY: CU PB MO V
MINE_DIST: DRIPPING SPRINGS DISTRICT
TOWNSHIP: 003S
RANGE: 014E
SECTION: 27
SECT_FRACT: SW
POSITION: 1/2 MI SW OF TROY SITE 1/4 MI NE OF BUCKEYE MINE
LOC_COMM:
MINE_TYPE: U
PROD_SIZE: S
EXPL_COMM:
DEP_TYPES: REPLACEMENT
DEP_SHAPE: LENTICULAR BUNCHES
OBY_STRIKE:
OBY_DIP:
DEP_DESCR:
WKGS_COMM: SHAFT WAS APPARENTLY 150 FT DEEP WITH 3 LEVELS
AGE_HOST: PREC
HOST_ROCK: LIMESTONE DIABASE QUARTZITE
AGE_IGN:
IGN_ROCK:
AGE_MINER: LCRET-TERT
MINERALOGY:
ORE_MATS: OXIDE MINERALS OF CU PB MO & V WULFENITE
ORE_CONTRL: ALONG BEDDING PLANES IN LIMESTONE AS INCLUSIONS
IN DIABASE
SIG_ALTER:
FORM_AGE: PREC
FORM_NAME: MESCAL LIMESTONE & DRIPPING SPRING QUARTZITE OF
APACHE GROUP
GEOL_COMM: WULFENITE IS AN OXIDATION PRODUCT OCCURRING IN
JOINTS IN QUART
GEOL_ENV:
GEOL_NOTES:
GEN_COMMS: THIS RECORD WAS TAKEN FROM RECORD M000385 OF JAN
WILT IN MOLY
REFERENCES: RANSOME 1923 USGS FOLIO 217|ABM FILE
DATA|CORNWALL ET AL
ALT_NAME:
QUADRANGLE: SONORA
SCALE: 24000
ALTITUDE: 3795 FT
YR_FST_PRD:
YR_LST_PRD:
LAST_OPER:
CUM_PROD: 1945-1955|1945-1955|1945-1955
CUM_P_COMM:
COMMENTS:
LATITUDE: 33-08-18N
LONGITUDE: 110-54-28W
UTM_N:
UTM_E:
UTM_ZONE:
COUNTRY: US
INFO_SRCE: 1
REPORTER: PETERSON JOCELYN A
REP_AFFIL: USGS
REP_DATE: 83 04
UPDATE:
REC_TYPE: X1M
REC_NO: M000385

DEP_NAME: RATTLER MINE
STATE_CODE: AZ
COUNTY: PINAL
COMMODITY: CU AG AU
MINE_DIST: DRIPPING SPRINGS DISTRICT
TOWNSHIP: 003S
RANGE: 014E
SECTION: 26
SECT_FRACT:
POSITION: 1 1/2 MILES NE OF TIGER MOUNTAIN
LOC_COMM: 1/4 MILE W OF PINAL-GILA COUNTY LINE
MINE_TYPE: U
PROD_SIZE: S
EXPL_COMM: 13 UNPATENTED CLAIMS IN 1956. ENTIRE GROUP INTO
GILA CTY INC
DEP_TYPES: VEIN/SHEAR ZONE
DEP_SHAPE: TABULAR/IRREGULAR
OBY_STRIKE:
OBY_DIP:
DEP_DESCR:
WKGS_COMM: 3 TUNNELS AT LEAST ONE ADIT 160 FT DEEP IN 1944
AGE_HOST: PREC
HOST_ROCK: LIMESTONE QUARTZITE
AGE_IGN: CRET-PALEO
IGN_ROCK: RHYOLITE OR RHYODACITE PORPHYRY
AGE_MINER: CRET-PALEO
MINERALOGY: QUARTZ SERICITE
ORE_MATS: CHALCOPYRITE CHALCOCITE BORNITE MALACHITE NATIVE
SILVER N
ORE_CONTRL: FISSURES AND VEINS NEAR RHYOLITE DIKES
SIG_ALTER:
FORM_AGE: PREC|CAMB
FORM_NAME: MESCAL LIMESTONE DRIPPING SPRINGS QUARTZITE TROY
QUARTZITE|
GEOL_COMM: PROBABLY ASSOCIATED WITH GRANODIORITE INTRUSION OR
AT LEAST
GEOL_ENV:
GEOL_NOTES:
GEN_COMMS:
REFERENCES: USGS GQ 1021 1971|ABGMT-USBM FILE DATA|ADMR TROY
AND RATTLE
ALT_NAME: TROY TROY-MANHATTAN TROY-ARIZONA
QUADRANGLE: SONORA (1964)
SCALE: 24000
ALTITUDE: 3750 FT
YR_FST_PRD: 1903
YR_LST_PRD: 1940'S
LAST_OPER: INSPIRATION COPPER COMPANY 1965
CUM_PROD:
CUM_P_COMM:
COMMENTS:
LATITUDE: 33-08-31N
LONGITUDE: 110-53-07W
UTM_N: 3666840
UTM_E: 510700
UTM_ZONE: +12
COUNTRY: US
INFO_SRCE: 1
REPORTER: GEST DON E.
REP_AFFIL: ABGMT
REP_DATE: 82 03
UPDATE:
REC_TYPE: X1M
REC_NO: M899871

DEP_NAME: TROY RANCH PROSPECT
 STATE_CODE: AZ
 COUNTY: PINAL
 COMMODITY: CU MO
 MINE_DIST: DRIPPING SPRINGS DISTRICT
 TOWNSHIP: 003S
 RANGE: 014E
 SECTION: 23
 SECT_FRACT: S/
 POSITION: 1/2 MI N OF RATTLER MINE
 LOC_COMM:
 MINE_TYPE: S
 PROD_SIZE: N
 EXPL_COMM: STAKED BY C. MOORE OF GLOBE IN LATE 1960'S. TESTED
 BY SHALLOW
 DEP_TYPES: PORPHYRY COPPER
 DEP_SHAPE:
 OBY_STRIKE:
 OBY_DIP:
 DEP_DESCr:
 WKGS_COMM:
 AGE_HOST: CRET
 HOST_ROCK: GRANODIORITE RHYODACITE PORPHYRY DIKES
 AGE_IGN: CRET
 IGN_ROCK: GRANODIORITE RHYODACITE PORPHYRY DIKES
 AGE_MINER: CRET ABOU
 MINERALOGY: QUARTZ SERICITE PYRITE K-SPAR BIOTITE
 ORE_MATS: CHALCOPYRITE MOLYBDENITE PYRITE BORNITE CU
 OXIDES
 ORE_CONTRL: E-W VEINS DIKES
 SIG_ALTER: QUARTZ-SERICITE-PYRITE VEINS CUTTING
 K-SPAR-BIOTITE-CHALCOPYR
 FORM_AGE:
 FORM_NAME:
 GEOL_COMM:
 GEOL_ENV:
 GEOL_NOTES:
 GEN_COMMS: THIS REPORT WAS TAKEN FROM RECORD M030475 OF JAN
 WILT IN MOLY
 REFERENCES: CORNWALL ET AL 1971 USGS MAP GQ-1021 KEITH S.B.
 UNPUB DA
 ALT_NAME: MARY ALICE CLAIMS 8-21 NEARBY CLAIMS 2 3 4 7
 QUADRANGLE: SONORA
 SCALE: 24000
 ALTITUDE:
 YR_FST_PRD:
 YR_LST_PRD:
 LAST_OPER:
 CUM_PROD:
 CUM_P_COMM:
 COMMENTS:
 LATITUDE: 33-08-53N
 LONGITUDE: 110-53-03W
 UTM_N: 3667500
 UTM_E: 510040
 UTM_ZONE: +12
 COUNTRY: US
 INFO_SRCE: 1
 REPORTER: PETERSON JOCELYN A
 REP_AFFIL: USGS
 REP_DATE: 83 04
 UPDATE:
 REC_TYPE: X1M
 REC_NO: M030475

DEP_NAME: BLUE COPPER MINE
 STATE_CODE: AZ
 COUNTY: PINAL
 COMMODITY: CU AGAU
 MINE_DIST: DURHAM-SUIZO DISTRICT
 TOWNSHIP: 008S
 RANGE: 012E
 SECTION: 17
 SECT_FRACT: SW
 POSITION: 1 MILE SE OF NORTH HILL; NORTHERN MOST OF THE
 DURHAM HILLS
 LOC_COMM: UTM IS WHERE 2 ADITS ARE SIDE BY SIDE IN SW 1/4
 SECTION 17
 MINE_TYPE: B
 PROD_SIZE: S
 EXPL_COMM: OPERATORS INCLUDED DRAKE ENTERPRIZES GUZMAN AND
 SCHWARTZ '61
 DEP_TYPES: VEIN/SHEAR ZONE
 DEP_SHAPE: IRREGULAR
 OBY_STRIKE: NW
 OBY_DIP: 10-
 DEP_DESCr: SOUTH ORE BODY 1200 FT LONG WIDTH IN RANGE OF 75 TO
 150 FT-AL
 WKGS_COMM: 2 SHAFTS OLD; TUNNEL DRIVEN WESTWARD INTO HILL.
 SHALLOW AN
 AGE_HOST: PREC
 HOST_ROCK: SCHIST GRANITE
 AGE_IGN: CRET-PALEO
 IGN_ROCK: QUARTZ DIORITE APLITE DIKES
 AGE_MINER: CRET-PALEO
 MINERALOGY: ALONG FRACTURES
 ORE_MATS: CHALCOCITE CHRYSOCOLLA MALACHITE AZURITE
 CHALCOPYRITE
 ORE_CONTRL: SCHIST- GRANITE CONTACT FRACTURES WITH VEINLETS
 SIG_ALTER: KAOLINIZATION AND SILICIFICATION
 FORM_AGE: PREC
 FORM_NAME: PINAL SCHIST
 GEOL_COMM: ORE HAS HIGHALUMNA CONTENT
 GEOL_ENV:
 GEOL_NOTES:
 GEN_COMMS:
 REFERENCES: ADMR BLUE COPPER FILE C.F. BARTER 1962. GEOLOGY
 OF THE OWL
 ALT_NAME: DURHAM HILLS MINE BLUEHILL COPPER MINE BIG BULL
 MINE BUSY B
 QUADRANGLE: TORTOLITA MOUNTAINS (1959)
 SCALE: 62500
 ALTITUDE: 2720 FT
 YR_FST_PRD: 1948
 YR_LST_PRD: 1962
 LAST_OPER: MCFARLAND AND MULLINGER 1966
 CUM_PROD:
 CUM_P_COMM:
 COMMENTS:
 LATITUDE: 32-43-51N
 LONGITUDE: 111-07-56W
 UTM_N: 3621250
 UTM_E: 487600
 UTM_ZONE: +12
 COUNTRY: US
 INFO_SRCE: 2
 REPORTER: GEST DON E.
 REP_AFFIL: ABGMT
 REP_DATE: 82 03
 UPDATE:
 REC_TYPE: X1M
 REC_NO: M899898

DEP_NAME: BLUESTAR MINE

DEP_NAME: RENFRO GROUP
 MINE_DIST: DRIPPING SPRINGS DISTRICT
 COUNTY: GILA
 COMMODITY: CU AG AU
 TOWNSHIP: 003S
 RANGE: 014E
 SECTION: 25
 SECT_FRACT: NW
 POSITION: 1 MILE NNW OF MANHATTAN MOUNTAIN (4417 FT) DRIPPING
 SPRINGS MOUNTAINS
 LOC_COMM: 1/4 MILE E OF GILA-PINAL COUNTY BOUNDARY 1 1/4 MILES E OF
 TROY RANCH
 MINE_TYPE: U
 PROD_SIZE: S
 EXPL_COMM: PINAL DEVELOPMENT COMPANY AND F. M. POOL WERE PAST
 OPERATORS. 47 CLAIMS IN 1922
 DEP_TYPES: REPLACEMENT/SHEAR ZONE
 DEP_SHAPE:
 OBY_STRIKE:
 OBY_DIP:
 DEP_DESCR: 15 FT THICK ORE HORIZONON LIMESTONE
 WKGS_COMM: MAIN TUNNEL 1600 FT. LONG DEPTH 600 FT (MINES
 HANDBOOK 1922) ADMR REPORT (1907) DESCRIBES VARIOUS SMALL SHAFTS;
 220 FT OF WORKINGS ON THE GOLDEN EAGLE A 150 FT TUNNEL AND A 43 FT.
 SHAFT IN GRANITE.
 AGE_HOST: PREC
 HOST_ROCK: DIABASE LIMESTONE
 AGE_IGN: LCRET-TERT
 IGN_ROCK: RHYODACITE PORPHYRY
 AGE_MINER: LCRET-TERT
 MINERALOGY:
 ORE_MATS: CUPRITE CHRYSOCOLLA
 ORE_CONTRL: DIABASE-LIMESTONE CONTACT DIABASE UNDERLIES
 LIMESTONE ALSO ALONG SMALL FAULT FISSURES
 SIG_ALTER:
 FORM_AGE: PREC
 FORM_NAME: MESCAL LIMESTONE
 GEOL_COMM: LCRET RATTLERGRANODIORITE INTRUSION IS 1/8 MILE W OF
 MINE AND MINERALIZATION AND/OR FISSURING IS PROBABLY RELATED TO
 THIS INTRUSION. RHYODACITE POSTDATES INTRUSION DOWNTHROWN SIDE
 IS THE SOUTHERN SIDE ON MOST OF THE FRACTURES IN THE AREA
 GEOL_ENV:
 GEOL_NOTES:
 GEN_COMMS:
 REFERENCES: ADMR RENFRO GROUP FILE|ABGMT-USBM FILE DATA|USGS
 GEOLOGIC QUADRANGLE GQ-10218 1971|REED W. H. MINES HANDBOOK
 VOLUME XV 1922 P364
 ALT_NAME: ROMAN EAGLE SHAFT
 QUADRANGLE: SONORA (1964)
 SCALE: 24000
 ALTITUDE: 3400 FT
 YR_FST_PRD: 1913
 YR_LST_PRD: 1918
 LAST_OPER:
 CUM_PROD:
 CUM_P_COMM:
 COMMENTS:
 LATITUDE: 33-08-51N
 LONGITUDE: 110-52-39W
 UTM_N: 36674540
 UTM_E: 5114240
 UTM_ZONE: +12
 INFO_SRCE: 1
 REPORTER: GEST DON E.
 REP_AFFIL: ABGMT
 REP_DATE: 82 05
 UPDATE:
 REC_TYPE: X1M
 REC_NO: M241244

REC_NO: M000500

DEP_NAME: COLUMBIA MINE
MINE_DIST: DRIPPING SPRINGS DISTRICT
COUNTY: GILA
COMMODITY: CU AG AU
TOWNSHIP: 003S
RANGE: 014E
SECTION: 25
SECT_FRACT:
POSITION: 2 MILES EAST OF TROY
LOC_COMM:
MINE_TYPE: U
PROD_SIZE: S
EXPL_COMM:
DEP_TYPES: VEIN
DEP_SHAPE: LINEAR
OBY_STRIKE: N 89E
OBY_DIP: 75
DEP_DESCR:
WKGS_COMM:
AGE_HOST: PREC
HOST_ROCK: LIMESTONE
AGE_IGN: ETERT
IGN_ROCK: QUARTZ DIORITE PORPHYRY
AGE_MINER: ETERT
MINERALOGY:
ORE_MATS: CU CARBONATES CUPRITE PYRITE CHALCOPYRITE
ORE_CONTRL:
SIG_ALTER:
FORM_AGE:
FORM_NAME:
GEOL_COMM:
GEOL_ENV:
GEOL_NOTES:
GEN_COMMS:
REFERENCES: ROSS C. P. 1925 ORE DEPOSITS OF THE SADDLE MOUNTAIN AND
BANNER MINING DISTRICTS ARIZONA: USGS BULLETIN 771 .|HEINEMAN R. E. S.
ELSING M. J. ARIZONA METAL PRODUCTION ARIZONA BUREAU OF MINES ECONOMIC
SERIES NO. 19 BULLETIN NO. 14
ALT_NAME:
QUADRANGLE: EL CAPITAN MTN.
SCALE: 24000
ALTITUDE: 3320 FT
YR_FST_PRD:
YR_LST_PRD:
LAST_OPER:
CUM_PROD: 1882|1932 - 1936|1932 - 1936
CUM_P_COMM:
COMMENTS:
LATITUDE: 33-08-36N
LONGITUDE: 110-51-46W
UTM_N: 36669800
UTM_E: 5128100
UTM_ZONE: +12
INFO_SRCE: 1
REPORTER: GERE W.
REP_AFFIL: USGS
REP_DATE: 72 04
UPDATE: 79 05
REC_TYPE: X1M
REC_NO: M000383

Tracy Ranch Sec. 23, 26, T3S, R14E

Rancho Gray Sec. 25 T3S R14E

Chulito T4S, R15E Secs. 22, 23
- ASARCO + Inspir. Claims
+ Velasco claims

3 S 14 E 22 SW

D 16

REPORT DATE: AUG 8, 1991
ADMINISTRATIVE STATE: ARIZONAUNITED STATES DEPARTMENT OF THE INTERIOR
BUREAU OF LAND MANAGEMENTPAGE NO: 12555
PCN: LT892PPTGEOGRAPHIC INDEX
ALL CLAIMS

MERIDIAN: GILA-SALT R.

-LEGAL DESCRIPTION- - GEO BLM SERIAL CASE									
TOWNSHIP	RANGE	SEC	SUBDY	CITY	DIST	NO.	TYPE	CLAIM NAME/NUMBER	CLAIMANT(S)
3 S	14 E	22 SW	7	2	56053 LD	TROY MTN 12	MOORE CARLEY	56020 398:541	8/17/1964 1985 4/13/1987
SW	21	7	56054 LD	TROY MTN 13	56020 398:542	8/17/1964 1985 4/13/1987			
SW	21	7	56055 LD	TROY MTN 14	56020 398:543	8/17/1964 1985 4/13/1987			
SW	21	7	56056 LD	TROY MTN 15	56020 398:544	8/17/1964 1985 4/13/1987			
SW	21	7	56057 LD	TROY MTN 16	56020 398:545	8/17/1964 1985 4/13/1987			
SW	21	7	56062 LD	TROY MTN 22	56020 398:551	8/17/1964 1985 4/13/1987			
S2	21	7	56063 LD	TROY MTN 24	56020 398:553	8/17/1964 1985 4/13/1987			
SE	21	7	56064 LD	TROY MTN 26	56020 398:555	8/17/1964 1985 4/13/1987			
SE	21	7	56065 LD	TROY MTN 28	56020 398:557	8/17/1964 1985 4/13/1987			
SE	21	7	56066 LD	TROY MTN 29	56020 398:558	8/17/1964 1985 4/13/1987			
SE	21	7	56067 LD	TROY MTN 30	56020 398:559	8/17/1964 1985 4/13/1987			
SE	21	7	56068 LD	TROY MTN 31	56020 398:560	8/18/1964 1985 4/13/1987			
SE	21	7	56069 LD	TROY MTN 32	56020 398:561	8/18/1964 1985 4/13/1987			
W2	21	7	67233 LD	YORT NO 107	66892 673:480	5/24/1972 1985 4/03/1987			
NW	21	7	67244 LD	YORT NO 129	66892 673:491	5/17/1972 1985 4/03/1987			
NW	21	7	67245 LD	YORT NO 131	66892 673:492	5/17/1972 1985 4/03/1987			
NW	21	7	67246 LD	YORT NO 133	66892 673:493	5/17/1972 1985 4/03/1987			
NW	21	7	67247 LD	YORT NO 135	66892 673:494	5/17/1972 1985 4/03/1987			
ALL	21	7	67248 LD	YORT NO 192	66892 676:111	7/26/1972 1985 4/03/1987			
23 S2	21	7	56074 LD	HORSE 5	56020 525:77	10/02/1967 1985 4/13/1987			
SE	21	7	56075 LD	HORSE 6	56020 221:196	10/02/1967 1985 4/13/1987			
SE	21	7	56076 LD	HORSE 7	56020 221:197	10/02/1967 1985 4/13/1987			

* * DISCLOSURE * * ALL INFORMATION RECEIVED IN THIS OFFICE MAY NOT YET BE LISTED ON THIS REPORT. NAMES AND ADDRESSES ARE ENTERED AS THEY APPEAR ON THE LOCATION NOTICE OR ARE ABBREVIATED TO FIT LIMITED SPACE: THEREFORE THEY MAY NOT APPEAR IN THE EXPECTED SEQUENCE. A BLANK LATEST ASSESSMENT YEAR IN THIS REPORT DOES NOT CONSTITUTE AN ABANDONED CLAIM. * AFTER S/N INDICATES LAND STATUS CHECKED.

3 S 14 E 23 SE

E 16

REPORT DATE: AUG 8, 1991
ADMINISTRATIVE STATE: ARIZONAUNITED STATES DEPARTMENT OF THE INTERIOR
BUREAU OF LAND MANAGEMENTPAGE NO: 12556
PCN: LT892PP1GEOGRAPHIC INDEX
ALL CLAIMS

MERIDIAN: GILA-SALT R.

-LEGAL DESCRIPTION- GEO BLM SERIAL CASE		CLAIM NAME/NUMBER	CLAIMANT(S)	LEAD FILE	COUNTY BOOK:PAGE	LOCATION DATE	LATEST ASSMT-YR	CASE CLOSED
TOWNSHIP	RANGE SEC SUBDY CTY DIST	NO.	TYPE					
3 S	14 E	23 SE	7 4	56077 LD	HORSE 8	MOORE CARLEY	56020 221:198	10/02/1967 1985 4/13/1987
			21	56078 LD	HORSE 9		56020 221:199	10/02/1967 1985 4/13/1987
			SE	116639 LD	YELLOW ROSE GR NO1	ANGELIN ALBERT	1037:441 11/13/1980	0000 8/08/1985
			NW	126325 LD	YELLOW ROSE NO 2	ANGELIN ALBERT		
			21		ANGELIN WOODROW			
			NW		ANGELIN WOODROW			
			21		REITSNYDER SYLVESTER			
			24 S2	11392*LD	CEDAR TREE NO 1		11392 152:157	6/16/1963 1990
			S2	11393*LD	CEDAR TREE NO 2		11392 152:158	6/16/1963 1990
			SE	11394*LD	CEDAR TREE NO 3		11392 152:159	6/16/1963 1990
			SE	11395*LD	CEDAR TREE NO 4		11392 152:160	6/29/1963 1990
			SE	11396*LD	CEDAR TREE NO 5		11392 152:161	6/29/1963 1987
			SE	11397*LD	CEDAR TREE NO 6		11392 152:162	6/16/1963 1987
			N2	67197*LD	RAY SOUTHERN NO 40	ASARGO	66892 455:87	6/07/1956 1990
			SW	232425*PL	DSW #4	MAXWELL DANIEL	232422 636:886	10/27/1984 1987 5/19/1989
					MAXWELL LAWRENCE			
					MAXWELL NELLIE			
					MAXWELL BRIAN			
					VILLALOBOS LARRY			
					SAUTTI SCOTT			
					OERTER PAUL			
					JOHNSON WAYNE			
			SE	232426*PL	DSW #5	MAXWELL DANIEL	232422 636:888	10/27/1984 1987 5/13/1989
					MAXWELL LAWRENCE			
					MAXWELL NELLIE			
					MAXWELL BRIAN			
					VILLALOBOS LARRY			
					SAUTTI SCOTT			
					OERTER PAUL			
					JOHNSON WAYNE			
			25 SW	55960 LD	TROY #29		321:831	3/21/1972 1984 5/23/1986
			21		MOORE CARLEY			
			SW	55961 LD	TROY #30		55960 321:832	3/21/1972 1984 5/23/1986
			21		MOORE CARLEY			
			SE	55962 LD	TROY #31		55960 321:833	3/21/1972 1984 5/23/1986
			21		MOORE CARLEY			
			SE	55963 LD	TROY #32		55260 321:834	3/21/1972 1984 5/23/1986
			21		MOORE CARLEY			
			SE	55964 LD	TROY #33		55960 321:835	3/21/1972 1984 5/23/1986
			21		VIA CLARENCE			
			SE	55965 LD	TROY #34		55960 321:836	3/21/1972 1984 5/23/1986
			21		MOORE CARLEY			
			SE	55966 LD	TROY #35		55960 321:837	3/21/1972 1984 5/23/1986
			21		VIA CLARENCE			

* * DISCLOSURE * * ALL INFORMATION RECEIVED IN THIS OFFICE MAY NOT YET BE LISTED ON THIS REPORT. NAMES AND ADDRESSES ARE ENTERED AS THEY APPEAR ON THE LOCATION NOTICE OR ARE ABBREVIATED TO FIT LIMITED SPACE; THEREFORE THEY MAY NOT APPEAR IN THE EXPECTED SEQUENCE. A BLANK LATEST ASSESSMENT YEAR IN THIS REPORT DOES NOT CONSTITUTE AN ABANDONED CLAIM. * AFTER S/N INDICATES LAND STATUS CHECKED.

3 S 14 E 25 SE

F 16

REPORT DATE: AUG 8, 1991
ADMINISTRATIVE STATE: ARIZONAUNITED STATES DEPARTMENT OF THE INTERIOR
BUREAU OF LAND MANAGEMENTPAGE NO: 12557
PCN: LT892PPIGEOGRAPHIC INDEX
ALL CLAIMS

MERIDIAN: GILA-SALT R.

-LEGAL DESCRIPTION- - GEO BLM SERIAL CASE									
TOWNSHIP	RANGE	SEC	SUBDY	CITY	DIST	NO.	TYPE	CLAIM NAME/NUMBER	CLAIMANT(S)
3 S	14 E	25 SE	21	4		55966 LD	TROY #35		MOORE CARLEY
			SE			55967 LD	TROY #36		VIA CLARENCE
			21			55968 LD	TROY #37		MOORE CARLEY
			SE			55969 LD	TROY #38		VIA CLARENCE
			21			55970 LD	TROY #39		MOORE CARLEY
			SE			55971 LD	TROY #40		VIA CLARENCE
			21			56116 LD	YORT NO. 162		MOORE CARLEY
			SE			106168 PL	WATER		ROBINSON S
			21			106169 PL	AIR		
			NE			232423*PL	DSW #2		MAXWELL DANIEL
			21						MAXWELL LAURENCE
			NE						MAXWELL NELLIE
			21						MAXWELL BRIAN
			NE						VILLALOBOS LARRY
			21						SAUTTI SCOTT
			NE						JOHNSON WAYNE
			21						MAXWELL DANIEL
			NE						MAXWELL LAURENCE
			21						MAXWELL NELLIE
			NE						MAXWELL BRIAN
			21						VILLALOBOS LARRY
			NE						SAUTTI SCOTT
			21						JOHNSON WAYNE
			NE						RUSSELL CARL
			21			249192*LD	WASP #11		
			SE			249193*LD	WASP #V		
			21			249194*LD	WASP #V1		
			SE			249195*LD	WASP #V11		
			21			268063*LD	DCM #1		VANSLAMBRUCK THOMAS
			SE			268064*LD	DCM #2		LYNN FRANK
			21						VANSLAMBRUCK THOMAS
			SE						

* DISCLOSURE * * ALL INFORMATION RECEIVED IN THIS OFFICE MAY NOT YET BE LISTED ON THIS REPORT, NAMES AND ADDRESSES ARE ENTERED AS THEY APPEAR ON THE LOCATION NOTICE OR ARE ABBREVIATED TO FIT LIMITED SPACE. THEREFORE THEY MAY NOT APPEAR IN THE EXPECTED SEQUENCE. A BLANK LATEST ASSESSMENT YEAR IN THIS REPORT DOES NOT CONSTITUTE AN ABANDONED CLAIM. * AFTER S/N INDICATES LAND STATUS CHECKED.

3 S 14 E 25 SE

G 16

REPORT DATE: AUG 8, 1991
ADMINISTRATIVE STATE: ARIZONAUNITED STATES DEPARTMENT OF THE INTERIOR
BUREAU OF LAND MANAGEMENTPAGE NO: 12558
PCN: L1892PP1GEOGRAPHIC INDEX
ALL CLAIMS

MERIDIAN: GILA-SALT R.

-LEGAL DESCRIPTION- - GEO BLM SERIAL CASE									
TOWNSHIP	RANGE	SEC	SUBDY	CTY	DIST	NO.	TYPE	CLAIM NAME/NUMBER	CLAIMANT(S)
3 S	14 E	25 SE	21	4	268064*LD	DSM #2		LYNN FRANK	
		NE	7		270779*LD	CB NO 1		MURPHY ROBERT	
		ALL	21		270780*LD	CB NO 2		WITHERS CHARLES	
		E2	21		270781*LD	CB NO 3		MURPHY ROBERT	
		E2	21		270782*LD	CB NO 4		WITHERS CHARLES	
		S2	21		271863*LD	CB #5		MURPHY ROBERT	
		SE	21		271864*LD	CB #6		WITHERS CHARLES	
		SE	21		271865*LD	CB #7		MURPHY ROBERT	
		SE	21		271866*LD	CB #8		WITHERS CHARLES	
		E2	21		271867*LD	CB #9		MURPHY ROBERT	
		SW	21		271868*LD	CB #10		WITHERS CHARLES	
		SW	21		271869*LD	CB #11		MURPHY ROBERT	
		SE	21		271870*LD	CB #12		WITHERS CHARLES	
		SE	21		271871*LD	CB #13		MURPHY ROBERT	
		SE	21		274839*LD	CB #14		WITHERS CHARLES	
		SE	21		274840*LD	CB #15		MURPHY ROBERT	
		SE	21		274841*LD	CB #16		WITHERS CHARLES	
		SE	21		274842*LD	CB #17		MURPHY ROBERT	
		S2	21		274847*LD	CB #22		WITHERS CHARLES	
		SE	21		274848*LD	CB #23		MURPHY ROBERT	
		E2	21		274849*LD	CB #24		WITHERS CHARLES	
		N2	21		274885*LD	TROY 1		NICHOLS MONY	
		N2	21		274886*LD	TROY 2		NICHOLS MONY	

* * DISCLOSURE * * ALL INFORMATION RECEIVED IN THIS OFFICE MAY NOT YET BE LISTED ON THIS REPORT. NAMES AND ADDRESSES ARE ENTERED AS THEY APPEAR ON THE LOCATION NOTICE OR ARE ABBREVIATED TO FIT LIMITED SPACE. THEREFORE THEY MAY NOT APPEAR IN THE EXPECTED SEQUENCE. A BLANK LATEST ASSESSMENT YEAR IN THIS REPORT DOES NOT CONSTITUTE AN ABANDONED CLAIM. * AFTER S/N INDICATES LAND STATUS CHECKED.

3 S 14 E 25 N2

H 16

REPORT DATE: AUG 8, 1991
ADMINISTRATIVE STATE: ARIZONAUNITED STATES DEPARTMENT OF THE INTERIOR
BUREAU OF LAND MANAGEMENTPAGE NO: 12559
PCN: LTR92PP1GEOGRAPHIC INDEX
ALL CLAIMS

MERIDIAN: GILA-SALT R.

-LEGAL DESCRIPTION- GEO BLM SERIAL CASE CLAIM NAME/NUMBER CLAIMANT(S) LEAD COUNTY LOCATION LATEST CASE												
TOWNSHIP	RANGE	SEC	SUBDY	CITY	DIST	NO.	TYPE	FILE	BOOK:PAGE	DATE	ASMT-YR	CLOSED
3 S	14 E	25 N2	21	4		274886*LD	TROY 2	274885	0715:0882	9/22/1987	1989	5/13/1991
		N2	21			274887*LD	TROY 3	274885	0715:0884	9/22/1987	1989	5/13/1991
		NW	21			274888*LD	TROY 4	274885	0715:0886	9/22/1987	1989	5/13/1991
		ALL	21			274889*LD	TROY 5	274885	0715:0888	9/22/1987	1989	5/13/1991
		S2	21			274890*LD	TROY 6	274885	0715:0890	9/22/1987	1989	5/13/1991
		NW	21			274891*LD	TROY 7	274885	0715:0892	9/22/1987	1989	5/13/1991
		NW	21			274892*LD	TROY 8	274885	0715:0894	9/22/1987	1989	5/13/1991
		NW	21			274893*LD	TROY 9	274885	0715:0896	9/22/1987	1989	5/13/1991
		NW	21			274894*LD	TROY 10	274885	0715:0898	9/22/1987	1989	5/13/1991
		W2	21			274895*LD	TROY 11	274885	0715:0900	9/22/1987	1989	5/13/1991
		SW	21			274896*LD	TROY 12	274885	1474:0434	9/22/1987	1989	5/13/1991
		SE	21			279857*LD	C B 30	279856	726:302	2/01/1988	1989	5/22/1991
		SE	21			305164*LD	C B NO 14	305164	0802:0672	6/13/1990	1990	
		SE	21			305165*LD	C B NO 17	305164	0802:0674	6/13/1990	1990	
		26 NW	21	2		56070 LD	HORSE 1	56020	508:210	4/07/1967	1985	4/13/1987
		NW	21			56071 LD	HORSE 2	56020	508:211	4/07/1967	1985	4/13/1987
		NW	21			56072 LD	HORSE 3	56020	508:212	4/07/1967	1985	4/13/1987
		NW	21			56073 LD	HORSE 4	56020	508:213	4/07/1967	1985	4/13/1987
		N2	21			56074 LD	HORSE 5	56020	525:77	10/02/1967	1985	4/13/1987
		NE	21			56075 LD	HORSE 6	56020	221:196	10/02/1967	1985	4/13/1987
		NE	21			56076 LD	HORSE 7	56020	221:197	10/02/1967	1985	4/13/1987
		NE	21			56077 LD	HORSE 8	56020	221:198	10/02/1967	1985	4/13/1987
		NE	21			56078 LD	HORSE 9	56020	221:199	10/02/1967	1985	4/13/1987

* * DISCLOSURE * * ALL INFORMATION RECEIVED IN THIS OFFICE MAY NOT YET BE LISTED ON THIS REPORT. NAMES AND ADDRESSES ARE ENTERED AS THEY APPEAR ON THE LOCATION NOTICE OR ARE ABBREVIATED TO FIT LIMITED SPACE; THEREFORE THEY MAY NOT APPEAR IN THE EXPECTED SEQUENCE. A BLANK LATEST ASSESSMENT YEAR IN THIS REPORT DOES NOT CONSTITUTE AN ABANDONED CLAIM. * AFTER S/N INDICATES LAND STATUS CHECKED.

3 S 14 E 26 NE

I 16

REPORT DATE: AUG 8, 1991
ADMINISTRATIVE STATE: ARIZONAUNITED STATES DEPARTMENT OF THE INTERIOR
BUREAU OF LAND MANAGEMENTPAGE NO: 12560
PCN: L1892PPIGEOGRAPHIC INDEX
ALL CLAIMS

MERIDIAN: GILA-SALT R.

-LEGAL DESCRIPTION- GEO BLM		SERIAL	CASE	CLAIM NAME/NUMBER	CLAIMANT(S)	LEAD	COUNTY	LOCATION	LATEST	CASE
TOWNSHIP	RANGE SEC SUBDY CITY DIST	NO.	TYPE			FILE	BOOK:PAGE	DATE	ASSMT-TR	CLOSED
3 S	14 E	26 NE	7	2	56079 LD	HORSE 10	MOORE CARLEY	56020 221:200	10/02/1967	1985 4/13/1987
			21		56082 LD	RUSTY NO. 3		56020 676:115	7/20/1972	1985 4/13/1987
			21		56083 LD	RUSTY NO. 4		56020 676:116	7/20/1972	1985 4/13/1987
			21		56084 LD	RUSTY NO. 5		56020 676:117	7/20/1972	1985 4/13/1987
			21		56085 LD	RUSTY NO. 6		56020 676:118	7/20/1972	1985 4/13/1987
			21		56086 LD	RUSTY NO. 7		56020 676:119	7/20/1972	1985 4/13/1987
			21		56087 LD	RUSTY NO. 8		56020 676:120	7/20/1972	1985 4/13/1987
			21		56088 LD	RUSTY NO. 9		56020 676:121	7/20/1972	1985 4/13/1987
			21		56089 LD	RUSTY NO. 10		56020 676:122	7/20/1972	1985 4/13/1987
			21		56090 LD	RUSTY NO. 11		56020 676:123	7/20/1972	1985 4/13/1987
			21		56091 LD	RUSTY NO. 12		56020 676:124	7/20/1972	1985 4/13/1987
			21		56092 LD	RUSTY NO. 13		56020 676:125	7/20/1972	1985 4/13/1987
			21		56093 LD	RUSTY NO. 14		56020 676:126	7/20/1972	1985 4/13/1987
			21		56109 LD	RUSTY NO. 148		56020 676:98	6/27/1972	1985 4/13/1987
			21		56110 LD	RUSTY NO. 150		56020 676:99	6/27/1972	1985 4/13/1987
			21		56111 LD	RUSTY NO. 152		56020 676:100	6/27/1972	1985 4/13/1987
			21		56112 LD	RUSTY NO. 154		56020 676:101	6/27/1972	1985 4/13/1987
			21		56113 LD	RUSTY NO. 156		56020 676:102	6/27/1972	1985 4/13/1987
			21		56114 LD	RUSTY NO. 158		56020 676:103	6/27/1972	1985 4/13/1987
			21		56115 LD	RUSTY NO. 160		56020 676:104	6/27/1972	1985 4/13/1987
			21		56117 LD	RUSTY NO. 187		56020 676:106	6/27/1972	1985 4/13/1987
			21		274897-10	TROY 13	RAMSEY CAROL ELDER LAWRENCE NICHOLS MONY	274885 1474:0436	9/22/1987	1989 5/13/1991

* * DISCLOSURE * * ALL INFORMATION RECEIVED IN THIS OFFICE MAY NOT YET BE LISTED ON THIS REPORT, NAMES AND ADDRESSES ARE ENTERED AS THEY APPEAR ON THE LOCATION NOTICE OR ARE ABBREVIATED TO FIT LIMITED SPACE; THEREFORE THEY MAY NOT APPEAR IN THE EXPECTED SEQUENCE. A BLANK LATEST ASSESSMENT YEAR IN THIS REPORT DOES NOT CONSTITUTE AN ABANDONED CLAIM. * AFTER S/N INDICATES LAND STATUS CHECKED.

3 S 14 E 26 NE

J 16

REPORT DATE: AUG 8, 1991
ADMINISTRATIVE STATE: ARIZONAUNITED STATES DEPARTMENT OF THE INTERIOR
BUREAU OF LAND MANAGEMENTPAGE NO: 12561
PCN: 11892PPIGEOGRAPHIC INDEX
ALL CLAIMS

MERIDIAN: GILA-SALT R.

-LEGAL DESCRIPTION- - GEO BLM SERIAL CASE									
TOWNSHIP	RANGE	SEC	SUBDY	CITY	DIST	NO.	TYPE	CLAIM NAME/NUMBER	CLAIMANT(S)
FILE	COUNTY	LOCATION	LATEST	CASE					
BOOK	PAGE	DATE	ASMT-YR	CLOSED					
3 S	14 E	26 NE	21	2	274897-LD	TROY 13	SUMPTER LAVERN	274885 0715:0904	9/22/1987 1989 5/13/1991
NE			7		274898-LD	TROY 14	NICHOLS MONY	274885 1474:0438	9/22/1987 1989 5/13/1991
			21		274899-LD	TROY 15	SUMPTER LAVERN	0715:0906	
NE			21		274900-LD	TROY 16	NICHOLS MONY	1474:0440	9/22/1987 1989 5/13/1991
			21		274901-LD	TROY 17	SUMPTER LAVERN	0715:0910	
E2			21		274902-LD	TROY 18	NICHOLS MONY	1474:0446	9/22/1987 1989 5/13/1991
SE			21		274903-LD	TROY 19	SUMPTER LAVERN	0715:0912	
SE			21		274904-LD	TROY 20	NICHOLS MONY	1474:0450	9/22/1987 1989 5/13/1991
SE			21		274905-LD	TROY 21	SUMPTER LAVERN	1474:0450	9/22/1987 1989 5/13/1991
N2			21		274906-LD	TROY 22	NICHOLS MONY	0715:0914	
N2			21		274907-LD	TROY 23	SUMPTER LAVERN	1474:0454	9/22/1987 1989 5/13/1991
N2			21		274908-LD	TROY 24	NICHOLS MONY	1474:0456	9/22/1987 1989 5/13/1991
ALL			21		274909-LD	TROY 25	SUMPTER LAVERN	1474:0460	9/22/1987 1989 5/13/1991
S2			21		274910-LD	TROY 26	NICHOLS MONY	1474:0462	9/22/1987 1989 5/13/1991
S2			21		274911-LD	TROY 27	SUMPTER LAVERN	1474:0464	9/22/1987 1989 5/13/1991
S2			21		274912-LD	TROY 28	NICHOLS MONY	1474:0466	9/22/1987 1989 5/13/1991
NW			21		274913-LD	TROY 29	SUMPTER LAVERN	1474:0468	9/22/1987 1989 5/13/1991
NW			21		274914-LD	TROY 30	NICHOLS MONY	1474:0470	9/22/1987 1989 5/13/1991
NW			21		274915-LD	TROY 31	SUMPTER LAVERN	1474:0472	9/22/1987 1989 5/13/1991
NW			21		274916-LD	TROY 32	NICHOLS MONY	1474:0474	9/22/1987 1989 5/13/1991
W2			21		274917-LD	TROY 33	SUMPTER LAVERN	1474:0476	9/22/1987 1989 5/13/1991
SW			21		274918-LD	TROY 34	NICHOLS MONY	1474:0478	9/22/1987 1989 5/13/1991
SW			21		274919-LD	TROY 35	SUMPTER LAVERN	1474:0480	9/22/1987 1989 5/13/1991

* * DISCLOSURE * * ALL INFORMATION RECEIVED IN THIS OFFICE MAY NOT YET BE LISTED ON THIS REPORT. NAMES AND ADDRESSES ARE ENTERED AS THEY APPEAR ON THE LOCATION NOTICE OR ARE ABBREVIATED TO FIT LIMITED SPACE. THEREFORE THEY MAY NOT APPEAR IN THE EXPECTED SEQUENCE. A BLANK LATEST ASSESSMENT YEAR IN THIS REPORT DOES NOT CONSTITUTE AN ABANDONED CLAIM. * AFTER S/N INDICATES LAND STATUS CHECKED.

3 S 14 E 26 SW

K 16

REPORT DATE: AUG 8, 1991
ADMINISTRATIVE STATE: ARIZONAUNITED STATES DEPARTMENT OF THE INTERIOR
BUREAU OF LAND MANAGEMENTPAGE NO: 12562
PCN: L1892PPIGEOGRAPHIC INDEX
ALL CLAIMS

MERIDIAN: GILA-SALT R.

-LEGAL DESCRIPTION- GEO BLM SERIAL CASE									
TOWNSHIP	RANGE	SEC	SUBDY	CITY	DIST	NO.	TYPE	CLAIM NAME/NUMBER	CLAIMANT(S)
3 S	14 E	26 SW	21	2	274919-LD	TROY 35		SUMPTER LAVERN	NICHOLS MONY
			21	2	274920-LD	TROY 36		SUMPTER LAVERN	NICHOLS MONY
			21	2	274921-LD	TROY 37		SUMPTER LAVERN	NICHOLS MONY
			21	2	274922-LD	TROY 38		SUMPTER LAVERN	NICHOLS MONY
			21	2	274923-LD	TROY 39		SUMPTER LAVERN	NICHOLS MONY
			21	2	274924-LD	TROY 40		SUMPTER LAVERN	NICHOLS MONY
			21	2	275214-LD	TROY 41		SUMPTER LAVERN	NICHOLS MONY
			21	2	275215-LD	TROY 42		SUMPTER LAVERN	NICHOLS MONY
			21	2	275216-LD	TROY 43		SUMPTER LAVERN	NICHOLS MONY
			21	2	275217-LD	TROY 44		SUMPTER LAVERN	NICHOLS MONY
			21	2	56035 LD	ALICE NO 4		RAMSEY CAROL	ELDER LAWRENCE
			21	2	56036 LD	ALICE NO 5		RAMSEY CAROL	ELDER LAWRENCE
			21	2	56037 LD	ALICE NO 6		RAMSEY CAROL	ELDER LAWRENCE
			21	2	56038 LD	ALICE NO 8		RAMSEY CAROL	ELDER LAWRENCE
			21	2	56039 LD	ALICE NO 9		RAMSEY CAROL	ELDER LAWRENCE
			21	2	56040 LD	ALICE NO 10		RAMSEY CAROL	ELDER LAWRENCE
			21	2	56041 LD	ALICE NO 11		RAMSEY CAROL	ELDER LAWRENCE
			21	2	56042 LD	MARY ALICE #1		MOORE CARLEY	56020 278:222 11/14/1960 1985 4/13/1987
			21	2	56043 LD	MARY ALICE #2		MOORE CARLEY	56020 278:223 11/14/1960 1985 4/13/1987
			21	2	56046 LD	MARY ALICE NO 7		MOORE CARLEY	56020 278:228 11/14/1960 1985 4/13/1987
			21	2	56047 LD	TROY MIN. 1		MOORE CARLEY	56020 403:275 12/14/1964 1985 4/13/1987

* * DISCLOSURE * * ALL INFORMATION RECEIVED IN THIS OFFICE MAY NOT YET BE LISTED ON THIS REPORT. NAMES AND ADDRESSES ARE ENTERED AS THEY APPEAR ON THE LOCATION NOTICE OR ARE ABBREVIATED TO FIT LIMITED SPACE. THEREFORE THEY MAY NOT APPEAR IN THE EXPECTED SEQUENCE. A BLANK LATEST ASSESSMENT YEAR IN THIS REPORT DOES NOT CONSTITUTE AN ABANDONED CLAIM. * AFTER S/N INDICATES LAND STATUS CHECKED.

3 S 14 E 27 NW

L 16

REPORT DATE: AUG 8, 1991
ADMINISTRATIVE STATE: ARIZONAUNITED STATES DEPARTMENT OF THE INTERIOR
BUREAU OF LAND MANAGEMENTPAGE NO: 12563
PCN: L1892PP1GEOGRAPHIC INDEX
ALL CLAIMS

MERIDIAN: GILA-SALT R.

-LEGAL DESCRIPTION-- GEO BLM SERIAL CASE										CLAIM NAME/NUMBER		CLAIMANT(S)		LEAD	COUNTY	LOCATION	LATEST	CASE
TOWNSHIP	RANGE	SEC	SUBDY	CITY	DIST	NO.	TYPE						FILE	BOOK	PAGE	DATE	ASSMT-YR	CLOSED
3 S	14 E	27 NW				21	2	56048 LD	TROY MIN. 2		MOORE CARLEY	56020	403:276	12/14/1964	1985	4/13/1987		
		NW				21		56049 LD	TROY MIN. 3			56020	403:277	12/14/1964	1985	4/13/1987		
		NW				21		56052 LD	TROY MIN 11			56020	398:540	8/17/1964	1985	4/13/1987		
		NW				21		56054 LD	TROY MIN 13			56020	398:542	8/17/1964	1985	4/13/1987		
		NW				21		56056 LD	TROY MIN. 15			56020	398:544	8/17/1964	1985	4/13/1987		
		NE				21		56070 LD	HORSE 1			56020	508:210	4/07/1967	1985	4/13/1987		
		E2				21		56080 LD	RUSTY NO. 1			56020	676:113	7/20/1972	1985	4/13/1987		
		E2				21		56081 LD	RUSTY NO. 2			56020	676:114	7/20/1972	1985	4/13/1987		
		NE				21		56093 LD	RUSTY NO. 14			56020	676:126	7/20/1972	1985	4/13/1987		
		SE				21		56094 LD	YORT NO. 12			56020	673:411	4/23/1972	1985	4/13/1987		
											RANSEY CAROL							
		SE				21		56095 LD	YORT NO. 13			56020	673:412	4/23/1972	1985	4/13/1987		
											ELDER LAWRENCE							
											MOORE CARLEY							
											RANSEY CAROL							
		SE				21		56096 LD	YORT NO. 14			56020	673:413	4/23/1972	1985	4/13/1987		
											ELDER LAWRENCE							
											MOORE CARLEY							
											RANSEY CAROL							
		SE				21		56097 LD	YORT NO. 15			56020	673:414	4/23/1972	1985	4/13/1987		
											ELDER LAWRENCE							
											MOORE CARLEY							
											RANSEY CAROL							
		SE				21		56098 LD	YORT NO. 18			56020	673:417	4/23/1972	1985	4/13/1987		
											ELDER LAWRENCE							
											MOORE CARLEY							
											RANSEY CAROL							
		SE				21		56099 LD	YORT NO. 19			56020	673:418	4/23/1972	1985	4/13/1987		
											ELDER LAWRENCE							
											MOORE CARLEY							
											RANSEY CAROL							
		SE				21		56100 LD	YORT NO. 20			56020	673:419	4/23/1972	1985	4/13/1987		
											ELDER LAWRENCE							
											MOORE CARLEY							
											RANSEY CAROL							
		E2				21		56101 LD	YORT NO. 138			56020	673:496	6/16/1972	1985	4/13/1987		
											ELDER LAWRENCE							
											MOORE CARLEY							
											RANSEY CAROL							
		E2				21		56102 LD	YORT NO. 139			56020	673:497	6/16/1972	1985	4/13/1987		
											ELDER LAWRENCE							
											MOORE CARLEY							
											RANSEY CAROL							
		ALL				21		56103 LD	YORT NO. 140			56020	673:498	6/16/1972	1985	4/13/1987		
											ELDER LAWRENCE							
											MOORE CARLEY							
											RANSEY CAROL							
		W2				21		56104 LD	YORT NO. 141			56020	673:499	6/16/1972	1985	4/13/1987		
											ELDER LAWRENCE							
											MOORE CARLEY							
											RANSEY CAROL							
		NW				21		56106 LD	YORT NO. 144			56020	673:502	6/17/1972	1985	4/13/1987		
											ELDER LAWRENCE							
											MOORE CARLEY							
											RANSEY CAROL							
											ELDER LAWRENCE							

* * DISCLOSURE * * ALL INFORMATION RECEIVED IN THIS OFFICE MAY NOT YET BE LISTED ON THIS REPORT. NAMES AND ADDRESSES ARE ENTERED AS THEY APPEAR ON THE LOCATION NOTICE OR ARE ABBREVIATED TO FIT LIMITED SPACE; THEREFORE THEY MAY NOT APPEAR IN THE EXPECTED SEQUENCE. A BLANK LATEST ASSESSMENT YEAR IN THIS REPORT DOES NOT CONSTITUTE AN ABANDONED CLAIM. * AFTER S/N INDICATES LAND STATUS CHECKED.

3 S 14 E 27 NW

M 16

REPORT DATE: AUG 8, 1991
ADMINISTRATIVE STATE: ARIZONAUNITED STATES DEPARTMENT OF THE INTERIOR
BUREAU OF LAND MANAGEMENTPAGE NO: 12564
PCN: L1892BPIGEOGRAPHIC INDEX
ALL CLAIMS

MERIDIAN: GILA-SALT R.

-LEGAL DESCRIPTION- - GEO BLM SERIAL CASE CLAIM NAME/NUMBER CLAIMANT(S)													
TOWNSHIP	RANGE	SEC	SUBDY	CITY	DIST	NO.	TYPE	FILE	COUNTY	BOOK/PAGE	LOCATION DATE	LATEST ASMT-YR	CASE CLOSED
3 S	14 E	27 NW	21	2	56107	LD	YORT NO. 145	56020	673:503	6/22/1972	1985	4/13/1987	
							MOORE CARLEY RAMSEY CAROL						
							ELDER LAWRENCE MOORE CARLEY	56020	673:504	6/22/1972	1985	4/13/1987	
							RAMSEY CAROL						
							ELDER LAWRENCE MOORE CARLEY	56020	676:107	7/21/1972	1985	4/13/1987	
							RAMSEY CAROL						
							ELDER LAWRENCE NICHOLS MONY	274885	1474:0484	9/22/1987	1989	5/13/1991	
							SUMPTER LAVERN						
							NICHOLS MONY	274885	1474:0486	9/22/1987	1989	5/13/1991	
							SUMPTER LAVERN						
							NICHOLS MONY	274885	1474:0488	9/22/1987	1989	5/13/1991	
							SUMPTER LAVERN						
							NICHOLS MONY	274885	1474:0490	9/22/1987	1989	5/13/1991	
							SUMPTER LAVERN						
							NICHOLS MONY	275214	1474:492	9/22/1987	1989	5/13/1991	
							SUMPTER LAVERN						
							NICHOLS MONY	275214	1474:494	9/22/1987	1989	5/13/1991	
							SUMPTER LAVERN						
							NICHOLS MONY	275214	1474:496	9/22/1987	1989	5/13/1991	
							SUMPTER LAVERN						
							NICHOLS MONY	275214	1474:498	9/22/1987	1989	5/13/1991	
							SUMPTER LAVERN						
							NICHOLS MONY	275214	1474:500	9/22/1987	1989	10/20/1987	
							SUMPTER LAVERN						
							NICHOLS MONY	275214	1474:502	9/22/1987	1989	10/20/1987	
							SUMPTER LAVERN						
							NICHOLS MONY	275214	1474:504	9/22/1987	1989	5/13/1991	
							SUMPTER LAVERN						
							NICHOLS MONY	275214	1474:506	9/22/1987	1989	5/13/1991	
							SUMPTER LAVERN						
							NICHOLS MONY	275214	1474:508	9/22/1987	1989	5/13/1991	
							SUMPTER LAVERN						
							NICHOLS MONY	275214	1474:510	9/22/1987	1989	5/13/1991	
							SUMPTER LAVERN						
							NICHOLS MONY	275214	1474:512	9/22/1987	1989	5/13/1991	
							SUMPTER LAVERN						
							NICHOLS MONY	275214	1474:514	9/22/1987	1989	5/13/1991	
							SUMPTER LAVERN						
							NICHOLS MONY	275214	1474:516	9/22/1987	1989	10/20/1987	
							SUMPTER LAVERN						
							NICHOLS MONY	275214	1474:518	9/22/1987	1989	10/20/1987	

* * DISCLOSURE * * ALL INFORMATION RECEIVED IN THIS OFFICE MAY NOT YET BE LISTED ON THIS REPORT. NAMES AND ADDRESSES ARE ENTERED AS THEY APPEAR ON THE LOCATION NOTICE OR ARE ABBREVIATED TO FIT LIMITED SPACE; THEREFORE THEY MAY NOT APPEAR IN THE EXPECTED SEQUENCE. A BLANK LATEST ASSESSMENT YEAR IN THIS REPORT DOES NOT CONSTITUTE AN ABANDONED CLAIM. * AFTER S/N INDICATES LAND STATUS CHECKED.

3 S 14 E 27 NW

N 16

REPORT DATE: AUG 8, 1991
ADMINISTRATIVE STATE: ARIZONAUNITED STATES DEPARTMENT OF THE INTERIOR
BUREAU OF LAND MANAGEMENTPAGE NO: 12565
PEN: 11892PPTGEOGRAPHIC INDEX
ALL CLAIMS

MERIDIAN: GILA-SALT R.

-LEGAL DESCRIPTION- - GEO BLM SERIAL CASE CLAIM NAME/NUMBER CLAIMANT(S) LEAD COUNTY LOCATION LATEST CASE												
TOWNSHIP	RANGE	SEC	SUBDY	CTY	DIST	NO.	TYPE	FILE	BOOK:PAGE	DATE	ASSMT-YR	CLOSED
3 S	14 E	27 NW	2	275227-LD	TROY 54	SUMPTER LAVERNUR	275214	1474:518	9/22/1987	1989	10/20/1987	
		NW	21	275228-LD	TROY 55	NICHOLS MONY	275214	1474:520	9/22/1987	1989	5/13/1991	
		NW	21	275229-LD	TROY 56	SUMPTER LAVERNUR	275214	1474:522	9/22/1987	1989	5/13/1991	
		W2	21	275230-LD	TROY 57	NICHOLS MONY	275214	1474:524	9/22/1987	1989	5/13/1991	
		SW	21	275231-LD	TROY 58	SUMPTER LAVERNUR	275214	1474:526	9/22/1987	1989	5/13/1991	
		SW	21	275232-LD	TROY 59	SUMPTER LAVERNUR	275214	1474:528	9/22/1987	1989	5/13/1991	
		SW	21	275233-LD	TROY 60	NICHOLS MONY	275214	1474:530	9/22/1987	1989	5/13/1991	
		NW	21	275234-LD	TROY 61	SUMPTER LAVERNUR	275214	1474:532	9/22/1987	1989	5/13/1991	
		NW	21	275235-LD	TROY 62	SUMPTER LAVERNUR	275214	1474:534	9/22/1987	1989	5/13/1991	
		NW	21	275236-LD	TROY 63	NICHOLS MONY	275214	1474:536	9/22/1987	1989	5/13/1991	
		NW	21	275237-LD	TROY 64	SUMPTER LAVERNUR	275214	1474:538	9/22/1987	1989	5/13/1991	
		W2	21	275238-LD	TROY 65	SUMPTER LAVERNUR	275214	1474:540	9/22/1987	1989	5/13/1991	
		SW	21	275239-LD	TROY 66	NICHOLS MONY	275214	1474:542	9/22/1987	1989	5/13/1991	
		SW	21	275240-LD	TROY 67	SUMPTER LAVERNUR	275214	1474:544	9/22/1987	1989	5/13/1991	
		SW	21	275241-LD	TROY 68	SUMPTER LAVERNUR	275214	1474:546	9/22/1987	1989	5/13/1991	
		NW	21	275242-LD	TROY 69	NICHOLS MONY	275214	1474:548	9/22/1987	1989	5/13/1991	
		W2	21	275243-LD	TROY 70	SUMPTER LAVERNUR	275214	1474:550	9/22/1987	1989	5/13/1991	
		SW	21	275244-LD	TROY 71	SUMPTER LAVERNUR	275214	1474:552	9/22/1987	1989	5/13/1991	
28 S2		SE	21	21126-LD	ALICE #3	GAYLOR JAMES E	21124	52:322	4/15/1942	1990		
		SE	21	21127-LD	ALICE #4		21124	52:327	4/15/1942	1990		
		N2, SE	21	21128-LD	ALICE #5		21124	52:327	4/15/1942	1990		
		SE	21	21129-LD	ALICE #6		21124	52:327	4/15/1942	1990		
		SE	21	21131-LD	WHITE TAIL DEER		21124	52:330	4/15/1942	1990		
		SW	21	21132-LD	PRAIRIE SPRING		21124	52:329	4/15/1942	1990		
		S2	21	21133-LD	JOHNNY BOY		21124	52:329	4/15/1942	1990		
		S2	21	21134-LD	ALTO		21124	52:328	4/15/1942	1990		
		SW	21	21135-LD	BURRO		21124	52:329	4/15/1942	1990		

DISCLOSURE: ALL INFORMATION RECEIVED IN THIS OFFICE MAY NOT YET BE LISTED ON THIS REPORT. NAMES AND ADDRESSES ARE ENTERED AS THEY APPEAR ON THE LOCATION NOTICE OR ARE ABBREVIATED TO FIT LIMITED SPACE; THEREFORE THEY MAY NOT APPEAR IN THE EXPECTED SEQUENCE. A BLANK LATEST ASSESSMENT YEAR IN THIS REPORT DOES NOT CONSTITUTE AN ABANDONED CLAIM. * AFTER SYN INDICATES LAND STATUS CHECKED.

DEPARTMENT OF THE INTERIOR
UNITED STATES GEOLOGICAL SURVEY

GEOLOGIC MAP OF THE HAYDEN QUADRANGLE,
PINAL AND GILA COUNTIES, ARIZONA

By Norman G. Banks
and Medora H. Krieger

Revised 11/25/91

Dripping Spgs, Banner
Steamboat Mtn. Dist.

Chilito, 79, New Year

London-Arizona Mines

QTz veins, Jasperoid near
Steamboat Mtn.

WESTMONT MINING INC.
4949 S. SYRACUSE ST. #4200
DENVER, CO 80237

outcrops of white gypsum filling joint cracks and extending beyond in masses of nearly pure gypsum. (See Pl. XII, A.) The mode of formation of these deposits and their relation, if any, to the bodies of disseminated pyrite near by are not known. There is plenty of sulphur, both as sulphide and as sulphate, deposited by hypogene solutions in neighboring ore bodies and calcium is widespread in the rocks of the region. The gypsum of these two deposits may have been precipitated from water of the vadose circulation that had derived its sulphate content from oxidizing sulphides, or from ascending solutions related to those which produced the ore bodies, or from an intermingling of such solutions and descending meteoric water.

CONTACT-METAMORPHIC DEPOSITS

The ore bodies described under the term contact-metamorphic deposits are replacement masses, largely in limestone, which owe their origin to emanations from near-by intrusive rocks. Most of them contain garnet and other characteristic contact-metamorphic minerals. In others such minerals were not found, but the deposits are otherwise so similar to the more typical contact-metamorphic deposits that it seems best to group them together. The deposits thus grouped include those of the Christmas mine, the London-Arizona properties, the Schneider group, and the Gold-Copper, Seventy-nine, Apex, and Columbia mines. The deposits of the Cowboy mine and the C. & B. and Premier prospects may also be related to this group. The deposits on the Scottish American Copper Co.'s property south of the London-Arizona claims are presumably similar, but this company was not operating at the time of visit and so far as could be learned had not done much development work. A large part are in the Tornado limestone, but there are also extensive replacement deposits in the Martin limestone, some in the Troy quartzite, and related deposits in the Mescal limestone and in diabase.

The deposits of this group show greater diversity than those previously described, partly because of "original" differences in the country rock and partly because of variations in the proportions of the minerals present. Some of the deposits are largely confined to certain limestone beds, others are of irregular outline, and still others are of veinlike form on shear zones. All these variations are due principally to differences in the effect of mineralization on different kinds of rock under what appear to be in other respects fundamentally similar conditions. Some of the deposits are mined for copper; others are mined for lead and silver, or for lead and gold. Parts of some deposits have been mined for the zinc contained, and oxidized portions of a few have been prospected for vanadium. All the deposits were formed by replacement, rather than by the

filling of open spaces, all lie near dikes or similar masses of quartz-mica diorite, and nearly all are known to contain typical contact-metamorphic silicate minerals. The evidence seems clear that most of the deposits on the properties mentioned above were formed by emanations that had their source in the quartz-mica diorite magma and that interacted with susceptible parts of the rocks in contact with the intrusions, forming replacement masses, parts of which contain sufficient quantities of metallic minerals to constitute ore. The diorite near the deposits is itself altered, and apparently a part of the mineralizing emanations welled up through it from the magma until they encountered conditions favorable for spreading out into the inclosing rocks. The alteration in the diorite consists in sericitization and the introduction of pyrite and quartz.

The shape and location of the ore bodies in limestone appear to be controlled in general by the shape, size, and position of the dioritic mass that was the source of mineralization, the character and position of fractures or shear zones in the limestone, and the size and position of beds susceptible of replacement. At some places, as at the Christmas mine, the ore bodies appear to follow the intrusive contacts rather closely. Elsewhere, especially where the dioritic masses exposed are small, the ores are farther from the contact. Most of the deposits afford evidence that the entrance of the mineralizing material was facilitated by fracturing of the rock, and the deposits are more or less localized along such fracture zones. In some places the existence of the fractures can only be inferred from the elongation of the ore bodies in certain directions. Elsewhere evidence of shearing and brecciation can be clearly seen. The places where premineral fracturing can now be discovered are in general those where replacement of the rock has not extended far beyond the fracture zones, and in a number of such places contact-metamorphic minerals were not observed. The fractures and shear zones, both inferred and observed, have steep dips and strike nearly east, many of them somewhat north of east, although there is considerable local variation. The average strike of the fractures thus corresponds to the average strike of the quartz-mica diorite dikes as shown on Plates I and XVI, and the fracturing may be related to the intrusion of the diorite and have taken place at the same time or just prior to the intrusion. At each of the deposits in limestone certain of the beds are more thoroughly replaced than others, and only fractions of the replaced portions are of value as ore. This probably results in part from original differences in composition and in permeability. At the Christmas mine, in the Tornado limestone, the comparatively pure and coarser-grained beds are more thoroughly replaced by ore minerals than the finer-grained rocks with a lower carbonate content, although the

latter are also greatly altered. Such obvious differences in lithologic character do not, however, account for some of the differences noted, as is strikingly illustrated by the bed locally known as the O'Carroll ore bed, near the base of the Martin limestone. Mineralized rock at this horizon occurs at a number of places on a line of outcrop about a mile east of Chilito, but there is little evidence of mineralization at other horizons in the formation where the original lithologic character would seem to have been similar. It is possible that this difference may be partly accounted for by the relations between the limestone and the intrusive rock. The diorite masses that presumably caused most of the contact metamorphism in this vicinity now crop out west of the O'Carroll ore bed and at lower altitudes, and the upper parts, now eroded, may never have reached much above that bed. There are, however, a number of smaller masses of similar diorite that did reach high enough to cut the Martin limestone above this horizon. The lateral limits of the replacement bodies are conceived to have been determined in each place largely by the distance the emanations traveled along a particular bed from the point of access to the point where chemical interaction ceased because of exhaustion of the emanation. The point of exhaustion was in turn dependent on the quantity, character, and physical condition of the material given off by the intrusive magma, and on the readiness with which it was able to react with and permeate the limestone. The points of access were apparently along the contact of the more replaceable bed with an intrusive body of quartz-mica diorite and along fracture zones that cut the bed. The valuable sulphides were in the main localized along the fracture zones and not at the intrusive contact. In the simplest case ore bodies produced as suggested above would have the form of elliptical disks with rounded edges, bounded above and below by beds of shale or other difficultly replaceable material. Between different ore bodies in a single bed and between the ore bodies and the intrusive mass the limestone would be more or less thoroughly replaced by minerals other than the valuable sulphides. As a matter of fact such a shape is never more than approximated, and some of the ore bodies are most irregular in form. The irregularities have resulted from the numerous possible variations in the character and condition of the bed affected, the emanations, the fractures, and the intrusive, as well as from interference between the emanations circulating from neighboring points of access to the same bed. At a number of places there are several beds which proved susceptible of replacement by ore minerals, and as a consequence more or less separate ore bodies have been formed one above another, separated by altered but valueless material. Some complexities resulting from postmineral faulting have been super-

imposed on the original irregularities, but so far as known these are of minor importance.

The minerals resulting from the contact metamorphism in limestone include quartz, two varieties of andradite garnet, vesuvianite, wollastonite, epidote, pargasite or a similar amphibole, chlorite, fluorite, chalcodony, magnetite, specularite, chalcopyrite, pyrite, sphalerite, galena, and perhaps bornite. Calcite has been recrystallized and redistributed. All these minerals, except fluorite, have been found in the Christmas mine. Most of the other deposits are known to contain some of the silicates mentioned, and more of these as well as other minerals as yet unknown in the region will probably be discovered in the course of future development and study. Many of the deposits in limestone visited contain most of the metallic sulphides listed above, but the proportions vary widely. Details of the occurrence of the several minerals are given in the mine descriptions (pp. 41-69).

At the Christmas mine and probably also elsewhere the replacement bodies grade, in most places abruptly, into limestone that has been recrystallized into marble but is otherwise little altered. The alteration in the beds between those containing the ore bodies is largely silicification, although most of the minerals mentioned above can be found in such beds in minor amounts, and epidote and chlorite are rather common in them. Many of these beds still show stratification and are plainly of sedimentary origin; these were doubtless shale and impure limestone. In places, according to Locke and Bjorge, there are sills of fine-grained quartz-mica diorite that are difficult to distinguish from metamorphosed shale. The original character of some of the beds between ore bodies can not be determined. In the Tornado limestone exposed along Gila River immediately south of the south boundary of the Christmas area there are fine-grained sills of dark-green trap like that shown in Plate XI, B. Megascopically this rock resembles the diabase that is widely distributed in the Ray quadrangle. Under the microscope it is seen to be altered to an aggregate of quartz, calcite, and indeterminate dark dust. Some of the barren strata in the Christmas mine may be composed of more thoroughly silicified rock of this character.

Little information is available regarding the ore deposits in the Troy quartzite. Ore of good grade has been mined from such rock on Schneider Hill, but the bodies found appear to have been all rather small. It is probable that these ore bodies are replacement deposits similar and closely related genetically to those in limestone just described. The major difference probably is that the quartzite was less favorable for replacement than the limestone, and consequently the ore bodies produced in it were smaller and more irregular.

The deposits in diabase at the Columbia mine differ somewhat from the other deposits described in this section but are probably related genetically to them. They are replacement deposits along shear zones near intrusive bodies of quartz-mica diorite. Contact-metamorphic minerals have not been found in the diabase, but calcareous shale in another part of the mine contains vesuvianite and garnet, and the mineralization is similar to that of the typical contact-metamorphic deposits.

Oxidation has attacked all the contact-metamorphic deposits of the area, and a large part of the ore mined was oxidized material. In most of the deposits, however, residual hypogene sulphides can be found at or near the surface. Oxidation probably nowhere extends more than 100 feet below the surface and in most places less than that. The oxidized minerals include malachite, chrysocolla, azurite, native copper, gold, hydrous manganese oxide, limonite, and kindred iron oxides, copper pitch ore, cerussite, anglesite, plumbogjarosite, wulfenite, gypsum, and alunite. Concentration of the oxidized minerals of the valuable metals has taken place to some extent, especially in the lead deposits. The rich gold ore of the Apex and Cowboy mines may have been produced by concentration during oxidation. In both the gold was found in the oxidized part of lead ore bodies, but specimens were not available for examination. Vanadinite has been found associated with wulfenite in oxidized ore at three places in the Banner district. This mineral was not recognized in any of the deposits visited in the present investigation, but wulfenite is known to have been obtained from two of them. It may well be that the vanadinite was formed by the oxidation of lead deposits similar to those of the Seventy-nine mine, for example.

Enrichment by supergene sulphides has taken place in all the copper deposits seen but appears to be sporadic. The principal supergene sulphide is chalcocite. A little covellite was noted. Bornite is present in small veinlets in the ore bodies and might be interpreted as formed either in a late stage of hypogene deposition or early in the production of supergene sulphides, both chalcocite and covellite being later than the bornite. If the bornite is supergene, then enrichment of this character has greatly improved the tenor of the ore at Christmas. Veinlets of chalcocite, almost certainly of supergene origin, penetrate the ore down to at least the 300-foot level, but the amount of enrichment from this source in the ore seen is probably not very large. To judge by the available specimens massive chalcocite was present in the ore of the Schneider Hill deposits and in the tunnels of the London-Arizona mine. These specimens were, however, probably picked from the richer ore, and a considerable part of the ore shipped is believed

to have been oxidized material without large amounts of sulphides of any kind in it.

GENERALIZATIONS AND INFERENCES

A number of generalizations and inferences can be drawn from a study of the ore deposits of the Saddle Mountain and Banner districts, and most of them have a direct bearing on economic problems. The following statements are based on all available evidence, but they must be regarded as expressions of opinion rather than of proved facts.

The ore deposits of the two districts have so many points of similarity that they may be confidently regarded as genetically related products of the same period of mineralization. Some are certainly and the others probably not older than the end of the Cretaceous, and all are of types that are generally regarded as genetically related to igneous rocks. The only igneous rocks known in either district whose character and age would permit their consideration in this connection are the quartz-mica diorite and quartz-hornblende diorite, both believed to be of early Tertiary age. Many of the ore deposits are intimately associated with quartz-mica diorite, and nearly all the known deposits have outcrops of this rock near by, whereas the quartz-hornblende diorite is not known to be associated with any of the ore deposits. Thus the same magmatic source from which the quartz-mica diorite came also supplied the agents of mineralization. Although the mineralizing solutions presumably came for the most part from depths below anything yet exposed by erosion, they rose near and in part through masses of quartz-mica diorite which are so exposed, and ore bodies are therefore to be expected only in the vicinity of such masses. Of course, there may be dioritic dikes with associated ore bodies which through the vagaries of erosion do not crop out. A study of the geologic map of the Christmas area (Pl. I) and of that of the Ray quadrangle, reproduced in part in Plate XVI, brings out some features of the distribution of the intrusions. In the Saddle Mountain district most of the dikes of quartz-mica diorite are included in an area about 4 miles long and 1½ miles wide trending about N. 70° E., which is the average strike of the dikes. The same area includes nearly all the known mineral deposits in the district and all those known to be of any value. In the Banner district the diorite bodies are more scattered, more irregular, and on the average larger. Associated with these are contact-metamorphic and similar ore bodies, a number of which are larger than the deposits of the Saddle Mountain district. In the vicinity of Troy, partly in and partly beyond the limits of the Banner district, there is another group of dikes. These lie in an area about 5 miles long by 2½ miles

wide, trending about N. 80° E., which again is about the average strike of the dikes. The Columbia mine and a number of others are in this area. The strikes of the dikes in both groups are fairly consistent within the group but are independent of the direction of the major faults in the region. The dikes and the veins associated with them appear to occupy fissures and shear zones formed subsequently to much of the faulting in the region, although some of the dikes near Troy seem to follow fissures that form parts of the regional fault pattern. The set of fractures now occupied by the dikes and ore deposits was probably produced at an early stage in the structural disturbance in which the quartz-mica diorite was intruded. Where the fractures were closely spaced the diorite magma may have split up into smaller masses than it could elsewhere. Another possibility is that under the areas of closely spaced dikes there are diorite masses larger than any exposed at the surface and that the force of intrusion of these masses caused the production of fractures in the overlying rocks through which narrow apophyses from the diorite magma were thrust.

As the ore deposits are interrelated, it follows that gradations between them may be expected. It is conceivable that a vein containing no contact-metamorphic minerals might be found to grade into a contact-metamorphic deposit where the conditions were favorable. Typical contact-metamorphic deposits can be formed only in rock susceptible of replacement of that character. In these two districts the rocks known to be suitable for such replacement comprise the purer beds of the Tornado limestone and certain beds in the Martin limestone, especially one near its base. Deposits have also been found in the Troy quartzite and in a rock supposed to belong to the Mescal limestone, but they are of less value. It is probable that valuable contact-metamorphic deposits of any great size will not be found in any of the rocks in the region other than those enumerated above, and that most of the large ore bodies of this type are confined to the Tornado and Martin limestones.

Although all the deposits were produced in the same period of mineralization they were doubtless not all formed simultaneously, and different types were probably formed at slightly different times. The contact-metamorphic deposits were doubtless produced in general at an early stage. The order of formation of the pyritic deposits, disseminated and in shear zones, and the lead-silver veins is not known. If order of deposition of the minerals in the ore is a criterion, then it would seem that the pyritic deposits are younger than those of galena and sphalerite, but other evidence is lacking.

From the facts above set forth something can be inferred as to future possibilities in the exploitation of specific deposits, and

examples of such inferences are given below. The Cretaceous strata that form the wall rock of the veins of the Saddle Mountain district are underlain by the Tornado limestone, which is cut by quartz-mica diorite, perhaps by a large mass of this rock. Consequently contact-metamorphic deposits may be expected in the limestone, but it is by no means certain that they lie under the deposits now being worked, which may be underlain by quartz-mica diorite in which there has been little or no valuable mineralization. Before passing out of the Cretaceous rocks the lead-silver veins may grade into pyritic deposits in depth, or the reverse may be true, according to which is the older. At the Christmas mine the bottom of the No. 3 shaft is probably less than 200 feet above the lower limit at which valuable ore bodies are to be expected, but on the downthrown side of the Christmas fault limestone lithologically favorable to the production of ore bodies extends to much greater depths and has not yet been explored.

MINES AND PROSPECTS

ADJUST MINE

Location.—The Adjust mine, owned by the Adjust Mining Co., is on the south side of Deer Creek in secs. 34 and 35, T. 4 S., R. 16 E., about 2½ miles by trail from Christmas station on the Arizona Eastern Railroad. A branch of the road from Winkelman through Old Mill extends to the southern base of the hill on which the mine is situated, the distance being about 8½ miles. At present supplies are brought in by automobile on this road. When ore is shipped it is taken to Christmas station by burro pack train down Deer Creek.

Property.—The Adjust Mining Co., which has recently been reorganized, has 13 unpatented claims, on which there are a number of shafts, adits, and cuts. The principal developments are on the Blue Bird vein, shown in Figure 4, where there are more than 1,500 feet of workings. About 640 tons of ore,³⁰ principally oxidized material, has been shipped. In May, 1922, after a shutdown of several months, the mine was reopened with the intention of carrying out sufficiently extensive development to determine the value of the ore bodies to a depth of 300 feet or more.

Character of the deposits.—The country rock is principally Cretaceous andesite, but there are some small dikes of gray porphyry, probably quartz-mica diorite. These dikes are too small to show on Plate I, but larger ones have been mapped northwest and southeast of the property. Several veins crop out, but only one, called

³⁰ Heron, C. M., letter dated July 12, 1922.

Location.—The Seventy Nine mine is in the Ray quadrangle, a little less than 2 miles northwest of Tornado Peak and $4\frac{1}{2}$ miles northwest of Hayden in air lines. The principal workings and mine buildings are on the northwest side of a wash at an altitude of somewhat more than 3,300 feet above sea level, and there are prospect holes along the streamway above this.

History.—The deposit was located in 1879 by Mike O'Brien and his brother Pat, but little work was done on it until December, 1919. From that date until April 27, 1922, the mine was held by the Continental Commission Co., which is reported to have shipped about 3,000 tons of ore averaging 24 per cent of lead, 1.75 per cent of copper, and 4 ounces of silver and 80 cents in gold to the ton. In April, 1922, the mine was taken over by the Seventy Nine Mining Co., and in June, 1922, the managers reported that shipments of about 50 tons a day were being made of ore averaging about the same as that previously shipped. Recently⁵² the mine has been ordered reconveyed to the Continental Commission Co. as a result of litigation.

Property.—The property comprises 23 claims.⁵³ The mine has been developed by two tunnels, one above the other, with irregular slopes from them. Most of the stoping now in progress starts from the upper tunnel. The ore is sent down chutes to the lower tunnel and trammed to the surface. There is a winze down 70 feet on a 35° slope from the lower tunnel. A short distance up the gulch from the mine are the office, boarding house, and a few other buildings. Not far from the boarding house is a prospect shaft, and a few hundred yards farther upstream is a tunnel from which a little stoping has been done. Other prospect workings are scattered over the property.

Character of the deposits.—The country rock is the Tornado limestone. In the northeastern part of the mine is a dike of altered dioritic rock flanked by fault breccia, and near the mouth of the lower tunnel some irregular masses of comparatively fresh quartz-mica diorite are exposed. The ore has been formed as replacement bodies of galena along certain beds in the limestone separated by silicified beds, perhaps originally somewhat shaly, which are also more or less mineralized in places. About five different beds of ore from 3 to 5 feet thick are reported to have been found. There appears to be more unaltered galena in the upper stopes than below, but throughout the mine a large part of the ore is cerussite. Angle-site and small amounts of azurite and malachite are also present.

⁵² Eng. and Min. Jour.-Press, vol. 115, p. 862, 1923.

⁵³ Weed, W. H., Mines Handbook, vol. 15, p. 254, 1922.

At the shallow shaft near the boarding house the ore, which is an irregular body formed by the replacement of limestone, is thoroughly oxidized. The minerals recognized are wulfenite, cerussite, limonite, andradite garnet, quartz, calcite, and a little hydrous manganese oxide and malachite. There is also a small amount of a yellow powdery material, probably an oxidized lead mineral, but it was impossible to separate enough for a conclusive test. Garnet is also reported in outcrops of some of the ore beds at the main mine. The tunnel farther up the gulch contains oxidized ore, but here copper is the metal sought, and no lead is reported to be present. The ore is stated to carry several ounces of silver to the ton. This deposit is also in the Tornado limestone but appears to have been formed along a fissure instead of along the bedding. The minerals recognized are chrysocolla, malachite, cuprite, black copper oxide, hematite, magnetite, quartz, garnet, and calcite.

COLUMBIA MINE

Location.—The Columbia mine of the Dripping Springs Mines Corporation is in a tributary gulch on the south side of Dripping Spring Valley about three-quarters of a mile west of the point where the highway to Globe swings north out of the valley and 2 miles east of Troy. It is about $18\frac{1}{2}$ miles from Winkelman and 12 miles from the terminus of the Arizona Eastern Railroad at Christmas station.

History.—Most of the claims were located by J. W. Read, part of them in 1901. The Dripping Springs Copper Co. acquired the property about 1915 and was reorganized into the present company about 1920. Most of the development work was done in this year, and some copper ore was shipped. When visited in June, 1922, development work was in progress under Mr. A. T. Copley, superintendent, who kindly furnished the historical and other data regarding the mine. In the fall of 1922 exploration by diamond drilling from the floor of the principal tunnel was started, and an ore body 10 feet wide is reported to have been encountered.⁵⁴

Property.—The Columbia group comprises about 20 claims. The Dripping Springs Mines Corporation also owns the Pearl group⁵⁵ of claims, near Mammoth. At the Columbia mine there are a number of buildings, an air compressor, and other equipment. The principal underground workings consist of a tunnel about 550 feet long with a crosscut at the end of the tunnel extending 950 feet south and 350 feet north and a shallow winze at the north end.

⁵⁴ Eng. and Min. Jour.-Press, vol. 115, p. 688, 1922.

⁵⁵ Weed, W. H., op. cit., p. 265.

Some stoping has been done, and there are several branch drifts off the crosscuts and smaller workings elsewhere.

Character of the deposits.—The rocks in this vicinity are a complex of small fault blocks of Paleozoic sedimentary rocks and intrusive diabase, with intrusions of quartz diorite porphyry and granodiorite only short distances away. Plate XVI shows the general features of the geology, but is on too small a scale to illustrate fully the complexity of the faulting. The rock mapped by Ransome as quartz diorite porphyry is probably equivalent to that called quartz-mica diorite in the present report. Most of the rock exposed underground is diabase, but there is a block of limestone supposed to belong to the Mescal limestone, and at the north end of the crosscut some metamorphosed shaly rock is exposed, perhaps also a part of the Mescal limestone. Mr. Copley states that five fissures showing mineralization are known. They strike in general nearly east and dip 70°–80° S. Gouge and altered country rock without much quartz in belts 3 to 8 feet wide occur along these fissures, widening out into the limestone where this rock forms one of the walls. Small bodies of copper carbonate ore have been found and stoped in places. Mr. Copley states that cuprita was noted in cracks in one of the veins, and small amounts of pyrite and chalcopyrite have been observed. In the altered shale at the north end of the workings a bed about 4 feet thick has been replaced by vesuvianite, magnetite, and doubtless other minerals. Picked samples from this bed are reported to contain 3 to 7 per cent of zinc, but no zinc minerals appear to be present in the specimen collected. Garnet is probably present in places, and some specularite is associated with the magnetite showing incipient replacement along octahedral partings.

McHUR PROSPECT⁵⁷

In Seventy-nine Gulch there was in 1917 a vanadium prospect owned by Mac McHur. The exact location is not known, but it is possible that the deposits are on the ground of the present Seventy Nine Mine. Irregular masses of gossanlike material containing siderite, wulfenite, and vanadinite crop out in the gulch for about a quarter of a mile, and there are shallow workings at intervals. The country rock is the Tornado limestone, with diabase coming in near by and possibly against it in places. About a mile above this is an outcrop of ferruginous quartz at the contact of limestone and diabase, called the Iron Spike vein. A little wulfenite and vanadinite and some copper staining occur here. About a mile below the McHur prospect a small mill has at one time been operated by Mr. Boykin, of Tucson.

⁵⁷ From notes by F. L. Hess, of the U. S. Geol. Survey, taken in April, 1917.

PREMIER GROUP⁵⁷

The Premier group, owned by S. O. Stewart, comprises 10 unpatented claims on the south side of Dripping Spring Valley 2 miles northwest of Christmas. The location shown on Plate X is only approximate. The deposit is developed by several pits and by an open cut about 25 feet long, with a drift from one end extending 20 feet to the east and a winze 12 feet deep at the end of the drift. The country rock is the Tornado limestone. The vein strikes about magnetic east, dips nearly vertically, and can be traced for about half a mile on the surface. A dike of quartz porphyry about 60 feet wide cuts the vein at a small angle. It is stained with iron oxide and appears to contain a little vanadinite. At the open cut mentioned the spongy gossan-like material extends out north of the vein and lies nearly flat along the bedding of the limestone. Above the drift the limestone is crushed and recemented by caliche-like material. The vein ranges from 10 inches to 3 feet in width and contains iron-stained, spongy quartz with vugs and cracks lined with thin coatings of vanadinite and wulfenite.

C. & B. GROUP⁵⁷

The C. & B. group, owned by E. E. Cutler and Calvin Bywater, is on the south side of Dripping Spring Valley, about 9 miles above its mouth. It comprises five claims and is developed by a 20-foot shaft and two tunnels, 30 and 60 feet long, with a 10-foot winze below the shorter one. The tunnels are on a vein of oxidized material, striking approximately N. 60° E. and dipping southeast along the contact between limestone and diabase, and the shaft is on a branch of the vein, also on the contact between the limestone block and diabase but striking N. 35° W. and dipping northeast. Near by is a small outcrop of red granitic rock. Movement and brecciation have taken place along the vein fissures. The vein matter is well oxidized and for the most part soft, although silicified in places. Vanadinite and wulfenite are present in the soft material and also form coatings on brecciated diabase for a width of a few feet. Cavities left by galena crystals that had been oxidized and removed were noted at one place. These contain vanadinite. There are some small veins on the property. A little ore has been mined, and some is reported to have been shipped during the war.

⁵⁷ From notes by F. L. Hess, of the U. S. Geol. Survey, taken in April, 1917.

Tr



Tog

Tqd

Ka

db

Figure 1. Schematic diagram of the experimental setup. The subject is seated in a chair and views the target through a video camera. The target is a vertical rod with a horizontal bar at the end. The subject's hand is positioned at the end of the bar. The distance between the subject's hand and the target is 100 cm. The target is 10 cm in diameter and 100 cm in height. The subject's hand is positioned at the end of the bar. The distance between the subject's hand and the target is 100 cm. The target is 10 cm in diameter and 100 cm in height.

Known fault

Probable fault

concealed fault
(covered by younger
deposits)

15^a *Dip of fault plane*

Overthrust side of
thrust fault

0° *Strike and dip of stratified rocks*

Horizontal bed

Mine


Prospect


EARLY TERTIARY?

CRETACEOUS?

MESOZOIC ?

PRE-CAMBRIAN





(CASA GRANDE)

Edited and published by the Bureau of Land Management

Base map prepared by the U.S. Geological Survey

Compiled from USGS 1:24 000 and 1:62 500-scale topographic maps
dated 1945-1968. Planimetry revised from aerial photographs

INDEX 7

1	2	3	4	5
---	---	---	---	---

steeply dipping Apache Leap Tuff of Miocene age in the north-
west corner of the quadrangle. A steep normal fault bounds the
west margin of Dripping Spring Valley and has a stratigraphic dis-
placement of more than 2,900 feet, east side down. Diamond-drill
holes 800 feet east of bedrock in the northern part of the quad-
rangle have penetrated more than 2,900 feet of Pliocene and possibly
older fanglomerate along the west margin of the valley. At the
southeastern corner of the quadrangle, a churn drill hole 900 feet
east of bedrock bottomed in gravel at 1,470 feet.

ORE DEPOSITS

The disseminated copper deposit at Ray is the major mineral
deposit in the quadrangle. The following description of the deposit
is based on recent publications of Metz and Rose (1966), and Metz
and others (1968). The Ray deposit covers an area 2 miles long in
an east-west direction, and 1½ miles north-south, bounded on the
east by the Broken Hill fault, on the west by the West End fault
(half a mile west of the quadrangle boundary), on the north by the
North End fault, and on the south by the contact between the Pinal
Schist and Ruin Granite. Mining activities began about 1870, and
since 1911 over 1½ million tons of copper, 40,000 ounces of gold,
and 4 million ounces of silver have been recovered. The Kennecott
Corp. has been mining the deposit since 1933.

The bulk of the copper thus far produced at Ray has come from
secondary, enriched chalcocite in an irregular flat blanket, a few
feet to several hundred feet thick, mostly in Pinal Schist. Overlying
the chalcocite blanket was an average of 200 feet of leached and
hematite-stained schist, which has been removed in open-pit
operations. The primary hypogene minerals are pyrite, chalcopyrite,
minor bornite and molybdenite, and traces of galena, sphalerite
and tennantite. Primary copper, occurring mainly as chalcopyrite,
averages 0.1-0.2 percent in the Pinal Schist and Granite Mountain
Porphyry and forms large bodies of ore grade (more than 0.4 percent
Cu) in the diabase.

Ore bodies in diabase sills underlie the eastern part of the Ray
secondary chalcocite deposit and extend eastward from the Diabase
fault to the Broken Hill fault. Mining in this area began recently by
an eastward extension of the Pearl Handle pit. Schist or quartzite
adjacent to ore-grade mineralized diabase is commonly poorly
mineralized. Supergene enrichment in the diabase has been negli-
gible, but in the shallower parts of the ore bodies the chalcopyrite
has been oxidized to chrysocolla, azurite, and malachite.

The hypogene mineralization was controlled by the permeability
and type of host rock and by the arrangement of faults. Copper
mineralization is greatest in highly fractured rocks. Diabase, the
best host for copper, presumably reacted more strongly with the
mineralizing fluids than did the more siliceous rocks.


The zone of supergene copper enrichment is related to major
structures, as well as to lithology and distribution of primary
sulfides. Copper was leached most thoroughly from areas rich in
pyrite, which formed sulfuric acid under oxidizing conditions and
acted as a copper solvent. The copper solutions moved downward
and laterally along structures in the schist and porphyry, but not in
the diabase, which tended to react with and precipitate the copper.
The copper precipitated at moderate depths as chalcocite replacing
pyrite and chalcopyrite.

Small deposits of copper sulfides and oxides have been mined in
and around the Rattler Granodiorite in the Troy district (southeast
corner of quadrangle), and at the Monitor mine, northeast of where
the Rustler and Broken Hill faults converge. These deposits consist
mainly of sulfides, partly oxidized, in steeply dipping faults and
fissures with quartz and calcite gangue. In the Troy district a sizable
area in the eastern part of the Rattler Granodiorite and in the
host rocks along the southern contact contains disseminations of
azurite, malachite, chrysocolla, and copper sulfides.

It is reported that approximately 200,000 tons of ore averaging
2.5 percent Cu have been extracted from the Monitor mine, and the
Rattler mine in the Mescal Limestone and diabase near the south-
east margin of the Rattler Granodiorite is estimated to contain
2,500,000 tons of material containing 1 to 1.5 percent Cu, mainly
as copper oxides.

The Ray silver mine, near the center of the quadrangle, is reported
to have produced small amounts of silver, galena, and sphalerite.
The sulfides occur along near-vertical west- to northwest-trending
faults and fissures, mainly in the Martin Limestone.

Several other showings of copper occur in the quadrangle. The
most notable occur along the faults parallel to and north of the
Rustler fault system, and along the Broken Hill fault in the central
part of the quadrangle.



Base from U.S. Geological Survey, 1964
10,000-foot grids based on Arizona coordinates

aplite, the stock cooled, fractured, and was intruded by dikes of porphyritic rock (Krp) that is intermediate in composition between granodiorite and quartz diorite.

The main rock type consists of 45-65 percent subhedral to euhedral andesine as much as 6 mm long, 15-30 percent subhedral to anhedral quartz, 2-7 mm long, 10-20 percent anhedral orthoclase, 0-13 percent subhedral biotite as much as 5 mm in diameter, 0-11 percent hornblende as much as 4 mm long, 1-4 percent euhedral magnetite-ilmenite as much as 1 mm in diameter, and accessory apatite, sphene, and zircon. K-Ar age determinations by M. L. Silberman (written commun., 1971) give an age of 70 ± 2 m.y. for a biotite separate, and 74 ± 2 m.y. for a hornblende separate.

KT TORTILLA QUARTZ DIORITE—Small stock-like masses in the southern half of the quadrangle that are cut by rhyodacite porphyry (TKr) and that intrude all rocks older than and including the Pennsylvanian Naco Limestone in the quadrangle. Many stocks of identical rock are found south of Sonora quadrangle, especially along the axis of the Tortilla Mountains. This quartz diorite, previously informally called Sonora diorite by local geologists, is here formally named the Tortilla Quartz Diorite for widespread exposures in the Tortilla Mountains in the adjacent Kearny quadrangle. The rock, although slightly variable in appearance and composition from stock to stock, is for the most part a quartz diorite, both modally and chemically. Its type section is in sec. 2, T. 4 S., R. 13 E., north of the Gila River and west of Mineral Creek in the Kearny quadrangle. Within and between separate stocks, composition ranges from pyroxene-hornblende diorite through biotite-hornblende quartz diorite (tonalite).

An abruptly gradational porphyritic facies with euhedral phenocrysts of pyroxene and hornblende up to 3 cm across is a common but not dominant rock type. The most common type is medium-gray fine- to medium-grained hypidiomorphic-granular rock, composed of approximately 50-60 percent subhedral labradorite; 25-30 percent subhedral augite, hornblende, and biotite in varying amounts; 5 percent subhedral magnetite-ilmenite; 5-10 percent interstitial orthoclase that, like quartz, commonly occurs in optically continuous masses up to 10 times the average grain size of the rock; and 5-15 percent quartz. Accessory minerals are apatite, zircon, and sphene. M. L. Silberman (written commun., 1971) has obtained K-Ar ages of 71 ± 2 m.y. and 83 ± 2 m.y. for coexisting biotite and hornblende, respectively. Although these ages are discordant the true age is presumably Late Cretaceous since both fall in this range.

Kha HORNBLende ANDESITE PORPHYRY—Occurs in two small exposures in the NW¼ sec. 30, T. 3 S., R. 14 E., where it cuts diabase (db). In the Winkelman quadrangle it is cut by rhyodacite porphyry (TKr2) (M. Kreiger, oral commun., 1970); most mafic of Laramide dikes; K-Ar age is 128 m.y. (M. L. Silberman, written commun., 1971). Phenocrysts make up 10-30 percent of rock as follows: partly glomeroporphyritic, blocky euhedral partially altered labradorite up to 2 mm long (10-25 percent), subhedral magnetite-ilmenite up to 0.5 mm in diameter (0.5-1.5 percent), euhedral and rounded glomeroporphyritic hornblende up to 1.5 cm long (0-4 percent), euhedral colorless augite up to 1 mm in diameter (0-1 percent). Groundmass comprises 70-90 percent of rock and is gray to black, aphanitic with trachytic to hyalopilitic texture, and consists of plagioclase laths (40-70 percent), anhedral sphene (3 percent), subhedral magnetite-ilmenite (6-7 percent), devitrified light-green glass (10-15 percent).

bas BASALT PORPHYRY—Occurs as sills with rare connecting dikes, usually at contacts between the Abrigo, Martin, Escabrosa, and Naco Formations, and at several horizons in the Escabrosa and near and usually below the fossiliferous limestone bed in the Martin; usually too thin to map. Thin sills and selvages on sills more than 5 feet thick are very variable in color and appearance owing to variations in degree of alteration. They range from punky yellowish-white rocks through harder gray and brown

db DIABASE—Diabase sills and some discordant bodies common to abundant in all the Precambrian rocks, but in the Mescal Limestone and Dripping Spring. The sills range in thickness from a few feet to 1,000 feet. The thicker sills are probably composite. The rock is to olive gray, fine to coarse grained with diabasic texture. The thicker sills have pegmatitic schlierenular masses. The diabase disintegrates on weathering forms slopes and flat areas veneered by yellowish-brown soil and rounded boulders. The least altered composed of laths of labradorite partly altered to kaolinite poikilitically included in augite and pigeonite plus accessory magnetite-ilmenite. The ferromineral minerals are partly altered to bowingite, antigorite, iddingsite, and magnetite-ilmenite. Much of the diabase has been altered to some degree, with the development of hornblende, biotite, quartz, epidote, and pumpellyite. Diabase contains graphic intergrowths of quartz and K-feldspar.

t TROY QUARTZITE (300-800 ft)—The Troy Quartzite Sonora quadrangle and surrounding area has been named by Shride (1967, p. 44-46) with the Chediski Sandstone of the formation, and the three sections we measured to test this correlation. On the top of Troy Mountain, however, the formation is 800 feet thick, remnants of an upper member (Shride, 1967, p. 51-52). The main body of this quadrangle is quartzite and sandstone that range from grayish pink and white to light brown and gray and consist of medium to coarse subangular to rounded grains with variable amounts of feldspar, limonite, and dolomite. The quartzite is interbedded in varying proportions with thin layers and lenses of poorly to well-sorted granule conglomerates. Matrix is siliceous with sandstone. Beds range in thickness from laminated to very thin, arranged in thin to thick composite tabular to lenticular. Crossbedding is locally prominent.

The base of the formation consists of 5-10 feet of coarse-grained sandstone with angular to rounded pebbles and cobbles of which several types of sandstone and quartzite, Mescal Limestone, chert, red and pink jasper, basalt, and Pinal Schist are included. Sorted reddish-orange to light-brown matrix, medium-grained, well cemented, siliceous and ferruginous. Conglomerate beds, 2-6 feet thick, are also scattered throughout the formation, particularly toward the top. The predominant lithology of the Troy form bold cliffs with intervening slopes covered by more friable sandstone.


APACHE GROUP:

The Apache Group includes the Pioneer Formation, Spring Quartzite, Mescal Limestone, and basalt.

b BASALT (0-100 ft)—One or more flows of porphyritic basalt, grayish to blackish red or brown, with vesicular, and some with columnar jointing. Phenocrysts are plagioclase, partly glomeroporphyritic, 2-8 mm long. Groundmass is grayish or blackish red, fine grained, and consists predominantly of plagioclase, 0.1-0.3 mm long in intersertal relations with pyroxene and magnetite. Plagioclase has composition of about An₅₀ basalt has been intensely altered to chlorite, sericitite, epidote, quartz, hematite, kaolinite, limonite and feldspar; pyroxene and olivine have been almost completely replaced.


m Mescal LIMESTONE (270-340 ft)—Formation composed of pale-pink to light-brown and brownish-gray dolomitic calcareous, thinly laminated to thin-bedded, submicritic to fine grained, partly medium to coarse grained. Black and pinkish-gray chert, aphanitic to microcrystalline, abundant in some beds as uneven layers, lenses, and nodules. Matolitic algal beds are abundant above the middle of the formation, and massive bedlike layers of sandstone and

upthrown side. Sawteeth on
plate of thrust fault



Anticline

Syncline



Horizontal beds



Pearl Handle pit

x


Prospect pit

DDH
2919'

Diamond drill hole
Showing vertical depth

CDH
1470'

Churn drill hole
Showing vertical depth





5000'
- 4000'
- 3000'
- 2000'
- 1000'

ES, ARIZONA

percent), magnetite-ilmenite (5 percent), K-feldspar partially as plagioclase replacement (10 percent); accessory sphene, apatite, and zircon.

Kr RATTLER GRANODIORITE—A pear-shaped stock in the south-eastern corner of the quadrangle; cuts Paleozoic sedimentary rocks and is cut by many of the Laramide dikes. It is here named the Rattler Granodiorite for exposures in the vicinity of the Rattler mine, sec. 26, T. 3 S., R. 14 E., its type locality, on the south-center margin of the intrusion. The intrusive is composite and ranges from quartz diorite to sodic granite aplite. The main rock type is granodiorite, granitic in texture, and fairly constant in composition except for the amount of hornblende and biotite, which vary inversely. The main rock type grades into very small patches of a melanocratic facies (Krm), and into a porphyritic facies that is chemically identical with the main rock and is found in the eastern lobe of the stock. Prior to the intrusion of



GEOLOGIC QUADRANGLE MAP
SONORA QUADRANGLE, ARIZONA
GQ-1021

fractured, and was intruded by dikes
p) that is intermediate in composition
quartz diorite.

sts of 45-65 percent subhedral to euhedral
mm long, 15-30 percent subhedral to
n long, 10-20 percent anhedral ortho-
hedral biotite as much as 5 mm in
hornblende as much as 4 mm long, 1-4
etite-ilmenite as much as 1 mm in
apatite, sphene, and zircon. K-Ar age
L. Silberman (written commun., 1971)
y. for a biotite separate, and 74 ± 2 m.y.

ORITE—Small stock-like masses in the
quadrangle that are cut by rhyodacite por-
trude all rocks older than and including
Limestone in the quadrangle. Many
are found south of Sonora quadrangle,
of the Tortilla Mountains. This quartz
ormally called Sonora diorite by local
ally named the Tortilla Quartz Diorite
es in the Tortilla Mountains in the adja-
The rock, although slightly variable in
tion from stock to stock, is for the most
h modally and chemically. Its type section
3 E., north of the Gila River and west of
earny quadrangle. Within and between
osition ranges from pyroxene-hornblende
ornblende quartz diorite (tonalite).

porphyritic facies with euhedral pheno-
ornblende up to 3 cm across is a common
type. The most common type is medium-
grained hypidiomorphic-granular rock,
ately 50-60 percent subhedral labra-
ubhedral augite, hornblende, and biotite
percent subhedral magnetite-ilmenite;
orthoclase that, like quartz, commonly
nuous masses up to 10 times the average
; and 5-15 percent quartz. Accessory
con, and sphene. M. L. Silberman (writ-
obtained K-Ar ages of 71 ± 2 m.y. and
ing biotite and hornblende, respectively.
e discordant the true age is presumably
oth fall in this range.

TE PORPHYRY—Occurs in two small
sec. 30, T. 3 S., R. 14 E., where it cuts
Vinkelman quadrangle it is cut by rhyo-
(M. Kreiger, oral commun., 1970); most
; K-Ar age is 128 m.y. (M. L. Silberman,
). Phenocrysts make up 10-30 percent
rtly glomeroporphyritic, blocky euhedral
lorite up to 2 mm long (10-25 percent),
nenite up to 0.5 mm in diameter (0.5-1.5
rounded glomeroporphyritic hornblende
percent), euhedral colorless augite up to
percent). Groundmass comprises 70-90
gray to black, aphanitic with trachytic to
l consists of plagioclase laths (40-70 per-
(3 percent), subhedral magnetite-ilmenite
d light-green glass (10-15 percent).

Occurs as sills with rare connecting dikes,
ween the Abrigo, Martin, Escabrosa, and
at several horizons in the Escabrosa and
w the fossiliferous limestone bed in the
n to map. Thin sills and selvages on sills
are very variable in color and appearance
degree of alteration. They range from
rocks through harder gray and brown

db DIABASE—Diabase sills and some discordant bodies are com-
mon to abundant in all the Precambrian rocks, but especially
in the Mescal Limestone and Dripping Spring Quartzite.
The sills range in thickness from a few feet to 1,000 feet or more.
The thicker sills are probably composite. The rock is dark gray
to olive gray, fine to coarse grained with diabasic to ophitic
texture. The thicker sills have pegmatitic schlieren and irreg-
ular masses. The diabase disintegrates on weathering and
forms slopes and flat areas veneered by yellowish-brown gran-
ular soil and rounded boulders. The least altered diabase is
composed of laths of labradorite partly altered to sericite and
kaolinite poikilitically included in augite and pigeonite, and
olivine plus accessory magnetite-ilmenite. The ferromagnesian
minerals are partly altered to bowlingite, antigorite, limonite,
iddingsite, and magnetite-ilmenite. Much of the diabase has
been altered to some degree, with the development of horn-
blende, biotite, quartz, epidote, and pumpellyite. Some
diabase contains graphic intergrowths of quartz and K-feldspar.

t TROY QUARTZITE (300-800 ft)—The Troy Quartzite in the
Sonora quadrangle and surrounding area has been correlated
by Shride (1967, p. 44-46) with the Chediski Sandstone Member
of the formation, and the three sections we measured support
this correlation. On the top of Troy Mountain, however, where
the formation is 800 feet thick, remnants of an upper medium-
grained very light gray vitreous quartzite resemble the quartzite
member (Shride, 1967, p. 51-52). The main body of Troy in
this quadrangle is quartzite and sandstone that range in color
from grayish pink and white to light brown and grayish orange
and consist of medium to coarse subangular to rounded sand
with variable amounts of feldspar, limonite, and dark acces-
sories. The quartzite is interbedded in varying proportions
with thin layers and lenses of poorly to well-sorted pebble and
granule conglomerates. Matrix is siliceous with some clay.
Beds range in thickness from laminated to very thin but are
arranged in thin to thick composite tabular to lenticular sets.
Crossbedding is locally prominent.

The base of the formation consists of 5-10 feet of conglomerate
with angular to rounded pebbles and cobbles of white quartz,
several types of sandstone and quartzite, Mescal Limestone,
chert, red and pink jasper, basalt, and Pinal Schist in a poorly
sorted reddish-orange to light-brown matrix, medium to coarse
grained, well cemented, siliceous and ferruginous. Conglom-
erate beds, 2-6 feet thick, are also scattered through the for-
mation, particularly toward the top. The predominant quartz-
ites of the Troy form bold cliffs with intervening slopes under-
lain by more friable sandstone.

APACHE GROUP:

The Apache Group includes the Pioneer Formation, Dripping
Spring Quartzite, Mescal Limestone, and basalt.

b BASALT (0-100 ft)—One or more flows of porphyritic basalt,
grayish to blackish red or brown, with vesicular, amygdular
tops. Phenocrysts are plagioclase, partly glomeroporphyritic,
2-8 mm long. Groundmass is grayish or blackish red or brown,
fine grained, and consists predominantly of plagioclase laths
0.1-0.3 mm long in intersertal relations with pyroxene, olivine,
and magnetite. Plagioclase has composition of about An_{50} . The
basalt has been intensely altered to chlorite, sericite, calcite,
epidote, quartz, hematite, kaolinite, limonite and some K-
feldspar; pyroxene and olivine have been almost completely
replaced.

m Mescal Limestone (270-340 ft)—Formation consists of
pale-pink to light-brown and brownish-gray dolomite, partly
calcareous, thinly laminated to thin-bedded, sublithographic
to fine grained, partly medium to coarse grained. Black, white,
and pinkish-gray chert, aphanitic to microcrystalline, is abun-
dant in some beds as uneven layers, lenses, and nodules. Stro-
matolitic algal beds are abundant above the middle of the for-
mation and massive bedlike layers of sandstone and quartzite

RENFRO GROUP

PINAL & GILA COUNTIES

See: BANNER DISTRICT MISCELLANEOUS (Geology File)

Approx. Sec. 25, 26, T3S, R14E

Dripping Springs Dist

PRELIMINARY REPORT ON RENFRO GROUP.

INTRODUCTION

The facts that neither a topographic map of the area covered by this report, nor satisfactory surveys of the claims themselves, were available, rendered its study difficult and conclusions tentative. Never the less, I shall attempt to depict the salient geological features in their bearing on ore deposition. The interpretation of the relations of the main rock masses is made easy by the rugged topography, bare slopes and numerous well defined contacts. As above stated the absence of an adequate map alone makes the presentation vague.

TOPOGRAPHY.

Rising precipitately from Dripping Springs Wash at a point where the canon begins to box, a continuous ridge comprised of thick bedded quartzites and limestones runs in a northerly direction and forms in general way a natural eastern boundary to the area. To the north further high ridges predominate. Westerly a series of smaller, usually rounded hillocks fade into the peneplain on which Troy is situated. In a southerly direction the Dripping Springs Wash defines the area. Erosion has carved a deep wash, which emanating from the ridges to the south has to the east exposed the entire sedimentary series, while to the west and south denudation has almost completely removed them. This eroded vally forms in a general way the center line of the area and enters the Dripping Springs Wash. Minor depressions characterize the steep eastern slope usually instigated by lines of faulting. As a result of the weathering of quartzites the slopes are usually covered with small angular fragments of rock. The relatively more rapid weathering of the limestone overlying the quartzite often throw the latter rock into sharp relief and produce steep projecting promontories. A large mass of intrusive diabase now exposed by erosion produces smoothly rounded slopes.

GENERAL GEOLOGY.

The sedimentary rocks attain a total thickness of some 800 feet and comprise quartzites, quartzite conglomerate, limestones and highly metamorphic slates and schists. The quartzite occupies nearly one half of the sedimentary area and was derived from the metamorphism of thick bedded, fine grained sandstone, although conglomeratic developments occur locally. It is overlain by 80-ft of fine bedded sediments originally consisting of alternating shale and limestone, with occasional quartz or sandy layers. Metamorphic processes have altered the shale partly to chlorite schist, the limestone to talc and other hydrous silicates of magnesia, and the sandstone to quartzite. Capping this heterogeneous series and attaining in greatest development a thickness of 325-ft there obtains the massive, compact, dark colored dolomitic limestone series that forms the crest of all the higher ridges.

Beneath the quartzite there appears locally and eminently in the northern half of the area, a lenticular mass of older sediments composed of highly altered limestones and schists resting unconformably beneath the other sediments. This local series, a remnant left by erosion in the sea of igneous rock will be considered more in detail later.

The igneous rocks comprise granites, granite porphyry and diabase. The two former are strongly developed in the southern part of the area and are different textural phases of the same parent magma. They are the oldest igneous rocks in the region and were intruded while the area was still buried under the thicker mass of sediments. Portions of the magma were squeezed upward through the sedimentary rocks along lines of weakness, giving rise by more rapid cooling to porphyritic phases. The normal granite is a relatively fine grained, holocrystalline mass of quartz, orthoclase and biotite. Weather has either altered the mica to chlorite, or removed it entirely, giving the rock a vesicular appearance. The feldspar is quite often changed to sericite. At a considerably later period the region was invaded by a batholithic intrusion of diabase, which spread from the central stock in generally easterly and westerly directions, along lines of faulting produced during the earlier granitic intrusions. Where exposed by erosion it presents every textural gradation from a coarse crystalline, ophitic aggregate of plagioclase feldspar and augite to a compact, crypto-crystalline greenish mass. It was a most important mineralizing agent.

The combined effect of both plutonic invasions produced a tilting of the sediments to an angle of 20 to 30 degrees to the southeast, developed a series of approximately east-west fractures, and profoundly modified the chemical and mineral

characters of the sedimentary rocks. And it is along these fractures thus formed that ore bearing solutions have made their ingress. As mentioned under topography, erosion has to the north entirely removed the sedimentary series, in the central area exposed the diabase, while to the south the thick sedimentary capping still remains. For convenience in the discussion of the ore deposits, I shall group the forty-three claims under consideration as follows:

- (a) Those within the granite area
- (b) Those lying in the diabase zone
- (c) Those capped with sedimentaries

ORE DEPOSITS.

(a) Under this heading I include the Outlet, Los Angeles, Anaconda, Port Arthur, Saar and Garter, W.E. Bryan and Paris, and to a lesser degree the Mother Lode, and East Extension of the Bryan. All of those are full claims. The major portion of the included area is granite, with very subordinate diabase to the north, and quartzite to the east. Approximately 300-ft of development work has been done, of which nearly half is on the Port Arthur. The deepest opening is an inclined shaft 43-ft deep. The ore is developed along a series of fractures in the granite, striking north 70 degrees west, and dipping quite steeply to the north. These fractures are true faults although of relatively slight displacement, and not mere joint planes in the granite. The ore consists almost wholly of chrysocolla with subordinate cuprite. The fractures are mineralized for several inches on each side and samples taken at various points showed copper content from three to fifteen percent. The oxidized ore persists to the deepest workings. How extensive these deposits are cannot be predicted from the present shallow openings. The period of mineralization was earlier than on claims subsequently to be described and the character of the ore and gangue different. I would strongly suggest sinking a shaft preferably in the northeast corner of the Port Arthur claim on the quite prominent fault that passes through that point. This shaft would demonstrate the persistence of this type of ore body and determine that this mineralization which covers a large area is practically a continuation of the Rattler ore body on the Troy Arizona property adjoining the group. I regard these claims as a very favorable group in themselves and undoubtedly will warrant extensive development.

(b) Under this heading I shall group the West Extension of the Bryan, Oversight Cedar Springs Extension and a second eastward extension of the Cedar Springs, Copper Boom, Copper Fleece and Promotor. Also the Eastward Extension of the Prince of Wales and the Kentucky. All are full claims with the exception of the Oversight and the Kentucky. The former is a fraction 600 x 600 feet; the latter a small triangular section in the southwest corner of the Prince of Wales. The predominant rock mass is diabase, which intrudes itself into the granite to the north, and the quartzite to the east and south. From the field evidence it seems quite certain that the intrusion represents one single eruption. It tilted the sediments to the south and induced tensional strains and resulted in extensive fracturing and displacement of the sedimentaries along eastward planes. And it is along this series of major fractures seen in their strongest development to the southwest that the two large orebodies have been opened up, as will be considered more fully later. It is due to hot ascending solutions rising along these fault planes and metasomatically replacing limestone with ore that copper minerals have been introduced. And since in the area under consideration the sedimentaries have been largely removed, the importance from the economic standpoint hinges mainly on the persistence of those major fractures in the diabase and the degree to which precipitation has taken place from ascending solutions. In Globe the conditions are very much the same, and at that place the best development of sulphides has been along similar fractures and entirely in the diabase, this rock being apparently as favorable to ore deposition as the overlying sediments. Present development on the claims under consideration seems to make conditions analogous. Along the line of fracture there has in every case been developed masses of magnetite, hematite, calcite together with the copper minerals and from the abundance and richness of the latter it seems quite advisable to develop this group of claims beyond their present prospect stage.

In particular I would suggest the very careful prospecting of the Oversight claim and further work on the second eastward extension of the Cedar Springs.

(c) The third and by far the most important group of claims both from the standpoint of present development and immediate possibilities include the Cedar Springs No 2, Golden Rule, Roman Eagle, Good Luck, Old Year, New Year, Independent, Helen, Ashler, Golden Fleece and Uncle Sam. All with the exception of the Good Luck are full claims.

Development work embraces 220 feet of tunnels and adits on the Roman Eagle, a 40 ft shaft on the New Year and considerably more than the required amount of assessment work on the remainder. All lie eminently in the zone of sedimentary rocks. Three prominent and roughly parallel fractures strongly mineralized and accompanied by extensive replacement of the adjacent sedimentaries cross the area. The most powerful and northerly strikes in a northerly direction north 75° degrees west across the Roman Eagle, Golden Rule, and Cedar Springs No. 2. Some 400 ft south a second fracture determines the orebody in the New Year, Old Year and Good Luck. A third crosses the Helen and Independent. All three faults are normal with the downfall on the south or hangingwall side. I propose to discuss the features of the Roman Eagle fault at some length and the same remarks will apply with modification to the others. The fracture is exposed by a 150 ft tunnel driven along the hanging wall to a point 250 feet from the apex of the hill. At the foot of the 150 ft tunnel a winze 60 ft has been sunk on ore, while the lower tunnel is at present being extended to intercept the winze. A shorter tunnel exposed the footwall at a 25 foot lower elevation. Three short adits open up the foot wall immediately above the main tunnel. The foot wall rock is all of limestone. The hanging wall rock is all schistose material. The total displacement is approximately 80 feet. From three to six inches of gouge indicate the strength of the faulting movement. The ore consists of massive cuprite and chrysocolla disseminations in a calcareous gangue. The former mode predominates on the hanging walls, the latter on the foot walls. Ten feet of high grade ore lie exposed on the footwall and unquestionable 100 tons of ore are now ready for shipment. This claim is beyond the prospect stage. The ore body is characteristic of replacement deposits similar to Globe, Bisbee and other Arizona camps. The ore bearing solutions ascending along fault fissures have precipitated their metallic content either in the fissure or in the adjacent rocks. At a lower level any shaft must pass through a considerable thickness of quartzite but aside from the tightening of the fissure I see no reason for diminution in the value. This vein and its westward equivalent on the Golden Rule should be energetically developed.

CONCLUSIONS.

I have attempted to demonstrate in the above report that from a geological standpoint the included area is favorable for the development of copper deposits of considerable magnitude. That the conclusions drawn should be verified by a more elaborate study with the aid of maps I deem essential. It is a significant fact that mines of proven value exist both east and west along the line of the general system of fracturing.

Respectfully submitted,

(Signed) I. Win E. Adams, M.E.

Pinal Development Company

INCORPORATED UNDER THE LAWS OF ARIZONA

Capital Stock
\$1,500,000.00
Par Value
\$1.00

OFFICERS

PRESIDENT-TREASURER

J. C. DEYER

RAY, ARIZONA

VICE PRESIDENT

CHARLES ROBERTSON

RAY, ARIZONA

SECRETARY

C. E. CARPENTER

RAY, ARIZONA

GENERAL OFFICES

Ray

Arizona

WE want this prospectus to catch the eye, rivet the attention, and appeal to the best judgment of just one, big, broad-guaged business man, a man with enough discrimination in his make up to see an opportunity and sufficient push and enterprise to take advantage of it.

**FORTUNES ARE MADE,
not SAVED. A lifetime of
SAVING will not make you
as MUCH as ONE GOOD
I N V E S T M E N T .**

The Pinal Development Company

THE COMPANY

IS a newly organized corporation, incorporated under the laws of Arizona, with an authorized capital stock of One Million Five Hundred Thousand Dollars, divided into One Million Five Hundred Thousand shares of the par value of one dollar each. The stock is fully paid and non-assessable and all stock-holders of this corporation and their private property shall be exempt from the corporate debts and obligations of this corporation.

MANAGEMENT

The Board of Directors comprise the following:

**MR. J. C. DEVINE
MR. C. T. CARPENTER
MR. E. M. BLAKE
MR. J. H. ROBINSON
MR. R. H. BOXALL
MR. CHARLES HOLLISTER
MR. F. C. NORMAN**

Mr. Devine is President and Treasurer and is personally directing the development of the property, and stock-holders are assured that not only will every cent go in the ground, but that the Company will have the advantage of Mr. Devine's knowledge and experience in economical management. The Company is ably directed. Mr. C. T. Carpenter is Secretary. The members of the board of directors are men of extensive and varied mining experience and men of executive ability. The Company very respectfully refers to the Gila

Arizona mines during
year 1915 produced
450,000,000 pounds
:: of copper ::

Pinal Development Company

Developing

The Renfro Group of Mining Claims

LOCATION

IN the Mescal Mountains, 4000 feet above the sea, and lying in the County of Pinal, State of Arizona, is the Dripping Springs Mining District, a highly mineralized belt, copper predominating, the course of which embraces such large producing properties as Ray Consolidated Copper Co., Globe, Miami, Superior and others. **THE RENFRO GROUP OF MINING CLAIMS** is situated in this district and is being energetically developed by the Pinal Development Company. This property is four miles from the Arizona & Eastern R. R., being accessible by a good automobile road direct to the portal of the mine, and is fifteen miles from the smelter of the American Smelting & Refining Company, at Hayden, Arizona.

TOPOGRAPHY

The Renfro property consists of forty-three claims, or an area of about 860 acres. A general outline of the topography of the claims herein follows: Rising precipitately from Dripping Springs Wash, at a point where the canyon begins to box, a continuous ridge comprised of thick bedded quartzites and limestone runs in the southerly direction and forms in a general way a natural eastern boundary to the property. To the north, further high ridges predominate. Westerly a series of smaller, usually rounded hillocks grad-

**Invest in Copper.
Invest with new
Corporations.
Invest with Pinal
Development
Company.**

In Conclusion

THIS is a very attractive proposition and is placed before the public, not as a prospect for it is far beyond the prospect stage, but as a property only requiring sufficient development to make it another large producing Arizona Copper Mine. Without hesitation it may be said that as far as humanly possible with any mines in their early stages, the element of risk has been eliminated.


Therefore, to those desiring to invest in a good mining stock, we feel that in offering you this opportunity that not only are we expeditiously developing the mine but are eventually returning to the investor many times his investment. We respectfully invite all desiring to personally look over the ground to communicate with us and we will make arrangements accordingly. If there is any information you may desire that is not contained in this prospectus, kindly let us forward same to you.

Respectfully Submitted,

PINAL DEVELOPMENT COMPANY.


J. C. DEVINE,
President.

C. T. CARPENTER,
Secretary.



RENFRO GROUP.

Scale, 1"=1000.'





PINAL DEVELOPMENT COMPANY

RIVERSIDE MINING DISTRICT
PINAL AND GILA COS.

SCALE: 1 INCH = 600 FEET

JAN. 1914

DEPARTMENT OF MINERAL RESOURCES
STATE OF ARIZONA
FIELD ENGINEERS REPORT

Mine **RATTLER GROUP OF COPPER CLAIMS**

Date **April 17, 1944**

District **Riverside**

Engineer **A. Macfarlane**

Subject: **Report**

Owner: **H. R. Scott, Box 2893, Globe, Arizona**

Property & Location: Consists of 11 unpatented claims all contiguous, situated on the eastern slope of the Dripping Springs range, and about 2 miles westerly from the Dripping Springs wash, and adjoins the southeasterly limits of the old village of Troy.

The elevation at that point of the road crossing the creek or wash is 2,800 feet while the elevation at the main tunnel of the mine registered 3,500 feet a climb of 700' in this 2 miles.

The mining district of the locality is known as the Riverside and adjoins the south end of the Pioneer district, Pinal County, Arizona

History: These claims were formerly part of the Old Troy mine, discovered and worked many years ago, then in part abandoned by former owners, the present ownership being by location within the past few years.

Record of Ore Output: None of the old records are available now, but examination of the mine openings leads me to state that several hundred tons of hand sorted or selectively mined ores have been marketed.

The present owner has within the past few months, mined and trucked 3 small carlots, the settlement sheets giving copper grade ranging from 3.5% to over 5% copper.

Transportation Roads: From the International smelter at Miami, major and State highways pass within 5 miles of the mine and three or four miles up the Dripping Springs valley are county maintained. The last 2 miles to reach the cabins of the mine are broken with steep grades and much rocky terrain, passable with care for trucks of small capacity.

This last section of the road can be improved at moderate cost. The total distance from smelter to mine is about 36 miles. Another road from the buildings of the Troy village leads westerly and down grade for 6 or 7 miles to Ray Junction. Mr. Scott states that repairs are required to reopen this old road.

Mine Workings: Three tunnels driven from the northeast hill slope in a more or less southwesterly direction have cut the mineral system, and on which some drifting and stopping has at past time been done.

The general course of the stratas are from west to east and dip at varying degrees towards the south and as this examination was made for the purpose of ascertaining the feasibility of immediate copper production, the writer, after looking over the lower surface, decided that the vein as cut in the upper or Session tunnel, offers the best opportunity for copper production.

Session tunnel workings consist of an adit driven for about 100' southerly into the steep hillside, and the vein was cut at this point; thence a winze was sunk to

April 17, 1944

connect with the intermediate tunnel which is about 130' lower than the Session tunnel.

The vein where cut by the adit and at the top of this winze is fully 4' in width and at this point shows copper content stated by Mr. Scott to assay from 2.5% to about 4%.

The vein should be drifted on and an occasional upraise as lateral work of exploration is made, extending both easterly and westerly from the adit. I feel that this exploration will be justified by opening up an upper level along this horizon of the mineralization and may make available a substantial tonnage of copper.

Three or four hundred feet of development will require the installation of an air mill and compressor plant; this also for the subsequent stoping of the ores, together with ore bin, road improvement, mine trackage, and air pipe line to drilling faces.

Vein Structures: Surface croppings are plainly evident and surface cuts made show continuity of the mineralization. The lower tunnel fully 350' under the general surface of the vein proves the vein structure at that depth, but ores of a lower grade than the Session tunnel zone were opened by the lower explorations, and for that reason I suggest the mining be first directed along the upper ore area.

Geological Features: Primarily a limestone belt has through alterations and powerful intrusive action tilted the limes and formed bodies of the limestones irregular to dip, but generally with the long axis towards the northwest. The first intrusions of granite later altered and in turn in part dislocated by other igneous forces provided fractures and joints, wherein subsequent mineralization could deposit in the original form of cold and deeper hot solutions.

The immediate locality within this geological area is fertile with unconformable rock formations and veins and gashes.

Mineralization both of copper, silver, gold and vanadium has been discovered in mostly small but variable quantities within this immediate area. Only more extensive explorations may prove the value of the belt as an ore producer.

Tonnage: Many thousand tons of generally from 1 to 2% copper content are visible in the two lower tunnels and extending up to the surface, but as this report only deals with the possibility of mining copper of 3% and upwards, a grade required to cover all costs, the present visible exposures are now classed as Probable or Inferred.

Development of the drifts from near the heading of the Session tunnel and upraises therefrom should place a smelter grade of ore available.

Department of Mineral Resources
Globe Field Office

SUMMARY REPORT OF RATTLER MINE

February 27, 1945

Management

Mr. H. R. Scott of Globe, Arizona, is known as an experienced and industrious miner and small mine operator, who, being the owner of the Rattler copper property, proposes to manage the mining and leaching of the ores, assisted by a man with assaying experience.

Property Description

Consists principally of an adit tunnel and drifts, upraises, incline and winzes therefrom; in all several thousand linear feet of underground workings, the development of two copper bearing stratas. (See mine maps).

Plant

Just below the floor of the above stated adit tunnel is now installed a pilot copper leaching plant consisting of 3 Redwood tanks, diameter 10' x 5' in height, and just underneath the tank floors are two lines of compartmented Redwood precipitating boxes filled with cleaned cans and iron scrap.

A small sump just lower than the last scrap iron box is equipped with a small acid proof pump for the return of the solutions to top of the 3 leaching tanks.

It is proposed by the owner to build a copper leaching tankage with rock cement tanks with an 8-hour capacity of 50 tons, this proposed plant to consist principally of:

2 stone cement thick walled and floored tanks 20' wide by 25' to 30' long. Constructed with doors of plank and some type of packing sheet at the "ends centers" of the stone tanks, in order to discharge the leached and percolated tailings by slusher scraper to dump.

An 8" x 8" timber bridge floor above center of the 2 stone tanks, whereon is to be installed a jaw crusher of type to crush to 1/2" to 1" size, and high enough above the tanks to allow of their filling directly by distributing launder from under crusher jaw.

The present pilot tanks will serve as water and solution storage above the leaching. Beneath the leaching tank floors will be placed the scrap iron precipitation boxes and the sump with an acid proof pump for the return of all solution to mill head. On the level of the sump and precipitating boxes will be the drying and clean up floor and facilities.

DEPARTMENT OF MINERAL RESOURCES

REPORT

Of The RATTLER GROUP OF COPPER CLAIMS

H. R. Scott Owner, Box 2893 Globe, Arizona.

Globe, Arizona. April 17th, 1944

Property & Location ; Consists of 11 unpatented claims all contiguous, situated on the eastern slope of the Dripping Springs range, and about 2 miles westerly from the Dripping Springs wash, and adjoins the south-easterly limits of the old village of Troy.

The elevation at that point of the road crossing the creek or wash is 2,800 feet while the elevation at the main tunnel of the mine registered 3,500 feet a climb of 700' in this 2 miles.

The mining district of the locality is known as the Riverside and ~~there are 11 unpatented claims in the Rattler group.~~

adjoins the south end of the Pioneer district, Pinal Co. Ariz
History; These claims were formerly part of the Old Troy mine, discovered and worked many years ago, then in part abandoned by former owners, the present ownership being by location, within the past few years.

Record Of Ore Output; None of the old records are available now, but examination of the mine openings, leads me to state that several hundred tons of hand sorted or selectively mined ores have been marketed.

The present owner has within the past few months, mined and trucked 3 small carlots, the settlement sheets, giving copper grade ranging from 3.5% to over 5% copper.

Transportation Roads ; From the International smelter at Miami, major and State highways pass within 5 miles of the mine and three or four miles up the Dripping Springs valley, are County maintained, the last 2 miles to reach the cabins of the mine are broken with steep grades and much rocky terrain, passable with care for trucks of small capacity.

This last section of the road can be improved at moderate cost; the total distance from smelter to mine is about 36 miles. Another road from the buildings of the Troy village, leads westerly and down grade for 6 or 7 miles, to ray Junction, Mr Scott states that repairs are required to reopen this old road.

Mine Workings ; Three tunnels driven from the northeast hill slope in a more or less southwesterly direction, have cut the mineral system, and on which some drifting and stoping has at past time been done.

The general course of the ~~structure~~ *strata are* from west to east and dip at varying degrees towards the south and as this examination was made for the purpose of ascertaining, the feasibility of immediate copper production, the writer after looking over the lower surface, decided that the vein as cut in the upper or Session tunnel, offers the best opportunity for copper production.

Session Tunnel workings consists of an adit driven for about 100' southerly into the steep hill side, and the vein was cut at this point; thence a winz

RATTLER GROUP OF COPPER CLAIMS

Page 2

was sunk to connect with the intermediate tunnel which is about 130' lower than the Session tunnel.

The vein where cut by the adit and at the top of this winze is fully 4' in width and at this point shows copper content stated by Mr Scott to assay from 2.5% to about 4.5%

The vein should be drifted on and an occasional upraise as lateral work of exploration be made, extending both easterly and westerly from the adit. I feel that this exploration will be justified by opening up an upper level along ~~at~~ this horizon of the mineralization, may make available a substantial tonnage of copper.

Three or four hundred feet of development, will require the installation of an air drill and compressor plant, this also for the subsequent stoping of the ores, together with ore bin, road improvement, mine trackage, and air pipe line to drilling faces.

mineralization
Vein Structures ; Surface croppings are plainly evident and surface cuts made shows continuity of the ~~mineralization~~; the lower tunnel fully 550' under the general surface of the vein, proves the vein structure ~~to~~ that depth, but ores of a lower tenor than the Session tunnel zone, were opened by the lower explorations; and for that reason, I suggest the mining be first directed along the upper ore area.

Geological Features ; Primarily a limestone belt has thru alterations and powerfull intrusive action ~~has~~ tilted the limes and formed bodies of the limestones irregular as to dip, but generally with the long axis towards the northwest. The first intrusives of granite later altered and in turn in part dislocated by other igneous forces, all provided fractures and joints wherein subsequent mineralization, could deposit, both in the original form of cold and deeper hot solutions.

The immediate locality within this geological area is fertile with unconformable rock formations and veins and gashes.

Mineralization both of copper, silver, gold and vanadium has been discovered in mostly small but variable quantities, within this immediate area, only more extensive explorations may prove the value of the belt, as an ore producer.


Ore Tonnage ; Many thousand tons of generally from 1 to 2% copper content are visible in the two lower tunnels and extending up to the surface, but as this report only deals with the possibility of mining copper of 3% and upwards, a grade required to pay all costs, the present visible exposures are now classed as Probable or Inferred ore.

The extending of the drifts from near the heading of the Session tunnel and upraises therefrom, should place a smelter grade of ore available.

Department Of Mineral Resources
Globe Field Office.

DEPARTMENT OF MINERAL RESOURCES

REPORT TO OPA ON ACTIVE MINING PROJECT



Date May 28 1945
 Name of Mine Rattler
 Owner or Operator H.R. Smith
 Address Box 2893
 Mine Location Troy Gila Co. Ariz.

File System.....
 File No.....
 This chart to be used for gallons of gasoline required per month.

PRESENT OPERATIONS: (check X)

Production ☒; Development ☒; Financing.....; Sale of mine.....;
 Experimental (sampling) ☒; Owner's occasional trip ☒;
 Other (specify).....

PRODUCTION: Past and Future.

Tons

Approx. tons last 3 months
 Approx. present rate per 3 months 350
 Anticipated rate next 3 months 500
 If in distant future check (X) here

EQUIPMENT OPERATED:

Type	Quantity or Horse Power	Hours Per Month	Gallons Required Per Month
Personal Cars
Light or Service Trucks
air comp. Compressors	<u>40</u>	<u>220</u>	<u>200</u>
Compressors <u>crusher</u>	<u>15</u>	<u>220</u>	<u>210</u>
Other Mine or Mill Eqp. <u>Pump</u>	<u>3</u>	<u>240</u>	<u>80</u>
			<u>490</u>

PRODUCT PRODUCED OR CONTEMPLATED: Name metals or minerals.

Copper

REMARKS:

Operating Copper Leaching mill & mining. Processing ore 20 tons daily
Recommend for 3 mos. 1500 tons

ARIZONA DEPARTMENT OF MINERAL RESOURCES

By

A. Macfarlane

Market

The Hayden copper smelter situated 30 miles southerly on good road is a purchaser of cement or copper sludge, as well as ores and concentrates. The International smelter near Miami, Arizona, distant 55 miles, provides another market for this product.

Estimate of Production Costs

Mining, tramming, crushing	@ \$1.30 per ton
Leaching, labor, supplies, water, power	@ .75 " "
Overhead, accounting, etc.	@ .20 " "
Direct Production Costs	\$2.25 per ton amended cost estimate

Property Survey and Appraisal

Mines opened during 1880 to 1910.
Pilot leaching plant was built late 1944.
Buildings consist of 5 room mine cabin and small shop, situated about 1/2 mile west of adit tunnel.
Power at present - none used. Will be small, semi Diesel unit.
Present value and cost of surface improvements \$3,500.00.
Cost of proposed copper leaching plant 10,000.00.
Mine developments now usable estimated at 50,000.00.

Operations and Earnings

1. The proposed unit of 50 tons daily is believed to be adequate to recover 75% to 85% of the copper in the ores to be leached.
2. There are visible and available ores within the tunnel and it is believed that Mr. H. R. Scott is sincere and with sufficient ability to manage the work, and by his engaging an elderly man with sufficient experience in making copper determination and titrations the project can be operated efficiently.
3. The above estimate for completed 50-ton plant seems fair at \$13,500.00.
4. As only from 5 to 10 elderly miners and millmen are required, these are available in Globe or Winkelman - men rejects of the large copper producers.

NAME OF MINE: RATTLER

COUNTY: PINAL

DISTRICT:

METALS: CU

OPERATOR AND ADDRESS:

MINE STATUS

DATE:

5/1/44

H.R.Scott, Box ~~2893~~ 2893, Globe

DATE:

5/1/44

Mining

5/16/44

Shipping per Mact.

1/45

Shipping occasionally

10/45

Idle

4/46

Developing

Report on file by McFarlane

H. R. Scott, Box 872, Globe, Arizona, D. P. McGarvin, c/o Arizona Tours, First Street and Monroe, Phoenix, Arizona, and John A. Devine, Box 872, Globe, are reported to be installing a 30-ton leaching plant at the Troy Copper Company's Rattler group of claims located about one mile east of Troy, Arizona, in the Dripping Springs district of Pinal County. The mine is said to carry a quantity of copper sulphate which is amenable to treatment by the leaching process. Scott is manager of the project which is said to be financed by eastern interests. Production is expected to be started at the property in about 60 days. MINING JOURNAL - 10-15-44

RATTLER GROUP

Cu

Pinal

11 - 3

T 3 S, R 1 E

H. R. Scott, Box 2893, Globe

'44

DEPARTMENT OF MINERAL RESOURCES
STATE OF ARIZONA
FIELD ENGINEERS REPORT

Mine Rattler Group of Copper Claims and the
 Pilot Copper Leaching Plant Thereon
District Riverside

Date February 15, 1945

Engineer A. Macfarlane

Subject: Report

On April 17, 1944 the field engineer for the Eastern area of the State examined this property and his report of that date is now part of this second examination and report.

The second examination was confined to a study of the lower tunnel workings and the 8-ton Pilot copper leaching plant recently installed just beneath the outer portal of the main or adit tunnel.

Mine Workings: The attached maps show over 60' of adit driven southward from portal through what appears to be a diabase, barren of mineralization, until the limestone here is reached. About one hundred feet within tunnel from portal a strata conforming to the limestone bedding planes is visible as a body of low grade copper, having a dip of 10° towards the south and the lateral continuance being easterly and westerly from the stopes and tunnel sides.

There are approximately 900 linear feet of exploration under ground workings now open for inspection, and probably as much more at present caved and not now accessible.

Measurable Ore Tonnage: I estimate the visible workings of the adit tunnel to now contain upwards of 10,000 tons, and in order to obtain a practical estimate of the copper content, I dug 4 holes into the dump made from former stoping of the tunnel copper body.

This dump contains about 2,000 tons mostly copper in carbonate and sulfate form and the assay gave a copper content of 1.29%. As this dump was made by a long past operation, it is likely some higher grade was sorted out and shipped to one of the smelters then.

The presumption of 1.3% copper seems fair as a basis for further estimates covering the mining and leaching of the ores from the area and copper strata of the adit tunnel.

The Session ore body is stratigraphically about 250' higher than the adit tunnel and seems to have a much steeper dip again towards the south. These separate copper stratas are connected to the lower or adit level by a winze from floor of upper tunnel and by the Session shaft which has a depth vertically below collar of nearly 300 feet, thence drifts and crosscuts connecting with the adit tunnel.

In all many thousand tons of leachable copper ores are opened by the large amount of mine developments made during the latter part of the 19th Century on this property, all mineable through the lower or adit tunnel.

Mine Costs: No further mine developments need be considered now, the old workings containing both in the stope faces and gobbing an ample ore reserve for one or more years.

February 15, 1945

Mine tracking is required and a few ore chutes. The limestone hanging wall is firm and only occasional pillars and stulls will be required to safely sustain same. The copper ore stratas varies in thickness from 3' to 6'.

On basis of mining, tramming and crushing 100 tons in the day shift of 8 hours,	
cost @	\$1.10 per ton
Leaching, labor, supplies, water, power	.70
Overhead and emergency	.20
All direct operating expense	\$2.00 per ton

Estimate of Outcome: Assume a recovery of 25 pounds copper per ton of ore; copper @ 17 cents less marketing cost of 3 cents, we have 25# at 14 cents net, or a gross per ton of \$3.50, or if 20# recovered @ 14 cents, a gross per ton of \$2.80. In either assumption, a substantial margin of \$.80 to \$1.50 per ton may be earned in order to repay the plant investment.

Pilot Leaching Plant: During the past 4 months the owner of the Rattler Mine, Mr. H. R. Scott, has purchased and installed a small leaching unit consisting of 3 redwood tanks. Capacity of around 8 tons ore and solution per tank. Also 2 wooden precipitating boxes made into compartments with perforated shelf 4" above bottom, holding about 1,000 pounds scrap iron. The pregnant solution drawn from near bottom of the leaching tanks at 36 to 48 hour periods, circulates through the scrap and discharges into a small sump, thence pumped back into the leaching tanks. A 2% sodium bi-sulfate solution is used in the leaching tanks to liberate the copper from the crude ore.

The ore as trammed into the leaching tanks heretofore has not been crushed and ranges in size from fines to pieces barely passing a 4" ring. No accurate sampling and assaying has been made on the heads and tails other than a dump sample assaying 1.3% cu, probably a close assay of the 30 tons crude ore fed into the leaching tanks.

At request of the writer the sludge or precipitating boxes were cleaned up, the cement copper dried and weighed. A sample of this first lot assayed 52% cu and the total dried weight of 1,200 pounds figures 624 pounds metallic copper from 30 tons crude ore, equals 21 pounds per ton recovered and is 81% of the copper originally in the ore.

Recommendations: It is of paramount importance to the smaller copper mines of Arizona and elsewhere that a leaching unit of 50 to 100 tons daily capacity shall be designed and made workable through tanks. This method of copper leaching may be done with a small and limited water supply, the consumption of same may not exceed 1/2 ton water per ton crude ore, and by proper sampling and analysis the proper quantity of activating acid will be determined and the time of leaching required to convert the copper into solution.

The above practical on the ground experiment No. 1 indicates that better copper recovery can be made by crushing all ore through a 1/2" to 1" size and again the activating acid, whether the sodium bi-sulfate, or sulfuric acid, shall prove most efficient, are problems yet to be determined.

All the large copper operators of southern Arizona commonly known use sulfuric acid as the principal activating agent. The application of sodium bi-sulfate may not be affected by the very low percent of free lime in an altered limestone, and might well

be tried out further, and in the event the sulfuric acid method proves cheaper and more efficient it will cost very little to change the solvent.

Plant of Commercial Size: It is now desirable to instal a 50-ton "per 8 hour unit" and the following estimates may approximate a workable plant and costs thereof erected on a mill site underneath the adit tunnel.

Required 2 rock and cement leaching tanks 20' x 25' x 5' each. Each tank will have a capacity of 100 tons ore and solution.

Cost built on the mill site \$1,500 each	\$3,000.00
Ore crusher designed to crush fine installed above tanks, together with bridging and power	2,000.00
Additional scrap iron precipitating boxes	500.00
Hillside grading and retaining walls	500.00
Iron cleaning furnace and burner complete	500.00
Titration and cleaning up floor and building	500.00
2 acid proof solution pumps and power	700.00
1 drying tray and furnace burner	300.00
1 slusher with adequate guide and tail	1,000.00
(pulleys, cable, etc. installed to unload leaching tanks)	
Contingent needs and reserve	<u>2,000.00</u>

Total plant erected and maintained in operation for 2 months	\$8,000.00
--	------------

Mine Preparation: As the gob is cleaned out along the tunnel floor, 8# to 12# tee rail used, with 24" x 4" x 6" ties will be laid from crusher to mine headings, in all about 1,000 linear feet of mine trackage. Most of the mine gob left by a former operation is copper ore and will be trammed direct to the leaching tanks. Trackage will cost \$1,000.00.

On completing the mine cleaning a 3 or 4 drill capacity air compressor with all accessories, piping, etc. will be required, a used plant may be secured costing	2,000.00
Mine timbers, shop and reserve for 60 days operation	<u>2,000.00</u>


Total mine reconditioning	\$5,000.00
---------------------------	------------

Road: A good state road distant 5 miles from the Rattler leads to the Hayden smelter situated about 25 miles south. From the point on state highway nearest a piece of new road should be built to pass the Cockerham gold mine, thence westerly some 2 miles to the Rattler pilot mill. Application to the access road department will be made for this new piece of road. The property is now connected to this highway by a rough but passable road leading from the Rattler mine via Troy to the Winkelman-Globe road. Water is pumped from over a ridge distant 1/2 mile, in sufficient volume to provide the anticipated leaching plant.

Conclusion: Due to the extensive mine development work of many years ago in a former effort to ship higher grade copper to smelters. This work has made available and visible through the adit tunnel a large tonnage of low grade copper carbonate ores.

Within the lower workings where dampness contacts the ore chemical change to sulfate is very noticeable, suggesting adaptability to leaching.

It is a step in the right direction to develop a small sized copper leaching plant at this Rattler mine. Only such plants may treat copper carbonate ores of less than 2.5% to 3% copper. There is a very large tonnage of such ores in Arizona, which must be depended upon to provide work and the useful copper product as a post war measure.



PLAN OF RATTLER MINE WORKINGS
 Former Survey: Owner H.R. Scott, Globe, Ariz.
 Examination of the Field Engr. Feb. 2 1945
 Department of Mineral Resources
 Scale 1" To 30' Office 304 Home Builders, Phoenix
 Location, Riverside Mining District, Pinal County, Arizona
 Dripping Springs Range

LONGITUDINAL PROJECTION Looking Northwest

RATTLER MINE WORKINGS

Riverside Mining District, Dripping Spgs. Range


Pinal County, Arizona. Approx. Scale 1" To 30'

Owner H. R. Scott, Globe, Arizona


Examination Made By Globe Field Office

Department of Mineral Resources


February - 3 - 1945




Plan of Proposed
Copper Leaching Plant
Rattler Mine Dripping Spgs.
Pinal County, Arizona
For H.R. Scott, Globe "
Scale 1/4" to 20' Feb. 30th 1904
From Field Notes, Engr
Dept. Mineral Resources



canyon





Tailing
Disposal

PROFILE

Adv. T.M.

Plan of Proposed
Copper Leaching Plant
Rattler Mine Dripping Spgs.
Pinal County, Arizona
For H.R. Scott, Globe "
Scale 1/2" to 20' Feb. 20th 1945
From Field Notes, Engg
Dept. Mineral Resources

600 m. east side north.

One of these faults is Precambrian, as shown by the fact that it has a Precambrian diabase dike along part of its length. The greatest tectonic activity, however, must have occurred after the Paleozoic Era because of the major involvement of Paleozoic rocks, and faulting probably continued into middle or late Tertiary time as indicated by the substantial horizontal offset of the early Tertiary rhyodacite dike, mentioned above.

The southwest margin of the Dripping Spring Valley is bounded by a major normal fault (see section *D - D'*), north side down, probably of Miocene age, forming the basin that was filled with Tertiary alluvial and lakebed deposits. A churn drill hole in Dripping Spring Wash, 180 m west of the quadrangle boundary, penetrated 448 m of Tertiary conglomerate and did not reach the bottom of the basin even though the older rocks of the basement crop out 300 m to the southwest. The Tertiary alluvial basin in the northeast corner of the quadrangle must be underlain by a concealed west-trending normal fault, north side down, because of the position and attitude of Paleozoic sediments to the north (see section *E - E'*) in an area of Precambrian rocks.

ECONOMIC GEOLOGY

Mineral exploration in the El Capitan Mountain quadrangle started about 1870 and has continued intermittently to the present. Mining activity, especially in the early days, focused on silver and gold, which are associated with copper, lead, and zinc sulfides in fissure veins and replacements of carbonate sedimentary rocks. The deposits for the most part occur along or near major north- to northeast-trending high-angle faults in areas of moderate to intense deformation. There has been a small production from several mines. Between 1915 and 1926 one mine produced some vanadium associated with lead and silver. In the 1950's intensive uranium exploration in the western United States identified two deposits in this quadrangle, one of which produced a small amount. The mineral resource potential of this quadrangle is probably favorable for base metals, silver, and gold. The chances are quite good for the discovery of small- to moderate-sized deposits of these metals. Descriptions of individual deposits are given below.


The Cowboy gold mine (NE¼ sec. 31, T. 3 S., R. 15 E.) consists of two shallow shafts and several pits and adits along a shear zone striking N. 80° W. and dipping 80° S. occupied by a rhyodacite porphyry dike, and a vertical crosscutting shear zone striking N. 25° E. The deposit occurs in Mescal Limestone and the rhyodacite dike, and along the contact of a large diabase intrusion to the south. Vein material on the dumps and in the workings consists of vuggy quartz stained with iron and manganese oxides, and variable amounts of limonite, jarosite, specularite, malachite, chrysocolla, ankerite, wulfenite, and hemimorphite. Ransome (1923, p. 23), in discussing the mine, stated: "According to Mr. C. W. McGraw, the owner and operator, much of the gold is in coarse wire form and occurs erratically in pockets, particularly where the vein is in limestone. A sample of concentrate sent by Mr. McGraw to the Geological Survey shows the presence of a lead vanadate, probably descloizite. The total output of the Cowboy mine to the end of 1918 is estimated at about \$25,000 in gold, with a little silver and lead."

A gold placer was worked in the early 1930's in the wash that runs east from the Cowboy mine into Dripping Spring Wash (J. T. Eastlick, Chief Resident Geologist, Inspiration Consolidated Copper Co., written commun., 1976). A small amount of gold was recovered from a 1-m streak at the bottom of the stream channel, 9 - 12 m below the surface, near where the wash enters Dripping Spring Wash. The gravel contained \$2.00 in gold per cubic yard.

The Dripping Spring mine (sec. 25, T. 3 S., R. 14 E., and NW¼ sec. 30, T. 3 S., R. 15 E.), called the Columbia mine of the Dripping Springs Mines Corporation by Ross (1925, p. 67 - 68) and, according to him, consisting of about 20 claims, was located by J. W. Read about 1901; it was taken over by the Dripping Springs Copper Co. about 1915, which then explored with a number of adits, shafts, stopes, and drill holes through the early 1920's. The workings are shallow in the oxidized zone, and mineralization consists mostly of copper oxides, carbonates, and silicates (mostly malachite and chrysocolla) together with some zinc (reported by Ross but minerals not identified). Limonite, specularite, quartz, and calcite are also present in the copper-bearing veins, which occur mainly in east-striking, steeply south-dipping faults and shear zones where they cross the contact between the Abrigo Formation and the Martin Limestone.

The Amax Exploration Company claims (sec. 30, T. 3 S., R. 15 E., and SE¼ sec. 25, T. 3 S., R. 14 E.) were explored by that company in 1972 with several diamond drill holes. In this area there are a number of older prospects, mainly adits near the contact between the flat-lying Martin Limestone and Abrigo Formation. Mineralization in these workings consists of oxidized copper minerals (chrysocolla and malachite) plus limonite, quartz, and calcite in fissures and faults, mostly dipping steeply.

The Vanadium mine (N½ sec. 32, T. 3 S., R. 15 E.), called C and B group by Ross (1925, p. 69), consists of a vertical shaft several hundred feet deep, two adits, and an open stope. Stoping was along two 1-m shear zones; the main one strikes N. 50° E. and dips 45° S., and the other strikes north-south and dips 50° E. Mineralization is mainly along contacts of lenses of Mescal Limestone included in Precambrian diabase. Exposed mineralized rock is in the oxidized zone and contains cerussite, vanadinite, wulfenite, quartz, and calcite. Ross (1925, p. 69) stated that some ore was mined and shipped during World War I. The property was examined by C. L. Beckwith of the Inspiration Consolidated Copper Company in 1926 (J. T. Eastlick, written commun., 1976). Beckwith stated that one small car of ore was shipped to the smelter that year. Beckwith also reported that the ore minerals cerussite and vanadinite occur in bunches through the vein material, which is altered limestone. He cut four channel samples in mineralized rock; three averaged 0.33 oz silver per ton, 1.92 percent vanadium, and 6.2 percent lead, and the fourth contained 0.17 percent vanadium and traces of silver and lead.



olidated Copper Company mapped and sampled the workings in 1968 (J. T. commun., 1976). Their investigation indicates that an adit follows the r zone for 120 m. Four channel samples 1 - 2 m long across the vein aver- per ton and 0.92 percent copper. One sample assayed 2.25 percent lead. e mine (NW¼ sec. 3, T. 3 S., R. 15 E.) consists of three adits, a shallow n the siltstone member of the Dripping Spring Quartzite, which is crosscut . Vein fragments on the dumps are quartz and altered siltstone with abun- ns, and limonite in part as pseudomorphs after pyrite. No production has

ek mine (W½ sec. 34, T. 2 S., R. 15 E.) consists of several adits and pits st-striking fault zone dipping 60° - 70° S. that separates the siltstone mem- ing Spring Quartzite on the north from diabase on the south. Mineraliza- ine-grained quartzite, consists of sphalerite, galena, azurite, malachite, and stlick (written commun., 1976) reported that a grab sample of vein material ayed 4.18 oz silver per ton, trace of gold, 2.5 percent lead, 2.3 percent zinc, t copper. There is no recorded production.


ine (NW¼ sec. 29, T. 2 S., R. 15 E.) was active in the 1880's, according to p. 23). There are two shafts, an adit, and two narrow open stopes along N. 30° - 50° E., dipping vertical to 75° W., and cutting across diabase, Quartzite, and Pioneer Shale. J. S. Conpal (written commun., 1927) re- ing started in the early 1870's; production through 1927 amounted to about nly silver, averaging about 25 oz per ton, from oxidized ores mined to 90 m. Below this depth unoxidized sulfides, mostly galena and sphalerite, ult to treat economically by methods available at that time. Conpal stated ree major, roughly parallel, northeast-striking veins plus several minor ones, ntaining silver, lead, zinc, and copper sulfides plus minor gold. Vein frag- mps contain sphalerite, galena, quartz, barite, and calcite. There was also vity in the area in the late 1940's, and several holes were diamond drilled. y uranium deposit (secs. 31, 32, T. 2 S., R. 15 E.), described by Granger , p. 472; 1969, p. 68 - 70, figs. 23, 36, 38), occurs in a chlorite-filled shear thick parallel to bedding in locally bleached dark-gray siltstone of the upper Dripping Spring Quartzite. Metatorbernite associated with limonite is abun- radioactive parts of the deposit. It is estimated that 2,000 - 4,000 tons g 0.1 - 0.2 percent U_3O_8 were mined between 1954 and 1957. Granger , p. 79) stated that the uranium in the unoxidized parts of these deposits Spring Quartzite "occurs principally in several varieties of uraninite but on nontronite, chlorite, and, more rarely, graphite." They believed that the rived from diabase that intruded the Dripping Spring Quartzite about 1,050 b (1969, p. 76). The uraninite was concentrated in the carbon-rich siltstone ation in deuteric solutions that emanated from the diabase.

um prospect (SW¼ sec. 2, T. 3 S., R. 15 E.), described by Granger and Raup occurs near a discordant diabase body in dark-gray shale and siltstone of the of the Dripping Spring Quartzite. Radioactivity is abnormally high for sev- et near the base of the upper member and has been explored by several pits ill holes. Metatorbernite, malachite, limonite, pyrite, and gypsum occur and bedding planes. No minable ore body has been found, however.



Base from U.S. Geological Survey, 1964
10,000-foot grid based on Arizona coordinate
system, east zone
1000-meter Universal Transverse Mercator
grid ticks, zone 12, shown in blue

4



GEOLOGIC QUADRANGLE MAP EL CAPITAN MOUNTAIN, ARIZONA GQ - 1442

TEAPOT MOUNTAIN PORPHYRY (Paleocene) - A discontinuous east-west-trending gray to light-brownish-gray quartz latite dike in the northwestern part of the quadrangle (SW¼ sec. 25, T. 2 S., R. 14 E.) in Precambrian Y diabase contains scattered 5- to 25-mm phenocrysts of pink potassium feldspar and 1 - 8 mm phenocrysts of plagioclase, quartz, and green biotite in a fine-grained to aphanitic groundmass of the same minerals plus magnetite-ilmenite, hornblende, apatite, calcite, and zircon. The feldspars are moderately to intensely altered to sericite, calcite, montmorillonite, and kaolinite, and the biotite and hornblende to chlorite, epidote, calcite, and hematite. S. C. Creasey (oral commun., 1970) has obtained a potassium-argon date of 63 ± 2 m.y. (Paleocene) for potassium feldspar from the porphyry in the Teapot Mountain quadrangle to the west.

RHYODACITE PORPHYRY - Vertical dikes of light-brownish to medium-gray rhyodacite with prominent but locally sparse quartz phenocrysts. Phenocrysts, 1 - 6 mm in diameter, making up approximately half of the rock, are in decreasing order of abundance plagioclase (An₂₅ - 40), partly glomeroporphyritic, biotite, hornblende, and quartz. The groundmass, anhedral-granular in 0.01 - 0.4 mm grains, consists of potassium feldspar and magnetite-ilmenite plus the other minerals listed above and accessory apatite, zircon, sphene, and allanite. The plagioclase is partly altered to sericite, montmorillonite, kaolinite, calcite, and epidote, and the biotite and hornblende to chlorite and epidote. A typical mode (volume percent) is: andesine 54, quartz 20, potassium feldspar 19, biotite 3.5, hornblende 3, and magnetite-ilmenite 0.5. These distinctive dikes are also common in nearby quadrangles to the west and south. In the Grayback quadrangle (Cornwall and Krieger, 1975b) dikes of this type cut the Tea Cup Granite diorite of Paleocene age and are thus Tertiary, probably Paleocene, in age.

MELANOCRATIC RHYODACITE PORPHYRY - Several vertical dikes in the southwest corner of the quadrangle cut the Troy Quartzite and Ratler Grandodiorite. Medium- to dark-gray and brownish-gray fine-grained porphyry with phenocrysts, making up roughly one-third of the rock, of andesine, hornblende, and quartz, as much as 1 cm in longest dimension. Groundmass consists of the same minerals plus magnetite-ilmenite, potassium feldspar, and accessory apatite, sphene, and zircon. Plagioclase (andesine) is rather intensely altered to montmorillonite, kaolinite, sericite, calcite, and epidote; hornblende and biotite are partly altered to chlorite, epidote, calcite, and sphene. These distinctive dikes are also common in nearby quadrangles to the west and south.

RHYODACITE PORPHYRY - Generally forms dikes but locally occurs as sill-like bodies in the Pioneer Formation. Light- to medium-gray and brownish-gray fine-grained porphyry with 1 - 3 mm phenocrysts of andesine, hornblende, biotite, and locally quartz in a groundmass of the same minerals plus potassium feldspar, magnetite-ilmenite, and minor apatite and calcite. The plagioclase (andesine) is partly to entirely altered to montmorillonite, kaolinite, sericite, and minor epidote; the biotite and hornblende are partly altered to chlorite and epidote.

RATLER GRANDODIORITE (Upper Cretaceous) - A pear-shaped composite

Y₁

TROY QUARTZITE (Precambrian Y) - 125 - 325 m of quartzite, sandstone, and minor conglomerate; probably correlates with the Chediski Sandstone Member as suggested by Shride (1967, p. 44 - 46). Sandstone and quartzite range in color from grayish pink and white to light brown and grayish orange and consist of medium to very coarse subangular to rounded quartz grains with variable amounts of feldspar, limonite, and dark accessory minerals in a matrix of finer quartz and clay. Thin layers and lenses of poorly to well-sorted pebbles and granules are interbedded in the sandstone and quartzite. Beds are laminated to thin bedded and arranged in thin to thick composite tabular to lenticular sets, locally crossbedded or convolute. Conglomerate 1 - 4 m thick occurs at the base of the formation with angular to rounded pebbles and cobbles of white quartz, various sandstones, quartzites, and cherts, basalt, Mescal Limestone, and Pinal Schist. Conglomerate beds 0.3 - 2 m thick are scattered through the formation, particularly toward the top. The predominant quartzites of the Troy form bold cliffs with intervening slopes underlain by more friable sandstones.

Y_b

APACHE GROUP (Precambrian Y): Includes in ascending order the Pioneer

Formation, Dipping Spring Quartzite, Mescal Limestone, and basalt

BASALT - 20 - 65 m of mostly porphyritic dark-gray to grayish-red or brown basalt, one or more flows, vesicular tops with amygdulæ of calcite,

Y_m


chlorite, potassium feldspar, and epidote. Plagioclase phenocrysts, partly glomeroporphyritic, are 2 - 8 mm long. Groundmass is fine grained (0.05 - 0.3 mm) and consists of plagioclase (An₅₀), pyroxene, olivine, and magnetite-ilmenite intensely altered to chlorite, sericite, calcite, epidote, quartz, hematite, limonite, bowlingite, kaolinite, and some potassium feldspar.

MESCAL LIMESTONE - 80 - 120 m of light-pink, brown, and brownish-gray dolomite, partly calcareous, thinly laminated to thin bedded, very fine grained to coarse grained. Some beds contain abundant black, white, and pink chert lenses and nodules. Stromatolitic algal beds occur above the middle of the formation, and layers of sandstone and quartzite near the middle in some areas. The formation has been metamorphosed by diabase to marble and calc-silicates in many places, and certain beds have been replaced locally by magnetite, tremolite, and serpentine.

DRIPPING SPRING QUARTZITE - 245 m of siltstone, sandstone, quartzite, and conglomerate; includes the following members in descending order:

Siltstone member (145 - 170 m). - Siltstone interbedded with shale and fine-grained arkosic sandstone, laminated to thin bedded with low-angle cross-bedding; light to dark gray, olive green, dusky brown, and grayish

Y_{ds}



3669

Hagen Ranch

3668

Radio Group

GILA CO
PINAL CO

Spring

Spring

3666

Cave

3655000m N.

110° 52' 30"

110° 07' 30"

(HAYDEN)
3850 III SE

● INTERIOR- GEOLOGICAL SURVEY, RESTON, VIRGINIA - 1982

510 280 000 FEET (EAST) 511000m E

ROAD CLASSIFICATION


Heavy-duty Light-duty

SCALE 1:24 000

(KEARNY)
3850 III SW

0 3000 4000 5000 6000 7000 FEET

1 MILE





Mapped, edited, and published by the Geological Survey

Control by USGS and USC&GS

Topography by photogrammetric methods from aerial photographs taken 1962. Field checked 1964

Polyconic projection. 1927 North American datum
10,000-foot grid based on Arizona coordinate system, east zone
1000-meter Universal Transverse Mercator grid ticks,
zone 12, shown in blue



Troy District Mining Area
Dripping Springs Dist.

Ray Folio - #217, 1923

ECONOMIC GEOLOGY.

GENERAL CHARACTER OF RESOURCES.

ic monzonite and dioritic
usion of these rocks was
l by faulting and was the

usion, at a time which can
t information but which is
ding with the earlier or
region, characterized by a
ently dry land and under-
ce was probably less rugged,
imate and erosional activity
different from those of the
cumulation of the Whitetail
etritus was washed down the
open valleys and gulches. It
and accumulation of waste
the Ray and Miami copper
place.

its hollows partly filled with
poured, probably in early or
e flow of dacite. The greater
appears to have been covered
a vent or vents as yet undis-
mination in places shows rude
ite in some localities is under-
probable that the region con-
r, probably in consequence of
by faulting before and during

acite eruption, and possibly as
eat faulting to which are chiefly
less directly the topography of
his faulting has been described
e country, especially the Drip-
ered to an extent very imper-
number of small fault blocks
f the Ray quadrangle.

factory evidence for connecting
epochs the events which took
Carboniferous period, the post-
trarily considered as ending the
nature of this and other post-
this region should not be for-
rably modified when the present
l and extended by the study of a
recognized appear to be distinct
ul history. They may not, how-
et places in the larger volume of

1.
un by a vigorous erosion of the
resulting from the superposition of
upon an earlier structure that was
quantities of coarse, rocky detritus
es and deposited as the Gila con-
at least, of structural origin. It
that the larger conglomerate-filled
riginal depression to faulting.

a conglomerate indicates that the
early Quaternary were not very
prevailing aridity and dominance of
over rock decay were prominent
ion apparently occurred in violent

one eruption of basalt during the

The rough and generally rocky or stony surface of the Ray quadrangle, with its scanty desert vegetation, although utilisable in part for the grazing of cattle and the browsing of goats, offers little inducement to human occupancy or industry. The economic development of the area depends almost entirely upon its mineral resources, and of these copper is supreme. The Ray copper district, which is in the northwestern part of the quadrangle and extends in small part into the adjoining Florence quadrangle, on the west, contains one of the largest deposits of copper ore in Arizona. This ore is being mined on a large scale, and most of it is reduced to metal within the Ray quadrangle. Some copper ore is produced also near Troy and at and near the London-Arizona mine, in the Dripping Spring Range; and on the south side of the Gila, near Kelvin.

The area also contains ores that yield gold, silver, lead, zinc, and vanadium.

HISTORY OF MINING.

Apparently the first mines to be worked in the Ray quadrangle were those which yielded silver-gold ore near Pioneer, in the northeast corner of the quadrangle, and perhaps the Ripsey mine, in the southwest corner. The argentiferous ore of the El Capitan mine, south of Old Baldy, in the Mescal Range, was probably also worked in this early period, during the eighties. Very little definite information is now obtainable concerning these pioneer efforts in what was then a wild and remote region. The Republic mine, at Pioneer, had a mill and apparently was operated successfully for a number of years.

Some mining was done also about 1880 on Mineral Creek, near the site of the present town of Ray. At that time the Mineral Creek Mining Co. built a 5-stamp mill and did some work, presumably on the Mineral Creek claim, north of Copper Gulch. The subsequent history of mining development on the the ground now owned by the Ray Consolidated Copper Co., is given elsewhere.⁶¹

The Arizona Hercules Copper Co., whose ore-bearing ground is almost inclosed by the claims of the Ray Consolidated Copper Co. and contains the eastward extension of the Ray ore body, began exploration by drilling a little later than the Ray company. The existence of ore was soon ascertained, but it was not until 1916 that active steps were taken to mine it. In that year two shafts were sunk and levels were run preparatory to extensive mining. The shafts were completely equipped with first-class machinery and a coarse-crushing plant was built at the mine, railway connections were made, and at the settlement of Belgravia, 6 miles from the mine, near Kelvin, a 2,700-kilowatt power house and a 1,500-ton concentration mill were constructed in 1917 and 1918. Production of copper on a large scale began in 1918.

Active mining development at Troy began about the year 1900 by the Troy Copper Co., a Boston corporation organized under the laws of Maine and capitalized at \$1,000,000. About the same time the Manhattan Copper Co., of New York, capitalized at \$1,500,000, entered the field. The Troy company had about 30 claims lying chiefly in the western part of the Troy district and including the '91, Buckeye, Climax, and Alice. The Manhattan had about 15 claims, including the Rattler. Most of the development work appears to have been done between 1901 and 1903. In 1902 the two companies consolidated as the Troy-Manhattan Copper Co., capitalized under the laws of Maine at \$3,000,000. A 60-ton smelter, at River-

tion has been active over the mountains and dissecting the Gila

The total production of metals in the Ray quadrangle can not be given, as no records of output prior to the extensive mining near Ray are available. The older yield, however, can not have been large.

ters probably carrying
d that these particular
fy this commonly held
s appear to have been
although the quantity
rge in the aggregate, it
of the metallized rock

URE.

between primary metal-
said that such metalli-
vicinity of the contact
at this is true only in a
found more than 1,000
s of considerable size.
s of the contact region
In the Ray district, for
porphyry on Granite
metamorphism, are not
ami district it is only
h lobe of the Schultze
disseminated pyrite and

e ascent of metallizing
re doubtless complex.
lity of the rocks affected
olutions.

f the protore, was due
ssuring. This fissuring
f larger fissures along

red also by irregularity
esence of little tongues
nto the schist. The act
isturbance of the schist,
d dikes, by introducing
bably made for further
communicating channels
al and the zone of sul-
given to this suggestion
ular protrusions of por-
ground. This is partic-
ill, where the ore body
thickness, however, it
ue more to enrichment

ns available at any place
ed to a large extent also
of magma from which
ntirely beyond our ken,
ent at least, unknown.

COUNTRY ROCK.

or significant difference
protore as regards tenor
not more abundant, is in
of the porphyry protore
ite also seems to be more
At Ray the mineralized
re pyrite and chalcopy-
opper than the average
1.) This is not surpris-
diabase contains a much

on general grounds rather than from any definite evidence, is supposed to have taken place at the end of the Mesozoic era. It appears reasonable to regard the intrusion of the granitic porphyries as an early Tertiary event, but it must be admitted that this is little more than conjecture. The deposition of the protore certainly took place after the laying down of the Tornado (Mississippian and Pennsylvanian) limestone and before the eruption of the dacite.

TROY DISTRICT.

The intrusion of the granodiorite of the Troy Basin was followed by pronounced contact metamorphism and by considerable metallization. The granodiorite is closely related to the Schultze granite and the Granite Mountain and Teapot Mountain porphyries and was intruded at about the same time as those rocks. It is associated with many dikes and is surrounded by much fissured rocks, including limestone, which differ in no essential respect from rocks elsewhere ore-bearing in the same general region. In short, Troy would seem to be a decidedly favorable place for ore deposition. Yet those who have acted on this apparently reasonable supposition have thus far been disappointed. There has been extensive prospecting, and some ore has been found, but the returns have not equaled the outlay.

The principal development has been on the Rattler claim, 1 mile in a direction a little south of east of Troy, and on the '91, Buckeye, and Alice claims, one-half, three-fourths, and 1½ miles southwest, respectively, of the now practically abandoned settlement. When the visits were made in 1910 and 1914 the underground workings were only in small part accessible.

The workings of the Rattler mine comprise the Sisson shaft, which is an incline of 65° to the south and 300 feet in depth, connected with three levels, the first of which is an adit. The levels run nearly east-northeast and west-southwest and open a section of ground about 700 feet long. A second shaft, east of the Sisson, extends only about 50 feet below the adit level. There is no vein, the ore occurring as bunches and lenses that follow more or less closely the bedding of the Mescal limestone in which it occurs. The limestone at the Rattler mine is an inclusion in diabase, the original igneous contact being modified in some places by faulting. Granodiorite was not seen in the workings, but as shown on the geologic map it is not far away and has effected considerable contact metamorphism in the diabase, which sparkles with secondary biotite. The ore zone dips 20° S. 15° W. and from a point in the main tunnel about 100 feet in has been followed in an inclined winze for about 95 feet to a point where it appears to be cut off by a rather obscure fault. There is apparently no large quantity of ore available. The Sisson shaft appears to be entirely in diabase below the first level.

The ore of the Rattler mine is chiefly a dark fine-grained aggregate of magnetite and chalcopyrite with varying quantities of silicate minerals derived by metamorphism from the inclosing limestone. With increasing proportions of these silicates the ore grades into the metamorphosed limestone. Analyses of the ore recorded in the books of the Troy Arizona Copper Co. and its predecessors show from 3 to 3.7 per cent of copper, a maximum of 0.04 ounce of gold and 0.7 ounce of silver per ton, from 27 to 30 per cent of silica, about the same proportion of iron, about 1 per cent of calcium oxide, and 20 per cent of magnesium oxide.

Very little could be seen of the '91 mine, as the shaft had caved in. It apparently is about 150 feet deep, and the maps show three short levels. The little copper ore that was found in this mine appears to have occurred, as at the Rattler,

at determined, the ascent of metallizing particular place were doubtless complex. have been permeability of the rocks affected by of the active solutions.

own by a study of the protore, was due minute irregular fissuring. This fissuring the formation of larger fissures along place.

believed, was favored also by irregularity and by the presence of little tongues extending out into the schist. The act have caused some disturbance of the schist, such tongues and dikes, by introducing the rock mass, probably made for further extent provided communicating channels of igneous material and the zone of sulfide probability is given to this suggestion of dikes and irregular protrusions of porphyry in the metallized ground. This is particularly Humboldt Hill, where the ore body thickness. That thickness, however, it is, is probably due more to enrichment than to the protore.

Generalizing solutions available at any place probably depended to a large extent also to a deep-seated mass of magma from which the ore, however, is entirely beyond our ken, and, for the present at least, unknown.

OF VARIATIONS IN COUNTRY ROCK.

There is no regular or significant difference between the protore and porphyry protore as regards tenor of pyrite, however, if not more abundant, is in the protore. Molybdenite also seems to be more abundant in the protore. At Ray the mineralized protore appears to carry more pyrite and chalcopyrite than in schist. At Ray the mineralized protore is higher in copper than the average protore. (See p. 21.) This is not surprising, considering that the diabase contains a much larger amount of copper originally present as oxide and silicate and also, as has been shown in a previous section, that the protore was originally a little copper. Practically no ore had been mined in diabase in the field work for this folio was completed, and there is no reason why, under suitable conditions, the mineralized diabase should not have been mined. Consequently it would not be surprising if ore was found in that rock east of Mineral Creek.

SURFACE EXISTENT AT THE TIME OF DEPOSITION.

It is at the time the protore was deposited at the surface that the protore lay above the present surface, and it is probable that the thickness was several times the figure shown. The crystallinity of the granite porphyry and the metamorphism that accompanied or followed the protore is indicative of the solidification of the protore by a thick cover. In the Miami district the protore, at the present day, is in places fully exposed to the ore.

GEOLOGIC AGE.

The protore probably followed closely the granite porphyry, but no facts are known that place this event definitely in geologic time. The protore is younger than the diabase, whose intrusion,

The principal development has been on the Rattler claim, 1 mile in a direction a little south of east of Troy, and on the '91, Buckeye, and Alice claims, one-half, three-fourths, and 1½ miles southwest, respectively, of the now practically abandoned settlement. When the visits were made in 1910 and 1914 the underground workings were only in small part accessible.

The workings of the Rattler mine comprise the Sisson shaft, which is an incline of 65° to the south and 300 feet in depth, connected with three levels, the first of which is an adit. The levels run nearly east-northeast and west-southwest and open a section of ground about 700 feet long. A second shaft, east of the Sisson, extends only about 50 feet below the adit level. There is no vein, the ore occurring as bunches and lenses that follow more or less closely the bedding of the Mescal limestone in which it occurs. The limestone at the Rattler mine is an inclusion in diabase, the original igneous contact being modified in some places by faulting. Granodiorite was not seen in the workings, but as shown on the geologic map it is not far away and has effected considerable contact metamorphism in the diabase, which sparkles with secondary biotite. The ore zone dips 20° S. 15° W. and from a point in the main tunnel about 100 feet in has been followed in an inclined winze for about 95 feet to a point where it appears to be cut off by a rather obscure fault. There is apparently no large quantity of ore available. The Sisson shaft appears to be entirely in diabase below the first level.

The ore of the Rattler mine is chiefly a dark fine-grained aggregate of magnetite and chalcopyrite with varying quantities of silicate minerals derived by metamorphism from the inclosing limestone. With increasing proportions of these silicates the ore grades into the metamorphosed limestone. Analyses of the ore recorded in the books of the Troy Arizona Copper Co. and its predecessors show from 3 to 3.7 per cent of copper, a maximum of 0.04 ounce of gold and 0.7 ounce of silver per ton, from 27 to 30 per cent of silica, about the same proportion of iron, about 1 per cent of calcium oxide, and 20 per cent of magnesium oxide.

Very little could be seen of the '91 mine, as the shaft had caved in. It apparently is about 150 feet deep, and the maps show three short levels. The little copper ore that was found in this mine appears to have occurred, as at the Rattler, as small lenticular bunches in the Mescal limestone. The wulfenite in the '91 mine, concentration of which was at one time attempted, occurs in joints in fractured Dripping Spring quartzite.

The Buckeye mine is situated on a nearly east and west branch of the same porphyry dike complex on which is the Alice mine, at a point where the dike complex cuts through Dripping Spring quartzite, Mescal limestone, and Troy quartzite. These stratified rocks appear to be underlain and overlain by intrusive sheets of diabase. The dump of the shaft, which is apparently about 150 feet deep, with three levels, is chiefly diabase. These levels trend generally west-northwest. The first level has a length of about 1,000 feet, but each of the other two is less than 300 feet. The Buckeye at one time had a little oxidized copper ore near the surface and alongside of the porphyry dike. This ore was reduced in a small furnace at the mine.

The Alice mine was worked through a shaft inclined at 45° with three levels, the lowest of which is about 200 feet vertically below the collar. As shown by the mine map the general trend of the levels is northeast and the length of the block of ground explored by them about 350 feet. The shaft is sunk on a porphyry dike at a point where Tornado limestone on the north is faulted down against Martin limestone on the south. A few small bunches of ore were found in limestone near the dike, but the mine did not pay expenses. It could not be entered when the geologic field work on which this folio is

about 1/4 mi.
north of
the Buckeye
mine?

based was in progress. The mine maps show that connection was made with the Pratt tunnel by an inclined raise of about 200 feet vertical height.

Since the last visit the Troy Arizona Copper Co. has done additional prospecting and is reported to have sunk a 500-foot shaft on the Climax claim, about three-quarters of a mile west-southwest of Troy, and in 1917 some ore was being shipped.

On the Renfro group of 47 claims, about 1½ miles east-southeast of Troy, considerable ore was visible in 1912 in the lower part of the Martin limestone at a point near the crest of a steep spur where the limestone beds, which dip at a low angle to the east, are stepped down toward the south by four or five small faults the throw of which is apparently nowhere over 40 to 50 feet. The ore, mostly chrysocolla and carbonates, occurs as irregular layers 6 inches in maximum thickness, which lie parallel with the bedding of the limestone and occur as small replacement masses near fissures. From the ravine to the west of these exposures of ore on the ridge, and about 400 feet below them, a tunnel 900 feet in length had been driven in 1912 entirely in diabase, which here underlies the Troy quartzite and Martin limestone of the crest of the ridge. The tunnel follows a nearly vertical fissure and runs southeast. No ore had been found in this tunnel in 1912. Since that year the ground has been worked by the Pinal Development Co., which began production in 1917. The ore shipped is oxidized and presumably comes from the replacement deposits in limestone previously mentioned. The main tunnel is stated⁷¹ to be 1,600 feet long and to connect with about 2,500 feet of underground workings with a maximum depth of 600 feet.

OTHER COPPER DEPOSITS.

London-Arizona mine.—The London-Arizona mine is in the southeastern part of the quadrangle, about 4 miles north of Hayden, on the north side of Tornado Peak.

The lowest rock exposed in the canyon, in which are the mine buildings, is diabase, apparently in a sheet several hundred feet thick, which was intruded at approximately the horizon of the Mescal limestone. Overlying the diabase in succession are the Troy quartzite, the Martin limestone, and the Tornado limestone. All these rocks are cut by dikes and small intrusive masses of quartz diorite porphyry. The diabase in the vicinity of the mine buildings is conspicuously metamorphosed by the porphyry and in places is a sparkling dark biotitic schist, generally containing disseminated pyrite and chalcopyrite.

On the south side of the ravine the lower part of the Devonian Martin limestone, as at other places in the quadrangle, shows metallization by copper, especially near dikes of quartz diorite porphyry, and since the time of visit considerable oxidized copper ore has been shipped to the Hayden smelter from flat-lying lenticular deposits in this limestone. In 1913 about 1,000 tons of 16 per cent ore was sent to the smelter and in 1916 about 6,000 tons of 4½ per cent ore. This ore was probably mined through tunnels or inclines on the south side of the ravine. A few thousand tons of lead ore was also mined. In 1910 exploration was in progress from the Curtin shaft, which was sunk in the Tornado limestone south of the ravine in order to reach the ore-bearing zone in the Martin limestone or to cut any ore bodies that might possibly occur at higher horizons in the limestones near one of the porphyry dikes. The shaft at that time was 270 feet deep, and no ore had been found in the workings connected with it.

Christmas deposits.—The interesting contact-metamorphic deposits at Christmas, although they lie in the adjoining Christmas quadrangle, about 4 miles east of the London-

Average of eight analyses of

Cu	-----
S	-----
Fe	-----
CaO	-----
SiO ₂	-----

Schneider group.—The and west of the ground was in progress on the 1910, but operations by Canon Consolidated Co.

The principal development the No. 1 tunnel, on the ravine. This tunnel runs to the east, and at that It is chiefly in diabase porphyry. Small irregular and quartz were observed of ore had been found a

Since then considerable 1917 the main tunnel v ore were shipped weekly of about \$200,000 to the

Kelvin-Sultana mine. the Kelvin-Sultana Co. River, nearly opposite feet deep, and the total mately 10,000 feet in le

The shaft is sunk in intrusive into pre-Cambrian quartz diorite porphyry that trend nearly east a tical or dip south at hi oxidized copper ore no ments, amounting to al these zones on the Wi workings, now abandon shaft is about 800 feet i

At the time of the fir the bottom level of the to the south perhaps 50 porphyry dike and thr zones but showed no ore

At the second visit, i although it was being k 1,000 gallons a day. and a wire-rope tramwa A power plant had als river.

The mine has since not been in continuous

GOLD A

Pioneer mines.—In the south base of Pione from Kelvin to Globe underground work don settlement of Pioneer. been the Republic, bu shaft was started in di below, and the ore app

Fragments from the and barite, but rich se have occurred near th narrow stopes, 3 to 4 f the existence of a bra

Rain Folio #217

AREAL GEOLOGY

